Correction to: Identifying Nonalcoholic Fatty Liver Disease Advanced Fibrosis in the Veterans Health Administration

Yuval A. Patel¹ · Elizabeth J. Gifford² · Lisa M. Glass^{3,4} · Marsha J. Turner⁵ · Byungjoo Han⁵ · Cynthia A. Moylan¹ · Steve Choi¹ · Ayako Suzuki¹ · Dawn Provenzale^{1,5} · Christine M. Hunt^{1,5}

Published online: 24 July 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

Correction to: Digestive Diseases and Sciences
<https://doi.org/10.1007/s10620-018-5123-3>

The original version of the article unfortunately contained errors in Table 3, Risk Factor column headings “Age > 50 ($n = 115$),” “Age > 50–64 ($n = 154$),” and “Age > 65 + ($n = 60$).” Corrected version of Table 3 is given below.

The original article can be found online at <https://doi.org/10.1007/s10620-018-5123-3>.

✉ Yuval A. Patel
Yuval.Patel@duke.edu

Elizabeth J. Gifford
Beth.Gifford@duke.edu

Lisa M. Glass
LisaGlas@med.umich.edu

Marsha J. Turner
Marsha.Turner2@va.gov

Byungjoo Han
Byungjoo.Han@va.gov

Cynthia A. Moylan
Cynthia.Moylan@duke.edu

Steve Choi
Steve.Choi@duke.edu

Ayako Suzuki
Ayako.Suzuki@duke.edu

Dawn Provenzale
Dawn.Provenzale@duke.edu

Christine M. Hunt
Christine.Hunt@va.gov

- ¹ Department of Medicine, Duke University Medical Center, Durham Veterans Administration Medical Center, Box 3913, Durham, NC 27710, USA
- ² Sanford School of Public Policy, Duke University, Durham, NC, USA
- ³ Department of Medicine, University of Michigan, Ann Arbor, MI, USA
- ⁴ VA Ann Arbor Healthcare Systems, Ann Arbor, MI, USA
- ⁵ Cooperative Studies Program Epidemiology Center, Durham Veterans Administration Medical Center, Durham, NC, USA

Table 3 Fibrosis prediction scores by age (using alternative thresholds for age ≥ 65)

Risk factor	AUROC 95% CI	ROC lower bound	ROC upper bound	Cutoff	Sensitivity	Specificity	PPV	NPV	LR +	LR –	Odds ratio	Prevalence
<i>Age < 50 (n = 115)</i>												
NAFLD fibrosis > 0.676	0.72	0.56	0.87	0.676	45.45	97.94	71.43	94.06	22.05	0.56	39.58	10.19
NAFLD fibrosis < – 1.455	0.22	0.12	0.32	– 1.455	9.09	35.05	1.56	77.27	0.14	2.59	0.5	10.19
FIB-4 > 2.67	0.62	0.49	0.74	2.67	25	98.04	60	91.74	12.75	0.76	16.67	10.53
FIB-4 < 1.3	0.25	0.10	0.39	1.3	33.33	15.69	4.44	66.67	0.40	4.25	0.09	10.53
BARD 2–4	0.74	0.62	0.86	2	83.33	65.05	21.74	97.10	2.38	0.26	9.31	10.43
BARD 0–1	0.26	0.14	0.38	1	16.67	34.95	2.90	78.26	0.26	2.38	0.11	10.43
<i>Age 50–64 (n = 154)</i>												
NAFLD fibrosis > 0.676	0.76	0.68	0.83	0.676	68	83.51	68	83.51	4.12	0.38	10.76	34.01
NAFLD fibrosis < – 1.455	0.32	0.27	0.37	– 1.455	2	61.9	2.63	55.05	1.58	0.04	0.03	34.01
FIB-4 > 2.67	0.76	0.69	0.83	2.67	54.90	98	93.33	80.99	27.45	0.46	59.65	33.77
FIB-4 < 1.3	0.25	0.19	0.32	1.3	7.84	43	6.56	47.78	0.14	2.14	0.06	33.77
BARD 2–4	0.67	0.62	0.73	2	96.08	38.61	44.14	95.12	1.57	0.10	15.41	33.55
BARD 0–1	0.33	0.27	0.38	1	3.92	61.39	4.89	55.86	0.10	1.57	0.06	33.55
<i>Age ≥ 65 (n = 60)</i>												
NAFLD fibrosis > 0.676	0.71	0.59	0.83	0.676	74.07	67.74	66.67	75	2.30	0.38	6	46.55
NAFLD fibrosis < – 0.12	0.33	0.22	0.44	– 0.12	14.81	51.61	21.05	41.03	0.31	1.65	0.19	46.55
NAFLD fibrosis < – 1.455	0.43	0.36	0.51	– 1.455	3.70	83.87	16.67	50	0.23	1.15	0.2	46.55
FIB-4 > 3.25	0.80	0.70	0.89	3.25	65.52	93.55	90.48	74.36	10.16	0.37	27.55	48.33
FIB-4 > 2.67	0.80	0.70	0.90	2.67	75.86	83.87	81.48	78.79	4.70	0.29	16.34	48.33
FIB-4 < 2.0 if age ≥ 65	0.33	0.22	0.44	2	13.79	51.61	21.05	39.02	0.29	1.67	0.17	48.33
FIB-4 < 1.3	0.49	0.42	0.56	1.3	6.90	90.32	40	50.91	0.71	1.03	0.69	48.33
BARD 2–4	0.63	0.54	0.72	2	96.55	29.03	56	90	1.36	0.12	11.45	48.33
BARD 0–1	0.37	0.28	0.46	1	3.45	70.97	10	44	0.12	1.36	0.09	48.33