CORRECTION



Correction to: Hepatitis B Surface Antigen Loss with Tenofovir Disoproxil Fumarate Plus Peginterferon Alfa-2a: Week 120 Analysis

Sang Hoon Ahn¹ · Patrick Marcellin² · Xiaoli Ma³ · Florin A. Caruntu⁴ · Won Young Tak⁵ · Magdy Elkhashab⁶ · Wan-Long Chuang⁷ · Fehmi Tabak⁸ · Rajiv Mehta⁹ · Jörg Petersen¹⁰ · William Guyer¹¹ · Belinda Jump¹¹ · Alain Chan¹¹ · Mani Subramanian¹¹ · Gerald Crans¹¹ · Scott Fung¹² · Maria Buti¹³ · Giovanni B. Gaeta¹⁴ · Aric J. Hui^{15,16} · George Papatheodoridis¹⁷ · Robert Flisiak¹⁸ · Henry L. Y. Chan¹⁹

Published online: 15 October 2018

© The Author(s) 2018

Correction to: Digestive Diseases and Sciences https://doi.org/10.1007/s10620-018-5251-9

The original version of this article unfortunately contained affiliation and textual errors. This has been corrected with this erratum.

Affiliation of eleventh author was incorrectly assigned. Dr. William Guyer is affiliated with Gilead Sciences Inc, Foster City, CA, USA.

In Results section, Safety Profile subsection, the sentence, 'Rates of serious adverse events were 11%, 10%, 7%, and

The original article can be found online at https://doi.org/10.1007/s10620-018-5251-9.

Patrick Marcellin patrick.marcellin@bjn.aphp.fr

Henry L. Y. Chan hlychan@cuhk.edu.hk

Sang Hoon Ahn ahnsh@yuhs.ac

Xiaoli Ma tangmali@yahoo.com

Florin A. Caruntu drcaruntu@mateibals.ro

Won Young Tak wytak@knu.ac.kr

Magdy Elkhashab melkashabmd@yahoo.ca

Wan-Long Chuang waloch@kmu.edu.tw

Fehmi Tabak fehmitabak@hotmail.com

Rajiv Mehta rmgastro@yahoo.com 10% in groups A, B, C, and D, respectively, and 6%, 3%, and 5% in groups A, B, and D, respectively, among those patients who were treated with TDF' should read as below:

Rates of serious adverse events were 11%, 10%, 7%, and 10% in groups A, B, C, and D, respectively, and 6%, 3%, and 5% in groups A, B, and D, respectively, among those patients who were retreated with TDF.

In Table 2, header 'Mean HBV DNA change from baseline, $\log_{10} IU/mL$ (SD)' row 3 is missing 'B' in genotype. Revised version of the Table 2 is given below.

Jörg Petersen petersen@ifi-medizin.de

William Guyer william.guyer@gilead.com

Belinda Jump belinda.jump@gilead.com

Alain Chan alain.chan@gilead.com

Mani Subramanian mani.subramanian@gilead.com

Gerald Crans gerald.crans@gilead.com

Scott Fung scott.fung@uhn.on.ca

Maria Buti mbuti@vhebron.net

Giovanni B. Gaeta giovannibattista@unicampania.it

Aric J. Hui arichui@yahoo.com



Table 2 Efficacy results at week 120 by baseline HBeAg status and genotype

Response	Group A (n=186) TDF+PEG-IFN for 48 weeks	Group B (n=184) TDF+PEG-IFN for 16 weeks, TDF for 32 weeks	Group C (n=185) TDF for 120 weeks	Group D (n=185) PEG-IFN for 48 weeks
HBsAg loss, Kaplan	n–Meier estimate (%)			
Overall	10.36	3.49	0	3.51
HBeAg positive	9.73	5.15	0	5.23
HBeAg negative	10.99	1.32	0	1.28
HBsAg seroconvers	sion, Kaplan-Meier est	imate (%)		
Overall	10.08	0.56	0	2.87
HBeAg positive	10.41	0.97	0	4.10
HBeAg negative	9.65	0	0	1.28
Mean HBsAg chang	ge from baseline, \log_{10}	IU/mL (SD)		
Overall	-2.4(2.4)	-0.8(1.5)	-0.4(0.7)	-1.1 (1.6)
Genotype A	-4.2(2.3)	_	-0.7(1.2)	0.0 (0.6)
Genotype B	-2.2(2.1)	-1.0(1.0)	-0.7(0.6)	-1.2 (1.2)
Genotype C	-1.5 (2.2)	-0.4(1.5)	-0.4(0.7)	-1.1 (1.7)
Genotype D	-2.0(2.3)	-1.6(2.6)	-0.2(0.4)	-1.5 (2.9)
HBeAg positive	-2.2(2.6)	-1.2(1.9)	-0.7(0.7)	-1.3 (2.1)
HBeAg negative	-2.5(2.1)	-0.4(0.9)	-0.1(0.4)	-1.0(1.1)
Mean HBV DNA cl	hange from baseline, lo	og ₁₀ IU/mL (SD)		
Overall	-4.2(2.1)	-4.3(2.0)	-5.7 (1.6)	-3.5(2.0)
Genotype A	-4.0(1.9)	_	-5.1 (2.1)	-1.9 (2.0)
Genotype B	-4.7(2.0)	-4.2(2.5)	-5.8 (1.4)	-3.3 (1.8)
Genotype C	-4.2(2.9)	-4.4 (1.5)	-5.9 (1.6)	-3.4(2.3)
Genotype D	-4.0(1.8)	-4.1(3.0)	-5.3 (1.7)	-5.1 (1.9)
HBeAg positive	-5.5 (1.6)	-5.3 (1.5)	-6.6 (1.2)	-4.9 (1.6)
HBeAg negative	-2.9(1.7)	-3.4 (1.9)	-4.5 (1.3)	-2.5(1.8)

George Papatheodoridis gepapath@med.uoa.gr

Robert Flisiak robert.flisiak@umwb.edu.pl

- Department of Internal Medicine, Yonsei University College of Medicine, Brain Korea 21 Plus Project for Medical Science, Seoul, Republic of Korea
- ² Hôpital Beaujon, University Paris-Diderot, Clichy, France
- Drexel University College of Medicine, Philadelphia, PA, USA
- ⁴ National Institute for Infectious Diseases "Matei Bals", Bucharest, Romania
- South Korea 5 Kyungpook National University Hospital, Daegu, South Korea 5 Kyungpook National University Hospital, Daegu, South Korea 6 Kyungpook National University Hospital, Daegu, South Korea 7 Kyungpook National University Hospital, Daegu, South National University Hospital, Daegu, Na
- Toronto Liver Centre, Toronto, Canada
- Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan
- ⁸ Cerrahpasa Medical Faculty, University of Istanbul, Istanbul, Turkey
- ⁹ Liver Clinic, Surat, India

- IFI Institute for Interdisciplinary Medicine, Asklepios Klinik St. George, University of Hamburg, Hamburg, Germany
- ¹¹ Gilead Sciences Inc, Foster City, CA, USA
- Toronto General Hospital, Toronto, Canada
- Hepatology Unit, Hospital Universitari Vall d'Hebron and CIBEREHD del Instituto Carlos III, Barcelona, Spain
- Infectious Diseases and Viral Hepatitis Unit, University of Campania "Luigi Vanvitelli", Naples, Italy
- ¹⁵ The Chinese University of Hong Kong, Hong Kong, China
- Alice Ho Miu Ling Nethersole Hospital, Hong Kong, China
- Medical School of National and Kapodistrian University of Athens, General Hospital of Athens "Laiko", Athens, Greece
- Department of Infectious Diseases and Hepatology, Medical University of Bialystok, Białystok, Poland
- Department of Medicine and Therapeutics and Institute of Digestive Disease, The Chinese University of Hong Kong, Hong Kong, China

