Correction to: Patients with Alcoholic Liver Disease Have Worse Functional Status at Time of Liver Transplant Registration and Greater Waitlist and Post-transplant Mortality Which Is Compounded by Older Age

Patrick McCabe¹ · Artin Galoosian² · Robert J. Wong³

Published online: 3 January 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Digestive Diseases and Sciences
<https://doi.org/10.1007/s10620-019-05891-1>

The original version of the article unfortunately contained an error in Figure 3. X-axis of Figures 3a–d should be ‘Time in months’. Corrected version of Fig. 3 is given below.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10620-019-05891-1>.

✉ Robert J. Wong
Rowong@alamedahealthsystem.org
Patrick McCabe
McCabP@sutterhealth.org

- ¹ Division of Gastroenterology and Hepatology, Department of Medicine, California Pacific Medical Center, San Francisco, CA, USA
- ² Department of Medicine, California Pacific Medical Center, San Francisco, CA, USA
- ³ Division of Gastroenterology and Hepatology, Alameda Health System – Highland Hospital, 1411 East 31st Street, Highland Hospital – Highland, Care Pavilion 5th Floor, Endoscopy Unit, Oakland, CA 94602, USA

Fig. 3 Kaplan–Meier curves for 5-year post-LT survival stratified by liver disease etiology and KPS