Correction to: Cost-Effectiveness of Empirical Bismuth-Based Quadruple Therapy and Tailored Therapy After Clarithromycin Resistance Tests for *Helicobacter pylori* Eradication

Young Woon Chang¹ · Ga Young Shin² · Jung-Wook Kim² · Jin-Chang Moon¹ · Eun Jee Chang³ · Chi Hyuk Oh² · Jae-Young Jang²

Published online: 20 May 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Digestive Diseases and Sciences
<https://doi.org/10.1007/s10620-021-06938-y>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original version of the article unfortunately contained an error in the Abstract. In the Abstract, under 'Methods', the sentence 'The tailored group (n = 282) consisted of patients treated with STT for 7 days and BQT for 10 days as per their CR test results.' should read as 'The tailored group (n = 292) consisted of patients treated with STT for 7 days and BQT for 10 days as per their CR test results.'

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s10620-021-06938-y>.

✉ Jung-Wook Kim
iloveact@hanmail.net

¹ Department of Internal Medicine, Cheonan-Woori Hospital, Cheonan, Korea

² Department of Gastroenterology, Kyung Hee University College of Medicine, 23 Kyung Hee Dae-ro, Dongdaemun-gu, Seoul 02447, Korea

³ Department of Preventive Medicine, Graduate School of Public Health, Seoul National University, Seoul, Korea