Correction to: Increased Fecal Bile Acid Excretion in a Significant Subset of Patients with Other Inflammatory Diarrheal Diseases

Priya Vijayvargiya¹ · Daniel Gonzalez Izundegui¹ · Gerardo Calderon¹ · Sarah Tawfic² · Sarah Batbold² · Hiba Saifuddin² · Patrick Duggan² · Valeria Melo² · Taylor Thomas² · Megan Heeney² · Adrian Beyde² · James Miller Jr.² · Kenneth Valles² · Kafayat Oyemade² · Joseph F. Brant² · Jessica Atieh¹ · Leslie J. Donato³ · Michael Camilleri¹

Published online: 16 October 2021

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Digestive Diseases and Sciences
<https://doi.org/10.1007/s10620-021-06993-5>

In Table 3 of this article, the data ‘20.4 (4.8–81)’ in the column ‘Ileal resection’ and row ‘% primary BA’ should have read as ‘95 (75–99)’.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10620-021-06993-5>.

✉ Michael Camilleri
camilleri.michael@mayo.edu

¹ Clinical Enteric Neuroscience Translational and Epidemiological Research (CENTER), Division of Gastroenterology and Hepatology, Mayo Clinic, 200 First St. S.W. Charlton Bldg., Rm. 8-110, Rochester, MN 55905, USA

² Mayo Medical School, Rochester, MN, USA

³ Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN 55905, USA