

## Erratum to: Condition factor, Length – Weight relationship, and the fishery of *Barbus altianalis* (Boulenger 1900) in Lakes Victoria and Edward basins of Uganda

C. C. Ondhoro · C. Masembe · G. E. Maes ·  
N. W. Nkalubo · J. K. Walakira · J. Naluwairo ·  
M. T. Mwanja · J. Efitre

Published online: 12 January 2017  
© Springer Science+Business Media Dordrecht 2017

### Erratum to: Environ Biol Fish DOI:10.1007/s10641-016-0540-7

The original version of this article was revised to standardize units of measurement. The changes are underlined as follows:

#### 1. 9th sentence under **Abstract**

Catch rates varied between River Nile ( $1.92 \pm 0.59$   $\text{Kg boat}^{-1} \text{day}^{-1}$ ) and the rest of the systems,  $6.20 \pm 1.86$  and  $6.85 \pm 1.49$   $\text{Kg boat}^{-1} \text{day}^{-1}$  in Lake Edward and Kazinga channel respectively.

#### 2. 11th sentence under **Abstract**

Dissolved oxygen was below the minimum of 5  $\text{mg l}^{-1}$  required for the physiology of freshwater fish.

#### 3. 12th sentence under **Abstract**

Conductivity was highest in Lake Edward ( $312$   $\mu\text{S cm}^{-1}$ ), followed by Kazinga channel and least in River Nile.

#### 4. 5th sentence, paragraph 2 under **Introduction**

Catch rates in the Lakes Victoria and Edward basins reduced to the lowest levels of 0.5 to 0.2  $\text{kg h}^{-1}$  (Bekkevold et al. 2005; Chande and Mhithu 2005; Ntakimazi 2006) in the 1970s, and further declined to 0.06  $\text{kg h}^{-1}$  in Lake Victoria (Chande and Mhithu 2005).

#### 5. 1st sentence, paragraph 2 under **Water quality data collection**

Dissolved oxygen ( $\text{mg l}^{-1}$ ), water temperature ( $^{\circ}\text{C}$ ), electrical conductivity ( $\mu\text{S cm}^{-1}$ ) and pH were determined using Yellow Spring Instrument (YSI, model 556) at

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10641-016-0540-7>.

---

C. C. Ondhoro (✉)  
Buginyanya Zonal Agricultural Research and Development,  
Institute, P.O. Box 1356, Mbale, Uganda  
e-mail: oconstantine88@gmail.com

C. C. Ondhoro · J. K. Walakira · M. T. Mwanja  
Aquaculture Research and Development Centre Kajjansi,  
P.O. Box 530, Kampala, Uganda

C. C. Ondhoro · C. Masembe · J. Efitre  
Department of Zoology, Entomology and Fisheries Sciences,  
Makerere University, P.O. Box 7062, Kampala, Uganda

G. E. Maes  
Laboratory of Biodiversity and Evolutionary Genomics,  
University of Leuven (KU Leuven), B-3000 Leuven, Belgium

G. E. Maes  
Centre for Sustainable Tropical Fisheries and Aquaculture,  
Comparative Genomics Centre, College of Science and  
Engineering, James Cook University, Townsville, Qld 4811,  
Australia

N. W. Nkalubo · J. Naluwairo  
National Fisheries Resources Research Institute, P.O. Box 343,  
Jinja, Uganda

mid-water and between 07:00 h to 08:30 h to generate monthly average water quality observations for each population.

6. 5th sentence, paragraph 4 under **Extinction upstream Owen falls dam and variation in Condition between populations** section

Whereas dissolved oxygen levels in River Nile was higher than in Lake Edward and Kazinga channel, the concentration recorded is still lower than the usual recommended minimum of  $\geq 5 \text{ mg l}^{-1}$ . Boyd (1998)

provides water quality benchmarks for fresh water fishes within the range of 5–15  $\text{mg l}^{-1}$ .

7. 7th sentence, paragraph 4 under **Extinction upstream Owen falls dam and variation in Condition between populations** section

Elsewhere,  $8.7 \text{ mg l}^{-1}$  (Abowei 2010); Arumlampalam et al. (1998), documented dissolved oxygen levels of  $5.54\text{--}7.98 \text{ mg l}^{-1}$  in fresh water fisheries of the upper Gulf of Thailand and Malaysia respectively.

The original article was corrected.