Correction to: Japanese catadromous fourspine sculpin, *Rheopresbe kazika* (Jordan & Starks) (Pisces: Cottidae), transferred from the genus *Cottus*

Akira Goto · Ryota Yokoyama · Izumi Kinoshita ·
Harumi Sakai

Published online: 17 January 2020
© Springer Nature B.V. 2020

Correction to: Environ Biol Fish
<https://doi.org/10.1007/s10641-019-00921-3>

The original version of this article unfortunately contained mistakes introduced during the production process. Tables 1 and 2 were presented incorrectly. The correct Tables are now given below:

The online version of the original article can be found at
<https://doi.org/10.1007/s10641-019-00921-3>

A. Goto (✉)
Department of Science of Environmental Education, Hokkaido
University of Education, Hachiman-cho 1-2, Hakodate, Hokkaido
040-8567, Japan
e-mail: akir@fish.hokudai.ac.jp

R. Yokoyama
Civil Engineering & Eco-Technology Consultants, Co., Ltd,
2-23-2 Higashi-Ikebukuro, Tokyo, Toshima 170-0013, Japan

I. Kinoshita
USA Institute of Marine Biology, Kochi University, 194, Tosa,
Kochi 781-1164, Japan

H. Sakai
Department of Applied Aquabiology, National Fisheries
University, 2-7-1 Nagata-honmachi, Shimonoseki, Yamaguchi
759-6595, Japan

Table 1 Comparison of morphological characteristics in adult and larvae to juvenile stages among the fourspine sculpin *Rheopresbe kazika*, *Cottus* species and *Trachidermus fasciatus*

	Adult stage			Larval to juvenile stages		
	Bony r. l. ¹⁾	Villiform teeth ²⁾	Preop. sp. ³⁾	Pariet. sp. ⁴⁾	Body depth	Melanoph. on p. f. ⁵⁾
<i>Rheopresbe kazika</i>	absent	present	4	present	markedly deep	densely distribution
<i>Cottus</i> species	absent	absent or rarely present	1 to 3	absent or rarely present	shallow	no distribution
<i>Trachidermus fasciatus</i>	present	present	4	present	shallow	densely distribution

1) Bony ridge lines, 2) Villiform teeth on the palatines, 3) No. of Preopercular spines, 4) Parietal spines and 5) Melanophores on the pectoral fins. Data on adult morphology cited from Jordan and Starks (1904), Watanabe (1960), Nakamura (1963), Yabe (1985) and Fujii (2001). Data on larval and Juvenile morphology cited from Goto (1977), Takeshita et al. (1997), Harada et al. (1999), Kinoshita et al. (1999) and Kojima (2014)

Table 2 Karyotypes of *Rheopresbe kazika*, *Trachidermus fasciatus* and *Cottus* species

Species	Biarmed chromosomes (M and SM)	Uniarmed chromosomes (A)	2N	NF
<i>Rheopresbe kazika</i> ¹⁾	18	22	40	58
<i>Trachidermus fasciatus</i> ¹⁾	24	16	40	64
<i>Cottus gobio</i> ³⁾	6	46	52	58
<i>C. hangiongensis</i> ²⁾	6	42	48	54
<i>C. nozawae</i> ²⁾	10	38	48	58
<i>C. poecilopus</i> ³⁾	8	40	48	56
<i>C. pollux</i> ¹⁾	10	38	48	58
<i>C. reini</i> ¹⁾	12	36	48	60

M: metacentrics, SM: submetacentrics, A: acrocentrics, 2N: number of chromosomes, NF: Total number of chromosome. Arm number cited, 1) from Abe (1972), 2) from Abe (1976), 3) from Starmach (1967)

The original article has been corrected.