Correction to: Spatio-temporal distribution of Green Sturgeon (*Acipenser medirostris*) and White Sturgeon (*A. transmontanus*) in the San Francisco Estuary and Sacramento River, California

Emily A. Miller · Gabriel P. Singer · Matthew L. Peterson · Eric D. Chapman · Myfanwy E. Johnston · Michael J. Thomas · Ryan D. Battleson · Marty Gingras · A. Peter Klimley

Published online: 14 July 2020
© Springer Nature B.V. 2020

Correction to: Environ Biol Fish 103(5):577–603
<https://doi.org/10.1007/s10641-020-00972-x>

Due to an oversight by the Publisher during the typesetting stage, the Fig. 8 has been omitted from the pdf version and is now presented in this article.

The online version of the original article can be found at
<https://doi.org/10.1007/s10641-020-00972-x>

E. A. Miller (✉)
Monterey Bay Aquarium, 886 Cannery Row, Monterey, CA
93940, USA
e-mail: eamiller@ucdavis.edu

E. A. Miller · G. P. Singer · M. L. Peterson ·
E. D. Chapman · M. E. Johnston · M. J. Thomas ·
R. D. Battleson · M. Gingras · A. P. Klimley
University of California, Davis, Wildlife, Fish, and Conservation
Biology Department, One Shields Ave., Davis, CA 95616, USA

M. L. Peterson
FISHBIO, 180 East 4th Street, Suite 160, Chico, CA 95928, USA

E. D. Chapman
ICF, 980 9th Street, Suite 1200, Sacramento, CA 95814, USA

M. E. Johnston
Cramer Fish Science, 1300 New Airport Road, Suite 103, Auburn,
CA 95602, USA

R. D. Battleson
Oregon Department of Fish and Wildlife, Salmon and Trout
Enhancement Program, 1495 E Gregory Road, Central Point, OR
97502, USA

M. Gingras
California Department of Fish and Wildlife, 2109 Arch Airport
Rd, Stockton, CA 95206, USA

Fig. 8 Median number of days spent on spawning grounds for adult Green (above rkm 411.8 at Irvine Finch, reach 9 in Fig. 1) and White Sturgeon (above rkm 239 at Verona, reach 8 in Fig. 1)

The Publisher apologizes for this error.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.