



# Correction to: Metabolic growth hypothesis for the evolution of the nuchal hump in swordtail fishes

Keith B. Tompkins · Meggan S. Lott · Oscar Rios-Cardenas · Sukanta Jash · Molly R. Morris

Published online: 10 March 2022  
© Springer Nature B.V. 2022

**Correction to: Environ Biol Fish (2021) 104(10):1195–1206**  
<https://doi.org/10.1007/s10641-021-01145-0>

The original version of this article unfortunately contained mistakes.

The corrections are given in the following list:

- (1) In the abstract, sentence four reads: “Factors that influenced the size of the nuchal hump included diet and genotype, with genotypes known to have faster juvenile growth rates forming larger humps, as males stop growing after sexual maturity, forming larger humps.” The sentence should read: “Factors that influenced the size of the nuchal hump included diet and genotype, with genotypes known to have faster juvenile growth rates forming larger humps.”
- (2) The graph listed in Fig. 5 was actually a duplicate of the graph listed as Fig. 3.


The online version of the original article can be found at <https://doi.org/10.1007/s10641-021-01145-0>.

K. B. Tompkins (✉) · M. S. Lott · M. R. Morris  
Department of Biological Sciences, Ohio University,  
Athens, OH, USA  
e-mail: kt414318@ohio.edu

O. Rios-Cardenas  
Instituto de Ecología A.C, Xalapa, Mexico

S. Jash  
Alpert Medical School, Brown University, Providence, RI,  
USA

The correct Fig. 5 is now given below.



**Fig. 5** Activity levels (number of crosses in test tank) in the cold-water treatment was positively related to the relative size of the nuchal hump (hump area/body area)

The original article has been corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.