



Publisher Correction: Impact of uncertainties in exposure assessment on thyroid cancer risk among cleanup workers in Ukraine exposed due to the Chernobyl accident

Mark P. Little^{1,5} · Elizabeth K. Cahoon¹ · Natalia Gudzenko² · Kiyohiko Mabuchi¹ · Vladimir Drozdovitch¹ · Maureen Hatch¹ · Alina V. Brenner³ · Vibha Vij¹ · Konstantin Chizhov¹ · Elena Bakhanova² · Natalia Trotsyuk² · Victor Kryuchkov⁴ · Ivan Golovanov⁴ · Vadim Chumak² · Dimitry Bazyka²

Published online: 20 August 2022

This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2022

Publisher Correction to: European Journal of Epidemiology
<https://doi.org/10.1007/s10654-022-00850-z>

The correct table is included in this correction and the paper has been updated.

The Publisher regrets that after publication errors were found in Table 2, with columns 2, 3, and 5 formatted incorrectly.

The original article can be found online at <https://doi.org/10.1007/s10654-022-00850-z>.

✉ Mark P. Little
mark.little@nih.gov

¹ Radiation Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA

² National Research Centre for Radiation Medicine, Kyiv 04050, Ukraine

³ Radiation Research Effects Foundation, Hiroshima, Japan

⁴ Burnasyan Federal Medical and Biophysical Centre, 46 Zhivopisnaya Street, Moscow, Russia 123182

⁵ Radiation Epidemiology Branch, Division of Cancer Epidemiology and Genetics, Department of Health and Human Services, National Cancer Institute, National Institutes of Health, 9609 Medical Center Drive, Bethesda, MD 20892-9778, USA

Table 2 Excess odds ratio/Gy by morphology subtypes using linear EOR model, using geometric mean doses, regression calibration adjusted doses, or Monte Carlo maximum likelihood

Regression adjustment	Follicular morphology			Papillary morphology			Other and unspecified thyroid cancer		
	Cases/controls	EOR (95% CI)	p-value	Cases/controls	EOR (95% CI)	p-value	Cases/controls	EOR (95% CI)	p-value
Unadjusted model									
Regression (geometric mean of 10,000 doses)	24/72	2.560 (-0.269, 21.458)	0.138	110/338	0.115 (-0.117, 1.104)	0.580	15/48	2.353 (-0.084, > 100)	0.074
Regression calibration (arithmetic mean of 10,000 doses)	24/72	1.826 (-0.227, > 100)	0.137	110/338	0.067 (-0.101, > 100)	0.671	15/48	2.003 (-0.073, > 100)	0.075
Full Monte Carlo maximum likelihood	24/72	2.072 (-0.247, > 100)	0.153	110/338	0.075 (-0.172, 0.988)	0.659	15/48	2.314 (-0.073, 17.476)	0.074
Adjusted baseline for proxy status									
Regression (geometric mean of 10,000 doses)	24/72	2.699 (-0.262, 24.083)	0.132	110/338	0.176 (-0.114, 1.428)	0.470	15/48	2.016 (-0.141, > 100)	0.100
Regression calibration (arithmetic mean of 10,000 doses)	24/72	1.733 (-0.245, 13.628)	0.152	110/338	0.109 (-0.099, 0.991)	0.552	15/48	1.670 (-0.131, 13.369)	0.108
Full Monte Carlo maximum likelihood	24/72	2.009 (-0.259, 23.456)	0.165	110/338	0.125 (-0.156, 1.298)	0.537	15/48	1.996 (-0.133, > 100)	0.104
Adjusted baseline for smoking status, cigarette packyears and proxy status ^a									
Regression (geometric mean of 10,000 doses)	24/72	5.461 (-0.015, 60.624)	0.052	110/338	0.212 (-0.112, 1.562)	0.416	15/48	2.025 (-0.174, 18.789)	0.117
Regression calibration (arithmetic mean of 10,000 doses)	24/72	3.224 (-0.082, 30.615)	0.068	110/338	0.131 (-0.098, 1.064)	0.499	15/48	1.653 (-0.155, 15.502)	0.126
Full Monte Carlo maximum likelihood	24/72	4.708 (-0.075, 85.143)	0.066	110/338	0.143 (-0.149, 1.338)	0.493	15/48	1.970 (-0.165, 21.697)	0.125
Adjusted baseline for smoking status, cigarette packyears, alcohol consumption and proxy status ^b									
Regression (geometric mean of 10,000 doses)	24/72	30.091 (1.274, > 100)	0.009	110/338	0.154 (-0.115, 1.333)	0.506	15/48	> 100 ^c (2.550, > 100)	0.003 ^c
Regression calibration (average of 10,000 doses)	24/72	12.154 (0.381, > 100)	0.020	110/338	0.100 (-0.099, 0.945)	0.574	15/48	> 100 ^c (3.085, > 100)	0.002 ^c
Full Monte Carlo maximum likelihood	24/72	42.119 (1.062, > 100)	0.010	110/338	0.102 (-0.161, 1.127)	0.577	15/48	> 100 ^c (2.114, > 100)	0.003 ^c

^a Adjusted using categories of cigarette smoking status (1) never smoked, (2) ex smoker/unknown, (3) current smoker, cigarette packyears, proxy status^b Adjusted using categories of cigarette smoking status (1) never smoked, (2) ex smoker/unknown, (3) current smoker, cigarette packyears, proxy status, categories of alcohol consumption (1) unknown, (2) never drank, (3) drank once/month or less, (4) drank 2–3 times/month, (5) drank once/week or more^c Indications of lack of convergence

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.