Correction to: Physiological and histopathological assessments of the susceptibility of different tomato (*Solanum lycopersicum*) cultivars to early blight disease

Nashwa M. A. Sallam · Heba-Alla S. AbdElfatah ·
Mona F. A. Dawood · Elhagag Ahmed Hassan ·
Mohamed S. Mohamed · Hadeel M. M. Khalil Bagy

Published online: 21 April 2021

© Koninklijke Nederlandse Planteziektenkundige Vereniging 2021

Correction to: Eur J Plant Pathol
<https://doi.org/10.1007/s10658-021-02263-2>

An incorrect version of fig. 8 was published in the original article and this erratum reflects the correct version.

The online version of the original article can be found at
<https://doi.org/10.1007/s10658-021-02263-2>

N. M. A. Sallam (✉) · H.-A. S. AbdElfatah ·
M. S. Mohamed · H. M. M. Khalil Bagy
Faculty of Agriculture, Plant Pathology Department, Assiut
University, Assiut 71526, Egypt
e-mail: nashwasallam@aun.edu.eg

M. F. A. Dawood · E. A. Hassan
Faculty of Science, Botany and Microbiology Department, Assiut
University, Assiut, Egypt

Fig. 8 Scanning electron micrographs showing healthy tomato CV1 (**a**), healthy tomato susceptible CV5 (**b**) compared with infected tomato CV1 (*A. solani* and *C. lunata* respectively) (**c** & **d**) that showing no colonization of fungal hyphae. Note the infected tomato susceptible cultivar (**e** & **f**) showing colonization of fungal mycelia and conidia formation