Correction to: *Bremia lactucae* populations on cultivated lettuce originate from prickly lettuce and are interconnected with the wild pathosystem

Fabian Runge · Ute Gärber · Aleš Lebeda · Marco Thines

Published online: 31 December 2021
© Koninklijke Nederlandse Planteziektenkundige Vereniging 2021

Correction to: Eur J Plant Pathol.
<https://doi.org/10.1007/s10658-021-02332-6>

This correction is published as typesetter overlooked several proofing corrections.

Original article has been updated.

The online version of the original article can be found at <https://doi.org/10.1007/s10658-021-02332-6>

F. Runge (✉)
IDENTXX GmbH, Maybachstr. 50, D-70469 Stuttgart, Germany
e-mail: fabian.runge@identxx.com

F. Runge · M. Thines
Senckenberg Biodiversity and Climate Research Center,
Senckenberg Gesellschaft für Naturforschung, Senckenberganlage
25, D-60325 Frankfurt am Main, Germany

F. Runge · M. Thines
Department of Biology, Institute of Ecology, Evolution, and
Diversity, Goethe University, Campus Riedberg,
Max-von-Laue-Str. 13, D-60439 Frankfurt am Main, Germany

U. Gärber
Julius Kühn-Institut, Federal Research Centre for Cultivated Plants,
Institute for Plant Protection in Horticulture and Forests,
Stahnsdorfer Damm 81, D-14532, Kleinmachnow, Germany

A. Lebeda
Faculty of Science, Department of Botany, Palacký University in
Olomouc, Šlechtitelů 27, CZ-78371 Olomouc, Czech Republic

M. Thines
Translational Biodiversity Genomics Centre (TBG),
Georg-Voigt-Str. 14-16, D-60325 Frankfurt am Main, Germany