



Relational Quantum Mechanics: Ozawa's Intersubjectivity Theorem as Justification of the Postulate on Internally Consistent Descriptions

Andrei Khrennikov¹

Received: 18 January 2024 / Accepted: 9 April 2024
© The Author(s) 2024

Abstract

The Ozawa's intersubjectivity theorem (OIT) proved within quantum measurement theory supports the new postulate of relational quantum mechanics (RQM), the postulate on internally consistent descriptions. But from OIT viewpoint postulate's formulation should be completed by the assumption of probability reproducibility. We remark that this postulate was proposed only recently to resolve the problem of intersubjectivity of information in RQM. In contrast to RQM for which OIT is a supporting theoretical statement, QBism is challenged by OIT.

Keywords Relational quantum mechanics · Ozawa intersubjectivity theorem · Postulate on internally consistent descriptions · Measurement process · Probability reproducibility · Unitary evolution · Entanglement of observables

1 Introduction

Recently Ozawa's intersubjectivity theorem (OIT) [1] proved within quantum measurement theory (e.g., [2–8]) is used as objection to QBism [9], to its basic interpretational statement about the individual agent perspective on measurement's outcomes [10–12]. (This privacy component of QBism was also criticized in my previous articles [13, 14], mainly from the probabilistic perspective.) Now QIT provides the strong argument that measurement's outcome is intersubjective and, although one can still interpret it as a private experience, such interpretation loses its elegance. QBists replied to my objection with two papers [15, 16] (the second one presents more elaborated counterargument).

In a few words the QBism's counterargument is that compatible measurement devices M_1 and M_2 considered in OIT should be related not to two distinct observers

✉ Andrei Khrennikov
Andrei.Khrennikov@lnu.se

¹ International Center for Mathematical Modeling in Physics and Cognitive Sciences, Linnaeus University, SE-351 95 Växjö, Sweden

Q_1 and O_2 as was done in my article [9], but to one observer O who performs these two measurements. As was clearly explained in papers [15, 16], this interpretation of such measurements is consistent with the general ideology of QBism. The important impact of article [9] to QBism is that this article attracted the attention (at least some) of QBists to quantum measurement theory [2–8]. QBism intensively explores one of the basic elements of this theory - POVMs. But QBists don't employ the important part of measurement theory - theory of *quantum measurement processes*. Each POVM is generated by a measurement process [3]. It is important to note that the same POVM can be generated by variety of measurement processes.

The discussions on the intersubjectivity problem in QBism stimulated my interest to this problem for other information interpretations of QM. A reviewer of one of my papers pointed that relational quantum mechanics (RQM) [17–21] was suffering of the same problem. It seems that it can't be resolved within "old RQM". Its resolution requires invention of an additional postulate on *internally consistent descriptions* (Postulate 6 in [21]). By treating this postulate within quantum measurement theory, one immediately see that OIT supports this postulate. So, in contrast to QBism which was challenged by OIT, RQM's Postulate 6 is mathematically justified via OIT. However, the OIT view on Postulate 6 leads to the conclusion that to match with quantum measurement theory, this postulate should be constrained by *the probability reproducibility condition* [1] (see section 3, equality (4)).

As well as QBism, RQM can earn from closer connection with quantum measurement theory [2–8]. Besides the present paper, an attempt to proceed in this direction was done in article [23]. However, the conclusions of this article are debatable. I am not an expert in RQM and I am not able to evaluate whether the authors of [23] correctly understood the conceptual premises of RQM.

We start with a brief recollection of the basics of RQM and the problem of intersubjectivity (section 2). We follow article [21]. Then we formulate OIT (section 3), see article [1] for its proof. In section 4 we couple OIT with Postulate 6 (on intersubjectivity) of RQM. Matching of quantum measurement theory and RQM is established in section 5. We should modify the interpretation and representation of the original von Neumann's measurement scheme to make it consistent with the RQM perspective on observer: "Any system, irrespectively of its size, complexity or else, can play the role of the textbook's quantum mechanical observer." (See [22]).¹ The crucial role of probability reproducibility (the basic condition of OIT) in analysis of the intersubjectivity problem in RQM is highlighted in section 6.

¹ This is the good place to cite another block from article [22]: "In textbook presentations, quantum mechanics is about measurement outcomes performed when an "observer" makes a "measurement" on a quantum system. What is an observer, if all physical systems are quantum? What counts as a measurement? Common answers invoke the observer being macroscopic, onset of decoherence, irreversibility, registration of information, or similar. RQM does not utilise anything of the sort."

2 Relational quantum mechanics and the problem of intersubjectivity

As was stated by Rovelli [17], RQM is based on the idea that “in quantum mechanics different observers may give different accounts of the same sequence of events.” It is crucial that “RQM is built on strong naturalistic intuitions; therefore, in RQM, the term ‘observer’ is understood in a broad sense, which allows that any physical system can be an ‘observer,’ so we do not have to accept that consciousness plays any fundamental role.” [21]. In the last aspect RQM differs crucially from QBism. As is emphasized in article [21], RQM has many attractive features and it resolves the basic quantum paradoxes (but QBism do this as well).

“However, some problems remain; in particular, there is a tension between RQM’s naturalistic emphasis on the physicality of information and the inaccessibility of certain sorts of information in current formulations of RQM. Thus, in this article, we propose a new postulate for RQM which ensures that all of the information possessed by a certain observer is stored in physical variables of that observer and thus is accessible by measurement to other observers. The postulate of cross-perspective links makes it possible for observers to reach intersubjective agreement about quantum events that have occurred in the past, thus shoring up the status of RQM as a form of scientific realism and allowing that empirical confirmation is possible in RQM.

Adding this postulate requires us to update some features of the ontology of RQM, because it entails that not everything in RQM is relational.”[21]

For readers convenience, here the list of all RQM’s postulates is presented [21]:

1. Relative facts: Events, or facts, can happen relative to any physical system.
2. No hidden variables: Unitary quantum mechanics is complete.
3. Relations are intrinsic: The relation between any two systems A and B is independent of anything that happens outside these systems’ perspectives.
4. Relativity of comparisons: It is meaningless to compare the accounts relative to any two systems except by invoking a third system relative to which the comparison is made.
5. Measurement: An interaction between two systems results in a correlation within the interactions between these two systems and a third one; that is, with respect to a third system W , the interaction between the two systems S and F is described by a unitary evolution that potentially entangles the quantum states of S and F .
6. Internally consistent descriptions: In a scenario where O_1 measures S , and O_2 also measures S in the same basis, and O_2 then interacts with O_1 to “check the reading” of a pointer variable (i.e., by measuring O_1 in the appropriate “pointer basis”), the two values found are in agreement.

OIT matches with Postulate 6, justifies it mathematically on the basis of quantum measurement theory, but with one important reservation: restriction of its applicability to measurement processes satisfying the probability reproducibility condition. Generally RQM can earn a lot from employing quantum measurement theory.

As was cited above in “RQM the term ‘observer’ is understood in a broad sense, which allows that any physical system can be an ‘observer’...”

In quantum measurement theory, ‘apparatus’ plays the key role. I would suggest to extend its meaning and operate in this theory with the notion observer. The latter has the RQM meaning, i.e., any physical system can be an observer. This the important novelty in application of quantum measurement theory and it will be discussed in more details (see section 5).

3 Ozawa intresubjectivity theorem

Von Neumann [2] described observables mathematically by Hermitian operators acting in Hilbert state space \mathcal{H} . They represent *accurate measurements*. Consider operators with totally discrete spectra $X \subset \mathbb{R} : A = \sum_{x \in X} x E_A(x)$, where $E_A(x)$ is projection on the eigensubspace for the eigenvalue x . The Born rule determines the probabilities:

$$P(A = x|\psi) = \langle \psi | E_A(x) | \psi \rangle.$$

The indirect measurement scheme involves the following components

- the states spaces \mathcal{H} and \mathcal{K} of the systems \mathcal{S} and the apparatus \mathcal{M} for measurement of some observable A ;
- the evolution operator $U = U(t)$ representing the interaction-dynamics for the system $\mathcal{S} + \mathcal{M}$;
- the meter observable M giving outputs of the pointer of the apparatus \mathcal{M} .

Let $E_M = (E_M(x))$ be the spectral family of the operator M ; here $E_M(x)$ are projections in Hilbert space \mathcal{K} . It is assumed that the compound system’s evolution is driven by the Schrödinger equation, so the evolution operator $U(t)$ is unitary.

Formally, an *indirect measurement model* for an observable A , introduced in [3] as a “measurement process”, is a quadruple

$$(\mathcal{K}, |\xi\rangle, U, M), \tag{1}$$

where $|\xi\rangle \in \mathcal{K}$ represents the apparatus state.

We explore the Heisenberg picture. To describe meter’s evolution, we represent it in the state space of the compound system, i.e., as $I \otimes M$. The meter observable evolves as

$$M(t) = U^*(t)(I \otimes M)U(t). \tag{2}$$

By the Born rule

$$P(M(t) = x|\psi\xi) = \langle \psi\xi | E_{M(t)}(x) | \psi\xi \rangle. \tag{3}$$

This is the probability distribution for the outputs of measurements done by the apparatus and given by the meter.

Definition A measurement process $(\mathcal{K}, |\xi\rangle, U, M)$ reproduces the probability distribution for quantum observable A (accurate von Neumann observable) if

$$P(A = x|\psi) = P(M(T) = x|\psi\xi). \tag{4}$$

Following [1], consider two remote observers O_1 and O_2 who perform joint measurements on a system S , in mathematical terms it means that the meter quantum observables of the corresponding measurement processes commute,

$$[M_1(t), M_2(t)] = 0.$$

Here each apparatus has its own state space, i.e., $\mathcal{K} = \mathcal{K}_1 \otimes \mathcal{K}_2$. We call such measurements local. In this situation the joint probability distribution is well defined

$$P(M_1(t) = x, M_2(t) = y|\psi\xi_1\xi_2) = \langle \psi\xi_1\xi_2 | E_{M_1(t)}(x)E_{M_2(t)}(y) | \psi\xi_1\xi_2 \rangle. \tag{5}$$

Theorem (OIT [1]) *Two observers performing the joint local and probability reproducible measurements of the same quantum observable A on the system S should get the same outcome with probability 1:*

$$P(M_1(T) = x, M_2(T) = y|\psi\xi_1\xi_2) = \delta(x - y)P(E = x|\psi) = \delta(x - y)\|E(x)\psi\|^2. \tag{6}$$

4 Postulate on internally consistent description in the view of Ozawa intersubjectivity theorem

In OIT, as in Postulate 6 [RQM], observer O_1 measures S , and observer O_2 also measures S in the same basis. In our terms the later means that they measure the same observable A . Then O_2 can interact with O_1 to “check the reading” of its pointer variable, the two values found are in agreement.

The important condition which is missed in Postulate 6 is the probability reproducibility condition, only under this conditions outcomes of measurements performed by O_1 and O_2 coincide. Therefore, it is natural to complete postulate 6 by this condition. But this is just my viewpoint and may be Carlo Rovelli would not support it.

Moreover, due to OIT observer O_2 even need not to perform a measurement on O_1 . If observers are sure in validity of quantum theory, then they can be sure that they get the same outcome.

I suggest the following modification of Postulate 6:

Postulate 6[OIT]. *Under the condition of probability reproducibility, in a scenario where O_1 measures S , and O_2 also measures S in the same basis, the two values found are in agreement.*

5 Matching von Neumann's theory of quantum measurement with Rovelli's interpretation of observer

As was emphasized in introduction, in this paper the meaning of the notion “apparatus” is extended, we operate with the notion “observer”. And in RQM any physical system can play the role of an observer. The following question naturally arises:

Are there some constraints on the notion of “apparatus” as it is defined in quantum measurement theory which might prevent it from encompassing all physical systems?

This paper is based on the modern version of von Neumann's approach to the problem of quantum measurement. Therefore it is natural to return to the original von Neumann's work [2] (chapter 6). Our aim is to show that, in spite some interpretational differences, are von Neumann's and Rovelli's viewpoints on the quantum measurement process generally close to each other. In fact, elements of such comparative analysis can be found in the original article of Rovelli [17]. Our discussion is more detailed. Moreover, we bring into our analysis a new theoretical construction - “entanglement of observables”, A_1A_2 -entanglement for a pair of compatible observables. This generalization of entanglement was presented in articles [28, 29]. It matches both with von Neumann's description of a measurement process and RQM's Postulate 5, on measurement. The probability reproducibility condition of OIT can also be formulated in terms of “entanglement of observables”, in the special case of von Neumann's model.

We start with some interpretational remarks on measurement theory presented in [2] (see section 1, chapter 6). In this theory both a quantum system and measurement apparatus are described within the quantum formalism. Thus, although the apparatus is macroscopic, it is, nevertheless, described as a quantum system. Moreover, the mathematical model is symmetric w.r.t. system-apparatus interchange. Thus, it is applicable to macroscopic systems, (cf. with RQM). This macro-applicability was recently supported by quantum-like modeling, applications of the mathematical formalism of quantum measurement theory outside of physics, e.g., in cognition and decision making; see [24, 25]. We also remark that von Neumann's description of the measurement process doesn't involve decoherence (again cf. with RQM).

At the same time von Neumann's interpretation differs from RQM, since he refers to subjective nature of an observer (cf. with QBism). However, in contrast to QBists he didn't emphasize the private character of observer's measurement experience, he neither use the subjective interpretation of probability. Moreover, he diminished the value of subjective experience by referring to psycho-physical parallelism. At the same time he states [2] (p. 420): “Indeed experience only makes statements of this type: an observer has made a certain (subjective) observation; and never any like this: a physical quantity has a certain value” This is the Bohr's viewpoint on quantum observables [26, 27]: their outcomes aren't the objective properties of systems under measurements. They are generated during the complex process of interaction between system S and measurement apparatus

\mathcal{M} used to measure quantum observable A . At this point the positions of von Neumann and Bohr coincide. In [2] interaction between the system and apparatus is mathematically described by unitary operator U , see below.

Now we turn to the formalism developed in chapter 6 [2]. The measurement process is described in Sect. 3. There are considered three systems: \mathcal{S} , observed system, \mathcal{M} , a measurement apparatus, and \mathcal{O} , observer. It is important to remark [2] (p. 421) that “... \mathcal{O} itself reminds outside of calculations.” (See also p. 439). Hence, all mathematical considerations are related to the pair of physical systems, $(\mathcal{S}, \mathcal{M})$. And in section 3, “measurement section”, von Neumann refers to calculations done in section 2 (“operator entanglement section”).

In fact, von Neumann considered a kind of entanglement which can be called “*entanglement of observables*” (see [28, 29]). Two compatible observables A_1 and A_2 can be called entangled in the state $|\Phi\rangle$, if their eigenvalues can be enumerated in such a way, $(a_{1k}, k = 1, 2, \dots)$ and $(a_{2k}, k = 1, 2, \dots)$ that probability

$$P(A_1 = a_{1k}, A_2 = a_{2m}|\Phi) = 0, k \neq m, \tag{7}$$

or

$$\sum_k P(A_1 = a_{1k}, A_2 = a_{2k}|\Phi) = 1, \tag{8}$$

or

$$P(A_1 = a_{1k}|\Phi) = P(A_1 = a_{1k}, A_2 = a_{2k}|\Phi), \tag{9}$$

$$P(A_2 = a_{2k}|\Phi) = P(A_1 = a_{1k}, A_2 = a_{2k}|\Phi), \tag{10}$$

or

$$P(A_1 = a_{1k}|A_2 = a_{2k}|\Phi) = 1, P(A_2 = a_{2k}|A_1 = a_{1k}|\Phi) = 1, \tag{11}$$

under the constraint $P(A_1 = a_{1k}, A_2 = a_{2k}|\Phi) \neq 0$. Here conditional probability w.r.t. state $|\Phi\rangle$ is defined by the Bayes formula, e.g., $P(A_1 = a_{1k}|A_2 = a_{2k}|\Phi) = P(A_1 = a_{1k}, A_2 = a_{2k}|\Phi)/P(A_2 = a_{2k}|\Phi)$ (since the operators commute we can operate within the classical probability framework).

From my viewpoint, the notion of such A_1A_2 -entanglement matches better with RQM, than the standard notion of the entangled state (see Postulate 5 of RQM). We remark that a state $|\Phi\rangle$ can be A_1A_2 -entangled, but not entangled in the ordinary sense. Moreover, it was shown that, for any state $|\Phi\rangle$, there exist pairs of operators which are entangled in this state [2] (section 2). We note that von Neumann didn't use the term “entanglement” for observables. He wrote about establishing “one-to-one correspondence between the possible values of certain quantities” in two systems. As suggested in articles [28, 29] that such correspondence can be called observables' entanglement (and it has some features of state entanglement).

In formal mathematical calculations of Sects.2,3 the notations \mathcal{S} and \mathcal{M} lose their meaning and we consider two arbitrary physical systems S_1 and S_2 in the

states $|\psi\rangle$ and $|\xi\rangle$ belonging to the state spaces \mathcal{H}_1 and \mathcal{H}_2 . Initially the compound system (S_1, S_2) is in the state

$$|\Phi_0\rangle = |\psi\rangle|\xi\rangle \tag{12}$$

belonging to the state space $\mathcal{H}_1 \otimes \mathcal{H}_2$. Consider observables for systems S_1, S_2 represented by Hermitian operators A_1, A_2 . Let $(|\psi_k\rangle)$ and $(|\xi_k\rangle)$ be corresponding bases consisting of eigenvectors, $A_1|\psi_k\rangle = a_{1k}|\psi_k\rangle, A_2|\xi_k\rangle = a_{2k}|\xi_k\rangle$, i.e.,

$$A_1 = \sum_k a_{1k}|\psi_k\rangle\langle\psi_k|, A_2 = \sum_k a_{2k}|\xi_k\rangle\langle\xi_k|. \tag{13}$$

To simplify considerations, we assume that these operators have non-degenerate spectra.

It is proved [2] (with the explicit mathematical formula) that there exist a unitary operator $U : \mathcal{H}_1 \otimes \mathcal{H}_2 \rightarrow \mathcal{H}_1 \otimes \mathcal{H}_2$, such that the initial state $|\Phi_0\rangle$ is transferred into A_1A_2 -entangled state:

$$|\Phi\rangle = U|\Phi_0\rangle = \sum_k c_k|\psi_k\rangle|\xi_k\rangle, \tag{14}$$

where

$$c_k = \langle\psi|\psi_k\rangle \tag{15}$$

(cf. with EPR article [30]). This mathematical result can be reformulated in the language of quantum observables, as following. *The outcomes of the observables A_1 and A_2 are perfectly correlated.* The outcome $A_1 = a_{1k}$ is perfectly correlated with the outcome $A_2 = a_{2k}$. Thus, measurement of A_2 can be considered as indirect measurement of A_1 and vice versa.

In the light of the above analysis of von Neumann’s construction [2] (chapter 6), the assignment to the systems S_1 and S_2 the meaning of a system under observation and a measurement apparatus is the purely interpretational issue.

Now turn to sect. 3 (chapter 6 [2]) and set $S_1 = \mathcal{S}, S_2 = \mathcal{M}$ and $A_1 = A$, observable on \mathcal{S} measured with \mathcal{M} and $A_2 = M$, pointer observable. Due to AM -entanglement M -outcomes can be identified with the corresponding outcomes of the observable A . Although the observer \mathcal{O} was excluded from calculations, it plays the role: the quantity M is observed by \mathcal{O} . This mathematical description of the measurement process matches with RQM Postulate 5:

Measurement: An interaction between two systems results in a correlation within the interactions between these two systems and a third one; that is, with respect to a third system W , the interaction between the two systems S and F is described by a unitary evolution that potentially entangles the quantum states of S and F .

Subjective nature of observable \mathcal{O} mentioned (but not so much highlighted) by von Neumann doesn’t play any role in the formal scheme presented in [2].

6 Probability reproducibility condition: Postulate 6 [RQM] vs. Postulate 6 [OIT]

We point out that the measurement processes modeled by von Neumann satisfy the probability reproducibility condition. We remark that (15) implies that

$$P(A_1 = a_{1k}, A_2 = a_{2k} | \Phi) = |c_k|^2 = |\langle \psi | \psi_k \rangle|^2 = P(A_1 = a_{1k} | \psi). \tag{16}$$

And by using equalities (9), (10) we obtain

$$P(A_1 = a_{1k} | \psi) = P(A_2 = a_{2k} | \Phi), \tag{17}$$

We remark that

$$P(A_2 = a_{2k} | \Phi) = |\langle (I \otimes E_{A_2}) U \psi_\xi | U \psi_\xi \rangle|^2 = |\langle U^* (I \otimes E_{A_2}) U \psi_\xi | \psi_\xi \rangle|^2.$$

Here $U = U(T) = e^{-iTH/\hbar}$. Set $A_2(T) = U^*(T)(I \otimes A_2)U(T)$. Thus, we obtain the probability reproducibility condition:

$$P(A_1 = a_{1k} | \psi) = P(A_2(T) = a_{2k} | \psi_\xi), \tag{18}$$

or by selection $a_{1k} = a_{2k} = a_k$, and setting $A_1 = A, A_2 = M$, we get

$$P(A = x | \psi) = P(M(T) = x | \psi_\xi), \tag{19}$$

for any $x = a_k$.

Thus, if one restricts measurement processes to those described in sects.2, 3 (chapter 6 [2]), then the condition of probability reproducibility in **Postulate 6[OIT]** is redundant.

Generally measurement process doesn't guarantee probability reproducibility. Such measurements are noisy. Therefore it is not surprising that intersubjectivity postulate can be violated. Measurements performed by O_1 and O_2 are disturbed by noises and their outcomes need not coincide.

It is important to note that the class of measurement processes satisfying the probability reproducibility condition doesn't coincide with von Neumann's measurement processes described in [2]. To discuss this important issue, we have to refer to the notion of generalized quantum observable, POVM.

Consider only discrete POVMs valued in finite sets, $X = \{x_1, \dots, x_m\}$. POVM is a map $x \rightarrow \Pi(x)$: for $x \in X, \Pi(x)$ is a positive contractive self-adjoint operator: $0 \leq \Pi(x) \leq I$, called an *effect*; the family of effects form the resolution of unity $\sum_x \Pi(x) = I$. This map defines an operator valued measure on algebra of all subsets of set X . For $O \subset X, \Pi(O) = \sum_{x \in O} \Pi(x)$. POVM Π represents statistics of measurements:

$$P(\Pi = x | \psi) = \langle \psi | \Pi(x) | \psi \rangle. \tag{20}$$

The resolution of unity condition is the operator-measure counterpart of the condition normalization by 1 for usual probability measures. Any observable A given by

Hermitian operator can also be represented as POVM of the special type – PVM $E_A = (E_A(x))$.

Any measurement process given by quadruple (1) generates a generalized observable given by POVM

$$\Pi(x) = \langle \xi | E_{M(T)}(x) | \xi \rangle. \quad (21)$$

As was shown by Ozawa [3], any POVM, generalized observable, can be generated in this way, from some quantum measurement process. The tricky point which is often ignored in quantum information theory is that the same observable, POVM, can be generated by a variety of measurement processes.

Now we turn to the probability reproducibility condition. As was shown in [1], a measurement process satisfies to this condition if and only if it generates POVM, as (21), which coincides with PVM of the operator A , i.e., for any $x \in X$, $\Pi(x) = E_A(x)$. We repeat once again that a plenty of measurement processes can generate the same PVM.

The violation of the probability reproducibility condition implies that a measurement process generates POVM which is not PVM of the von Neumann observable A .

The original Postulate 6 [RQM] doesn't involve the probability reproducibility condition. Thus, there are the following situations. The general theory of quantum measurement processes allows outcomes to be different, but Postulate 6 [21] would still require them to be the same. One of the possibilities to resolve this discrepancy is to proceed with the original Postulate 6 [RQM], but with von Neumann's measurement processes [2]. Another possibility is to proceed with Postulate 6 [OIT], i.e., restrict the class of quantum measurement processes to those satisfying the probability reproducibility condition.

We also remark that we considered measurement processes for von Neumann observables, operator A is Hermitian. As is pointed out in article [1], OIT does not hold for measurements of generalized observables. In [1] one can find an example of generalized observable A represented by POVM $A = (A(x))$, where $A(x), x \in X$, are effects, and two compatible measurement processes $M_1(t)$ and $M_2(t)$ satisfying the probability reproducibility condition, but violating the intersubjectivity condition. (Here 'compatibility' has the meaning of compatibility of generalized observables). Thus, for generalized observables, one should give up intersubjectivity. As was already mentioned, generalized observables are unsharp and their measurements are noisy. Therefore, one can't expect coincidence of the outcomes of measurements, even for measurement processes satisfying the probability reproducibility condition.

Acknowledgements This study was partially supported by the EU-project: CA21169 - Information, Coding, and Biological Function: the Dynamics of Life (DYNALIFE). I would like to thank Masanao Ozawa for numerous lessons on quantum measurement theory, discussions on intersubjectivity of quantum measurements, and hospitality during my visits to Nagoya during the last 20 years. I also would like to thank Christopher A. Fuchs for 25 years of discussions on the QBism's place within quantum theory. (In fact, my interest to intersubjectivity in RQM was ignited by analysis of the intersubjectivity problem in QBism in the view of OIT [9]; cf. [15, 16].)

Author contributions I am the only author of this paper. So, I wrote the main manuscript text and reviewed it, as well as its revisions.

Funding Open access funding provided by Linnaeus University.

Data availability No datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors declare no Conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ozawa, M.: Intersubjectivity of outcomes of quantum measurements. (2019) <https://arxiv.org/abs/1911.10893>
2. Von Neumann, J., *Mathematische Grundlagen der Quantenmechanik*. Springer-Verlag, Berlin,: English translation: *Mathematical Foundations of Quantum Mechanics*; Princeton University Press: Princeton. NJ, USA (1932) (1955)
3. Ozawa, M.: Quantum measuring processes for continuous observables. *J. Math. Phys.* **25**, 79–87 (1984)
4. Busch, P., Lahti, P., Mittelstaedt, P.: *The Quantum Theory of Measurement*, LNP m2, Springer, 1991, 2nd rev. ed., (1996)
5. Ozawa, M.: An operational approach to quantum state reduction. *Ann. Phys. (N.Y.)* **259**, 121–137 (1997)
6. Holevo, A.S.: *Statistical Structure of Quantum Theory*. Springer, Berlin (2001)
7. Busch, P., Lahti, P., Pellonpää, J.-P., Ylino, K.: *Quantum Measurement*. Springer, Berlin (2016)
8. Ozawa, M.: Soundness and completeness of quantum root-mean-square errors. *Quant. Inf.* **5**, 1 (2019)
9. Khrennikov, A.: Ozawa's Intersubjectivity Theorem as objection to QBism individual agent perspective. *Int. J. Theor. Phys.* **63**, 1–9 (2024)
10. Fuchs, C.A., Mermin, N.D., Schack, R.: An Introduction to QBism with an Application to the Locality of Quantum Mechanics. *Am. J. Phys.* **82**, 749 (2014)
11. Fuchs, C.A.: Notwithstanding Bohr, the reasons for QBism. *Mind Matter* **15**(2), 245–300 (2017)
12. Fuchs, C. A.: QBism, Where Next?. (2023) arXiv preprint [arXiv:2303.01446](https://arxiv.org/abs/2303.01446)
13. Khrennikov, A.: External Observer Reflections on QBism, Its Possible Modifications, and Novel Applications, In: *Quantum Foundations*, STEAM-H: Science, Technology, Engineering, Agriculture, Mathematics & Health; Khrennikov A. and Toni B. Eds.; Springer, Cham, pp. 93–118 (2018)
14. Khrennikov, A.: Towards better understanding QBism. *Found. Sc.* **23**(1), 181–195 (2018)
15. Stacey, B. C.: Whose Probabilities? About What? A Reply to Khrennikov. [arXiv:2302.09475](https://arxiv.org/abs/2302.09475)
16. Schack, R.: When will two agents agree on a quantum measurement outcome? Intersubjective agreement in QBism. [arXiv:2312.07728](https://arxiv.org/abs/2312.07728) [quant-ph]
17. Rovelli, C.: Relational Quantum Mechanics. *Int. J. Theor. Phys.* **35**(8), 1637–1678 (1996)
18. Rovelli, C.: Space is blue and birds fly through it. *Phil. Trans. R. Soc. A* **376**, 20170312 (2017)

19. Martin-Dussaud, P., Rovelli, C., Zalamea, F.: The Notion of Locality in Relational Quantum Mechanics. *Found. Phys.* **49**, 96–106 (2019)
20. Rovelli, C.: *Helgoland: making sense of the quantum revolution*, Riverhead Books, Penguin Random House LLC, 2021. Original, in Italy, (2020)
21. Adlam, E., Rovelli, C.: Information is physical: Cross-perspective links in relational quantum mechanics. *Philos Phys.* **1**(1), 1–19 (2023)
22. Laudisa, F., Rovelli, C.: Relational Quantum Mechanics, The Stanford Encyclopedia of Philosophy, Eds.: E. N. Zalta and Uri Nodelman, Metaphysics Research Lab Publisher, Stanford University; <https://plato.stanford.edu/archives/sum2024/entries/qm-relational/>
23. Lahti, P., Pellonpää, J.-P.: An attempt to understand relational quantum mechanics. *Int. J. Theor. Phys.* **62**, 170 (2023)
24. Ozawa, M., Khrennikov, A.: Application of theory of quantum instruments to psychology: combination of question order effect with response replicability effect. *Entropy* **22**(1), 1–9436 (2020)
25. Ozawa, M., Khrennikov, A.: Modeling combination of question order effect, response replicability effect, and QQ-equality with quantum instruments. *J. Math. Psych.* **100**, 102491 (2021)
26. Bohr, N.: *The Philosophical Writings of Niels Bohr*. Ox Bow Press, Woodbridge, UK (1987)
27. Plotnitsky, A.: *Niels Bohr and Complementarity: An Introduction*; Springer: Berlin. Germany; New York, NY, USA (2012)
28. Basieva, I., Khrennikov, A.: Conditional probability framework for entanglement and its decoupling from tensor product structure. *J. Phys. A: Math. Theor.* **55**(39), 395302 (2022)
29. Basieva, I., Khrennikov, A.: Entanglement of observables: quantum conditional probability approach. *Found. Phys.* **53**, 84 (2023)
30. Einstein, A., Podolsky, B., Rosen, N.: Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.* **47**, 777–780 (1935)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.