CORRECTION

Correction to: Nitrous oxide emission factors in conventionally and naturally simulated cattle urine patches

M. O'Neill · S. Saggar · K. G. Richards · J. Luo · B. P. Singh · P. Mehra ·
P. J. Forrestral

Published online: 7 October 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2021

Correction to: Nutr Cycl Agroecosyst
<https://doi.org/10.1007/s10705-021-10162-5>

data for only the NZ1 site as labelled. This has been corrected with this Correction.

In the original publication of the article Fig. 1 contains the meteorological data of all three experimental sites (NZ1, NZ2 and IRE) rather than the meteorological

The original article can be found online at <https://doi.org/10.1007/s10705-021-10162-5>.

M. O'Neill (✉) · K. G. Richards · P. J. Forrestral
Teagasc, Environment, Soils and Land Use Department,
Johnstown Castle Research Centre, Co. Wexford, Ireland
e-mail: macdara.oneill@teagasc.ie

S. Saggar
Manaaki Whenua –Landcare Research, Palmerston North,
New Zealand

J. Luo
AgResearch Ruakura Research Centre, Private Bag 3123,
Hamilton, New Zealand

B. P. Singh · P. Mehra
NSW Department of Primary Industries, Elizabeth
Macarthur Agricultural Institute, Menangle, Australia

Fig. 1 Daily air temperature (°C), rainfall (mm) and WFPS (%) data in NZ1. Air temperature and rainfall was obtained from the synoptic weather station adjacent to the site. WFPS was

obtained from the mean soil volumetric water content measured at 50 mm and 100 mm depths using time-domain reflectometers

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.