

A method for quantifying differentiation between populations at multi-allelic loci and its implications for investigating identity and paternity

David J. Balding · Richard A. Nichols

Published online: 9 August 2007
© Springer Science+Business Media B.V. 2007

Erratum to: Genetica 96: 3–12 (1995)
DOI 10.1007/BF01441146

This erratum is being published as a result of incorrect parameters of Table 2.

Table 2 should be now read as:

Table 2 Single-locus likelihood ratios for paternity when the mother's genotype is AA. Blank entries indicate that the alleged father is excluded

Alleged Father	Child	
	AA	AB
AA	$\frac{4F+(1-F)p_A}{1+3F}$	
AB	$2\left(\frac{3F+(1-F)p_A}{1+3F}\right)$	$\left(\frac{2F+(1-F)p_B}{1+3F}\right)$
BB		$\frac{2F+(1-F)p_B}{1+3F}$
BC		$2\left(\frac{F+(1-F)p_B}{1+3F}\right)$

The online version of the original article can be found under doi:
[10.1007/BF01441146](https://doi.org/10.1007/BF01441146).

D. J. Balding (✉) · R. A. Nichols
School of Mathematical Sciences and School of Biological
Sciences, Queen Mary & Westfield College, University
of London, Mile End Road, London E1 4NS, UK
e-mail: d.balding@imperial.ac.uk