

QTL analysis reveals the genetic architecture of domestication traits in Crisphead lettuce

Yorike Hartman · Danny A. P. Hooftman ·
M. Eric Schranz · Peter H. van Tienderen

Received: 19 July 2012 / Accepted: 12 November 2012 / Published online: 28 November 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract The genetic architecture of crop domestication is generally characterized by three trends: relatively few genomic regions with major QTL effects are involved, QTL are often clustered, and alleles derived from the crop do not always contribute to the crop phenotype. We have investigated the genetic architecture of lettuce using a recombinant inbred line population from a cross between a crop *Lactuca sativa* ('Salinas') and its wild relative *L. serriola*. Few genomic regions with major QTL, plus various intermediate QTL, largely control the

transition from wild to cultivated Crisphead lettuce. Allelic effects of all major QTL were in the expected direction, but there were intermediate QTL where the crop contributed to the wild phenotype and vice versa. We found two main regions with clusters of QTL, one on linkage group 3, where the crop allele induced lower seed output, another on linkage group 7, where the crop allele caused a delay in flowering time. Potentially, knowledge of genetic changes due to the domestication could be relevant for the chance that a transgene inserted in a crop genome will spread to wild relatives due to hitchhiking effects. If a transgene would be inserted in one of these regions, background selection on the crop alleles may lead to a reduced fitness of hybrids with the transgene. QTL research on the effects of domestication genes can thus indicate regions in the crop genome that are less likely to introgress, although these still need to be verified under field conditions.

Y. Hartman (✉) · D. A. P. Hooftman ·
M. Eric Schranz · P. H. van Tienderen
Institute for Biodiversity and Ecosystem Dynamics,
University of Amsterdam, PO box 94248,
1090 GE Amsterdam, The Netherlands
e-mail: y.hartman@uva.nl

D. A. P. Hooftman
e-mail: Dann1@ceh.ac.uk

M. Eric Schranz
e-mail: eric.schranz@wur.nl

P. H. van Tienderen
e-mail: P.H.vanTienderen@uva.nl

D. A. P. Hooftman
Centre for Ecology and Hydrology,
Wallingford OX10 8BB, UK

Present Address:
M. Eric Schranz
Biosystematics Group, Wageningen University,
6700 AP Wageningen, The Netherlands

Keywords Crop–wild hybrids · Domestication
traits · GMOs · *Lactuca*

Introduction

The study of crop domestication has received much interest from crop breeders and evolutionary biologists alike (Burger et al. 2008; Hancock 2005). Crops serve as a model for evolution and adaptation, because of the knowledge of crop history and the selective pressures

that accompanied domestication (Purugganan and Fuller 2009). Quantitative Trait Locus (QTL) studies of domestication-related traits using crosses between crops and their wild relatives have given insights into the origin of crops and speed of domestication (Gross and Olsen 2010). Furthermore, crop–wild crosses provide potential source material for crop improvement (Isemura et al. 2010; Vaughan et al. 2007), insight on hybridization and introgression between crops and their wild relatives (Hancock 2005; Hooffman et al. 2009), and show genomic regions under selection in crop–wild hybrids (Baack et al. 2008; Dechaine et al. 2009).

Three important trends in crop genetic architecture are often highlighted and are the focus of this study: effect size of QTL, clustering of QTL, and directionality of allelic effects. First, the transition from wild-type to cultivated crop is often controlled by relatively few genomic regions with major QTL effects (Burger et al. 2008; Gross and Olsen 2010; Ross-Ibarra 2005). Such regions contain very low variation among cultivars, indicating a rapid and uniform fixation caused by continuous artificial selection over many years (Burke et al. 2007). In maize (Doebley and Stec 1993), rice bean (Isemura et al. 2010), eggplant (Doganlar et al. 2002), tomato (Frery et al. 2000), and rice (Cai and Morishima 2002), a small number of QTL cause large phenotypic changes in seed shattering, seed and fruit size, and branching patterns. In contrast, the crop genetic architecture of sunflower domestication is characterized by a relative high number of QTL, many with minor or intermediate effect (Burke et al. 2002, 2007). Second, domestication-related QTL are not uniformly distributed over the genome, but are often clustered (Burger et al. 2008; Ross-Ibarra 2005). For example, clustering has been found in sunflower (Burke et al. 2002; Dechaine et al. 2009), azuki bean (Kaga et al. 2008), rice bean (Isemura et al. 2010), rice (Cai and Morishima 2002), eggplant (Doganlar et al. 2002), pearl millet (Poncet et al. 2000), and tomato (Prudent et al. 2009). Third, the majority of QTL usually show phenotypic effects in the expected direction; i.e. crop alleles contribute to a crop phenotype. However, the opposite also occurs, with crop alleles contributing to the wild phenotype and wild alleles to the crop phenotype (Ross-Ibarra 2005). For example, in sunflower the minority of plant height, number of branches and ray size QTL were in the expected direction (Burke et al. 2002).

The underlying cause of these three trends is not yet well understood. It has been suggested that the relative importance of major QTL, the presence of QTL clusters and phenotypic effects reflects a rapid domestication driven by strong directional selection (Rieseberg et al. 2002; Ross-Ibarra 2005), whereas a majority of intermediate and minor QTL might reflect a gradual process with multiple domestication events (Burke et al. 2002). The clustering seen in many crop species might be caused by the fact that species with beneficial genes in clusters are more easily domesticated than species with beneficial genes more dispersed over the genome (Burger et al. 2008), or alternatively, such clustering might be the result of single genes with pleiotropic effects over several traits. In any case, more detailed analyses are necessary to draw more definitive conclusions.

Domestication research might play an important role in the debate surrounding genetically modified (GM) crops. With the introduction of GM crops concerns have arisen about hybridization and the chances for transgene escape from crops to their wild relatives. A particular concern is possible negative ecological effects, such as increased invasiveness of the wild relative (Chapman and Burke 2006; Stewart et al. 2003). In order to minimize chances of transgene escape, mitigation strategies have been proposed where the transgene is in linkage with an allele that is selected against in the wild and therefore is more likely to be purged from the wild population (Gressel 1999; Kwit et al. 2011; Stewart et al. 2003). Locating domestication-related QTL might be a way to pinpoint genomic areas where the crop allele confers a fitness disadvantage to hybrid individuals. Many crops share a number of traits, termed the ‘domestication syndrome’, which were selected for in early stages of domestication and made crops easier to cultivate (Hammer 1984; Harlan 1992). Although many crops are reported to hybridize with their wild relatives (Ellstrand 2003), these domestication traits were until recently seen as unable to introgress into the wild and were generally believed to impose negative fitness effects on hybrid individuals in the wild (Hails and Morley 2005). However, Ellstrand et al. (2011) showed 13 examples of suggested evolution of weediness from domesticated ancestors of which seven are via hybridization with wild relatives.

Due to their economic importance, much of the recent, molecular marker based, domestication

research has focused on seed-propagated annual crop species (Burger et al. 2008; Gross and Olsen 2010), in particular cereal crops such as wheat, maize, and rice. Whether or not the three general patterns described earlier extend to other types of crop remains to be seen (Gross and Olsen 2010). Within the Compositae there are two major crops, namely sunflower and lettuce (Dempewolf et al. 2008). We study domestication traits in lettuce (*Lactuca sativa*), an annual vegetable crop grown for its leafy head rather than for its seeds (achenes). The recent availability of genomic sequences, genetic maps, and genotyped Recombinant Inbred Lines (RILs) makes extensive analyses of domestication-related traits in lettuce possible (<http://compgenomics.ucdavis.edu/>).

Lactuca serriola is fully cross compatible and interfertile with cultivated lettuce (de Vries 1990). It occurs in Europe, North America, South Africa, Asia and Australia, mainly in pioneer, and/or disturbed ruderal habitats, but also as a weed in agricultural fields (D'Andrea et al. 2009; Lebeda et al. 2001). In the past 50–60 years, it has expanded its range dramatically in Western Europe (D'Andrea et al. 2009; Hooftman et al. 2006). This invasiveness is probably closely linked to human activities (D'Andrea et al. 2009; Lebeda et al. 2001) although effects of introgressed crop genes to the wild population cannot be excluded (Hooftman et al. 2006). Lettuce is predominantly selfing, but also insect pollinated with a reported outcrossing rate of approximately 1–5 % (D'Andrea et al. 2008; Giannino et al. 2008). In a field experiment, Hooftman et al. (2005, 2007) showed that artificial hybrids of *L. serriola* and *L. sativa* had an increased rather than a decreased fitness compared to the wild-type. This suggests that hybridization can lead to novel genetic combinations that can be more vigorous than the original parent lines and that crop genes in lettuce are not necessarily negatively selected for.

It has already been shown that RILs of the cross between cultivated *L. sativa* and wild *L. serriola* used in this study have differentiated root architecture (Johnson et al. 2000), seed and seedling traits (Argyris et al. 2005), and leaf biophysical and developmental traits (Zhang et al. 2007). To our knowledge no classical or quantitative genetic data are available on the genetic basis of traits that differ between wild and cultivated lettuce: the available studies have mostly focussed on offspring fertility of different *Lactuca* species and among cultivars crosses (Thompson et al.

1941; Lindqvist 1960a, b; Prince et al. 1979; de Vries 1990). Generally, with regard to morphological domestication traits, lettuce shows a reduced natural seed (achene) dispersal and increased seed size compared to the wild-type, similar to cereal crops (de Vries 1997). More typical for a leafy vegetable, domestication in lettuce has been targeted at delay of bolting, increased head formation, loss of spines, decrease in latex content and bitter taste, and a change from long serrate leaves to broad almost circular leaves (de Vries 1997). The cultivar used in this study, Crisphead or Iceberg lettuce (*L. sativa* 'Salinas'), is mainly cultivated in the United States and was developed to form very tight dense heads (de Vries 1997).

In this study, we will focus on the identification of domestication- and fitness-related QTL for various life stages and compare the results with the genetic architecture of domestication found in previous lettuce research and in other crops. Our focus is on developmental, leaf shape, and seed output traits. We will discuss the results in the view of the development of new transgenic crops, and the likelihood of introgression of crop genes to wild relatives.

Materials and methods

Plant material and growing conditions

We used an existing Recombinant Inbred Line (RIL) population, selfed for nine generations, from a cross between the crop *L. sativa* 'Salinas' (*L. sativa* L. var. *capitata* L. nidus jæggeri Helm) and its wild relative *L. serriola* f. *serriola* (UC96US23). The RILs are from the same set as used in Argyris et al. (2005), Johnson et al. (2000), and Zhang et al. (2007). In total, 114 RILs were grown in the greenhouse under 6 h dark and 18 h light, a minimum of 18 °C, under 600W SON T-Agro lamps generating on average 160 $\mu\text{mol}/\text{m}^2/\text{sec}$ at plant level. One plant per line was grown from the end of November 2007 and followed through the entire lifecycle until the last plants produced seed in August 2008.

In addition, we performed a germination experiment in a germination cabinet under 16 h of light at 20 °C and 8 h dark with 15 °C. The experiment lasted 9 days in total until the majority of lines reached 90–100 % germination. Achenes (further referred to as seeds) were placed in Petri dishes on filter paper and

watered with sterilized water to induce germination. We added a small amount of TMTD (tetramethylthiuram-disulfide) powder to prevent the growth of fungi on the seeds. We phenotyped the 114 RIL and parent lines; all lines were replicated twice. Each Petri dish contained three to four lines with each ten seeds. Lines were randomized and, to prevent any position effects, trays and Petri dishes were shuffled around three times every day.

Trait measurements

At several life stages, we measured different domestication- and fitness-related traits focusing on developmental, leaf shape, flowering phenotype, spines, seed phenotype, and seed output traits (Table 1).

Development

Germination was counted daily for 9 days; seeds were considered as germinated if the two cotyledons were fully emerged from the seeds and roots were visible. Since most lines reached 90–100 % germination, we calculated the time point at which 50 % of seeds had germinated using an existing R-code (version 2.12.1; <http://lukemiller.org/index.php/2010/02/calculating-ld50-median-lethal-temperature-aka-ld50-quickly-in-r/>, March 11, 2011).

At 30 and 60 days after sowing, we measured several plant characteristics for one plant per RIL. Plant height was measured and the number of rosette and stem leaves was counted. From this we calculated the proportion of stem leaves by dividing the number of stem leaves by the total number of rosette and stem leaves. At 60 days, the majority of lines had bolted and produced shoots, therefore we only used the number of stem leaves in the final analysis. Flowering and seed set was measured as the number of days from sowing to the production of the first flower or seeds, respectively. At seed set, we also measured plant height, stem thickness of the main stem at 10 cm above the soil, and we counted the total number of stem leaves.

Leaf shape

Leaf shape was determined quantitatively as well as qualitatively. At 30 days after sowing, we measured the length and maximum width of the four biggest rosette leaves. At the flowering stage, we also

measured the length and maximum width of two leaves halfway the main stem. In both cases, the biggest leaf was scanned and images were analyzed with ImageJ v1.41 (Abramoff et al. 2004) to determine leaf surface and circularity as a measure of the amount of serration of the leaves. Circularity was calculated as: $\text{circularity index} = 4\pi \times (\text{leaf surface}/\text{perimeter}^2)$, ranging from 0 (infinitely narrow leaves) to 1 (perfectly circular leaves). In addition, we scored rosette and stem leaf serration categorically (no serration, little serration, much serration).

Flowering phenotype and spines

At the seed set stage, we measured the length of the main flowering stem from the first node at the base of the inflorescence until the top. In addition, we scored the inflorescence shape (spike-like: most capitula in a horizontal plane, or pyramid-like: capitula of main stem higher than those of axillary branches), the position of the involucre bracts of capitula that had set seed (erect, erect/reflexed, or completely reflexed), and the presence of spines on the stem base as well as on the stem leaves (no spines, few spines, or many spines).

Seed phenotype and seed output

We collected 15 capitula for seed (or the achene fruit to be precise) morphology and seed output measurements. We measured seed length and width, and also measured pappus length and width of 5 randomly selected seeds. The total number of seeds was counted for the 15 capitula to calculate the average number of seeds per capitulum. In order to calculate the total seed output, we counted the number of reproductive basal side shoots and the number of branches from the base of the main flowering stem to the top. Subsequently, we estimated the total number of capitula and the total seed output following Hooftman et al. (2005, eqn 1). Subsequently, the total seed output was calculated by multiplying the number of capitula and the average number of seeds per capitulum.

Statistical analysis

All statistical analyses were performed in PASW Statistics 17.0 (SPSS Inc. 2009), with the exceptions mentioned earlier. Several trait values were transformed to normalize the distribution and improve QTL

Table 1 Traits examined in a *L. sativa* ‘Salinas’ × *Lactuca serriola* recombinant inbred lines population

Plant stage	Trait	Abbreviation	Evaluation method	Type
Seedling	Germination time (days)	GT	No. of days when 50 % of seeds have germinated	Development
30 days after sowing	Rosette leaf length (cm)	RLL	Average distance from the base to leaf tip of four biggest leaves	Leaf shape
	Rosette leaf width (cm)	RLW	Average maximum width of four biggest leaves	Leaf shape
	Rosette leaf surface (cm ²)	RLSU	Surface area of the biggest rosette leaf determined by scanning the leaf	Leaf shape
	Rosette leaf serration	RLSE	No serration, little serration, much serration	Leaf shape
	Rosette leaf circularity	RLC	Proportion of serration of the biggest rosette leaf determined by scanning the leaf and calculated by $4\pi \times (\text{rosette leaf surface/perimeter}^2)$	Leaf shape
	Plant height (cm)	PH30	Length of main stem at 30 days after sowing, values log-transformed	Development
	Proportion of stem leaves (%)	PSL30	No. of stem leaves divided by the total no. of rosette and stem leaves at 30 days, values angular transformed	Development
60 days after sowing	Number of stem leaves (count)	SL60	No. of stem leaves at 60 days after sowing	Development
	Plant height (cm)	PH60	Length of main stem at 60 days after sowing, values log-transformed	Development
Flowering	Days to first flower (day)	FLD	No. of days from sowing to flowering of first flower, values log-transformed	Development
	Flower production	FLP	No flowering or flowering	Development
	Stem leaf length (cm)	SLL	Distance from the base to leaf tip of two leaves halfway the main stem	Leaf shape
	Stem leaf width (cm)	SLW	Maximum width of two leaves halfway the main stem	Leaf shape
	Stem leaf surface (cm ²)	SLSU	Surface area of a stem leaf halfway the main stem determined by scanning the leaf	Leaf shape
	Stem leaf serration	SLSE	No serration, little serration, much serration	Leaf shape
	Stem leaf circularity	SLC	Proportion of serration of a stem leaf halfway the main stem determined by scanning the leaf and calculated by $4\pi \times (\text{stem leaf surface/perimeter}^2)$	Leaf shape
Seed set	Days to first seed (day)	SSD	No. of days from sowing to seed set of first seed, values log-transformed	Development
	No. of stem leaves (count)	SLSS	Total no. of stem leaves	Development
	Plant height (cm)	PHSS	Length of main stem	Development
	No. of reproductive basal shoots (count)	SHN	No. of basal side shoots which have flower buds, flowers and/or seed heads	Seed output
	No. of branches main inflorescence (count)	BRN	No. of branches counted from the base of the main inflorescence to the top	Seed output
	Inflorescence shape	IS	Spike or pyramid	Flowering phenotype
	Inflorescence length (cm)	IL	Length of main inflorescence from the first node at the base until the top	Flowering phenotype
	Involucral bracts	IB	Erect, erect/reflexed, or reflexed	Flowering phenotype
	Stem thickness (cm)	STT	Diameter of the main stem 10 cm above the soil	Development
	Stem leaf spines	SLSP	No spines, few spines or many spines	Spines
Stem base spines	SBSP	No spines, few spines or many spines	Spines	

Table 1 continued

Plant stage	Trait	Abbreviation	Evaluation method	Type
After seed set	Seed length (mm)	SDL	Maximum distance from top to bottom of the seed	Seed phenotype
	Seed width (mm)	SDW	Maximum distance from side to side of the seed	Seed phenotype
	Seed weight (mg)	SDWT	Average weight of 50 seeds	Seed phenotype
	Pappus length (mm)	PL	Length of the stalk	Seed phenotype
	Pappus width (mm)	PW	Maximum distance between the outer tips of the radiating hairs	Seed phenotype
	No. of seeds per capitulum (count)	SDC	Average no. of seeds per capitulum based on 15 collected capitula	Seed output
	Total no. capitula	TC	Total no. of capitula developed, calculation following Hooftman et al. (2005)	Seed output
	Seed output	SDO	Total no. of seeds produced, calculation following Hooftman et al. (2005); values square-root-transformed	Seed output
	Survival until reproduction	SUR	No seeds formed or seed formation	Seed output

analyses. The fraction number of stem leaves at 30 days (relative to the total number of rosette and stem leaves) was arcsine-square-root-transformed. Plant height, days until first flowering, and days until seed set were log-transformed. Estimated seed output values ranged from 242 to 62,847 seeds; these values were square-root-transformed. For all other traits the data was normally distributed and hence, we used the trait values for the QTL analysis. We performed a Principal Component Analysis (PCA) using a Varimax rotation method to summarize the variation among RILs; missing values were replaced by the mean. The loading values of the RILs for the first two axes were saved and also used in the QTL analysis.

Quantitative trait loci analysis

The genetic map and marker data of the RILs, we used to conduct the QTL analysis, are publicly available from The Compositae Genome Project website (<http://compgenomics.ucdavis.edu>). The genetic map consisted of 1,513 predominantly AFLP and EST markers distributed over the nine chromosomal linkage groups (<http://cgpdb.ucdavis.edu/GeneticMapViewer/display/>; map version: RIL_MAR_2007_ratio). All QTL analyses were performed with Composite Interval Mapping (CIM) in QTL Cartographer version 2.5.008 (Wang et al. 2010). Tests for the presence of a

QTL were performed at 2 cM intervals using a 10 cM window and five background cofactors, which were selected via a forward and backward stepwise regression method. Statistical significance threshold values ($\alpha = 0.05$) for declaring the presence of a QTL were estimated from 1000 permutations (Churchill and Doerge 1994; Doerge and Churchill 1996). One-LOD support intervals and additive effects were calculated from the CIM results. The linkage map and QTL were drawn with MapChart 2.2 (Voorrips 2002).

The strength of QTL was described following Burke et al. (2002) where similar potentially multi-genic traits for sunflower were measured. In Burke et al. (2002) more than 25 % explained variance was considered as a major effect QTL and less than 10 % as a minor effect QTL.

Results

We identified 71 QTL for 37 traits and an additional five QTL when summarizing these traits in PCA axes (Table 2; Fig. 1). The range of Phenotypic Variation Explained (PVE) per QTL varied from 7.9 to 69.6 %. The majority of QTL was of intermediate effect (PVE between 10 and 25 %; definitions of QTL strength as defined by Burke et al. 2002). Only seven QTL were of

minor effect ($PVE < 10\%$) and fourteen QTL had a major effect ($PVE > 25\%$).

Quantitative trait loci were distributed over all nine linkage groups. For each trait one to six QTL were detected (mean = 1.9), except for total inflorescence length and number of seeds per capitulum, for which no QTL were detected. The 1-LOD support intervals ranged from 0.3 cM to 5.9 cM (mean = 2.6 cM). Although QTL were distributed over all linkage groups, the majority of QTL clustered with at least one other QTL. Only 17 (out of 71) QTL were uniquely located.

This high level of clustering occurred mainly in two regions on linkage group (LG) 3 and LG7 (Fig. 1), where nine out of the fourteen major QTL are localized. Additional major QTL were found on LG1 for rosette leaf circularity and on LG6 for involucre bracts. The cluster on LG3 is mainly determined by leaf shape and seed output traits. At this location, four major QTL co-localize, including rosette and stem leaf serration, stem leaf circularity, and total number of capitula. This region is also supported by QTL of intermediate effect for shoot number and total seed output. These same traits have the highest loading scores for the second PCA-axis that explained 10% of the variation (Table 2). A major PCA-2 QTL ($PVE = 29.7\%$) indeed co-localizes with the clustering on LG3, confirming this region as a major region for leaf shape and seed output traits. Another trait with a high loading score for the second PCA axis is stem base spines. Not surprisingly, another PCA-2 QTL co-localizes with two major spine-QTL found on LG5.

In contrast, QTL and PCA results show that the clustering on LG7 is the most important region for growth traits, where peak values of five major QTL co-localize within 9 cM. This includes proportion of stem leaves after 30 days, plant height after 60 days, days to first flower, days to first seed, and plant height at seed set (Table 2). These traits, except plant height at seed set, also showed the highest loading scores for the first PCA axis that explained 18% of the variation. A major PCA-1 QTL ($PVE = 35.2\%$) confirms this clustering. Therefore, we consider this the major region for growth related and earliness traits.

For 21 traits, we identified more than one QTL. Of these, QTL effects of seven traits were always increased by the wild parent allele. These included proportion of stem leaves after 30 days, plant height at 60 days, stem leaf width, stem thickness, and seed output traits such as

seed length, seed production, and seed output. In contrast, four traits (rosette leaf circularity, stem leaf circularity, days to first flower, and days to first seed) were always increased by the crop parent allele. This shows that the wild parent alleles cause a faster development with an earlier bolting and flowering time, and an increased seed output.

Ten traits showed opposing effects with some QTL where crop alleles resulted in higher values for a trait, but one or more others where they led to lower values. These were all QTL of intermediate effects, including a number of leaf shape traits, but also seed output traits such as seed weight, shoot number, and total number of capitula. Apparently, for each of these traits the crop allele contributed to an increase in trait values at one location, but at another location the increase came from the wild parental allele. Whether there are selective advantages of the direction of the effects could of course not be determined under greenhouse conditions.

Discussion

We identified 71 domestication- and fitness-related QTL for this lettuce cross of which 14 were of major effect ($PVE > 25\%$). Of the remaining QTL, the majority was of intermediate effect (50) and only seven were of minor effect ($PVE < 10\%$). Cultivated Crisphead lettuce appears to have been selected for no spines, erect involucre bracts, round almost circular leaves instead of serrate leaves, and a delay in bolting beneficial for vegetative biomass production (de Vries 1997). Several QTL of major effect could be linked directly to these assumed selection regimes. For instance, two closely linked major QTL for rosette leaf circularity on linkage group (LG) 1 are directly related to the cultivated allele inducing rounder leaves. On LG5, we found two major QTL for stem base spines and stem leaf spines and as expected, the wild allele induced spine production. Similarly, for involucre bracts we found one major QTL (LG6). Here, the cultivated parental allele induced erect involucre bracts that are important to prevent shattering of the seeds (achenes).

Two regions were identified—on LG3 and LG7—that have significant clustering of major as well as intermediate QTL. Our data show that the region on LG3 is involved in leaf shape and seed output traits,

Table 2 QTL detected by composite interval mapping in a *Lactuca sativa* ‘Salinas’ × *Lactuca serriola* recombinant inbred lines population and Principal Component Analysis (PCA) loadings for the first two axes

Trait	LG	Position (cM)	LOD	1-LOD interval	Additive effect	PVE (%)	Threshold 0.05	PCA-1	PCA-2
GT	7	65.9	3.9	65.4–67.7	0.39	12.1	3.2	−0.04	−0.00
RLL	7	17.4	5.1	15.5–18.5	−1.65	17.1	3.3	0.32	0.10
RLW	1	89.0	3.7	88.3–91.0	0.75	10.0	3.3	0.12	0.15
	3	37.2	5.5	36.8–40.9	−0.99	15.2			
	3	47.0	4.8	46.3–47.8	−0.90	13.4			
	7	97.8	7.1	97.1–99.0	1.03	18.5			
	7	114.6	5.1	112.8–114.7	0.89	13.9			
	8	132.0	3.4	130.9–132.9	0.71	7.9			
RLSU	1	1.7	3.8	1.6–1.9	−13.37	8.7	3.2	0.31	0.13
	1	89.0	7.8	87.9–89.7	20.53	20.4			
	3	38.8	5.7	37.9–41.0	−18.06	15.0			
RLSE	3	42.9	14.3	41.6–43.5	−0.48	34.7	3.5	−0.02	0.54
	7	97.8	7.6	96.6–100.0	0.31	16.4			
RLC	1	72.4	10.0	71.9–72.4	0.05	29.6	3.4	−0.03	−0.27
	1	76.6	12.7	76.4–77.3	0.05	35.9			
PH30	7	17.4	8.0	15.5–21.0	−0.22	23.7	4.2	0.63	0.03
PSL30	6	129.9	3.9	129.7–132.3	−0.20	8.8	3.5	0.74	0.20
	7	18.5	15.7	18.4–20.3	−0.45	43.9			
	8	33.5	3.8	32.2–34.3	−0.21	9.0			
SL60	7	19.2	5.5	18.7–22.4	−28.55	16.7	3.4	0.40	0.60
PH6	2	111.3	4.1	110.5–116.4	−0.18	8.7	3.5	0.73	0.32
	6	130.8	7.3	129.9–132.8	−0.26	17.4			
	7	21.9	15.3	19.6–22.4	−0.39	41.5			
	8	23.4	4.1	22.1–24.7	−0.19	8.9			
FLD	2	106.8	4.1	105.5–107.5	0.05	12.4	3.4	−0.82	−0.21
	7	19.2	11.0	18.5–19.9	0.08	35.0			
FLP	9	1.0	4.8	0.0–1.4	−0.18	17.0	3.4	0.10	0.05
SLL	2	17.8	3.7	15.9–18.1	1.62	11.3	3.4	0.31	−0.13
	7	109.4	4.8	108.8–109.8	−1.81	14.1			
	9	0.2	3.8	0.0–2.1	1.78	12.9			
SLW	3	37.2	6.7	36.7–39.0	−1.12	23.6	3.3	0.22	0.05
	7	7.2	3.4	6.2–7.4	−0.77	11.6			
SLSU	6	119.9	4.0	117.5–120.5	−12.44	19.3	3.4	0.08	−0.35
SLSE	3	42.9	16.2	41.6–45.8	−0.55	52.9	3.5	0.02	0.60
SLC	1	38.9	4.6	37.6–40.2	0.04	12.4	3.8	0.12	−0.37
	2	136.3	6.2	134.8–138.0	0.05	15.9			
	3	45.4	11.6	45.3–46.6	0.07	37.9			
SSD	1	30.8	3.9	30.7–31.6	0.04	13.9	3.3	−0.88	−0.25
	2	106.8	5.1	105.5–107.1	0.05	15.6			
	7	19.2	7.0	17.9–19.9	0.07	27.6			
SLSS	9	55.9	4.3	55.0–56.3	−31.92	16.5	3.5	0.18	0.58
PHSS	1	79.2	3.4	78.2–80.8	−0.04	13.0	3.4	−0.45	0.11
	6	119.9	3.5	118.2–124.1	−0.04	11.0			
	7	12.9	6.4	11.3–15.2	0.05	26.9			

Table 2 continued

Trait	LG	Position (cM)	LOD	1-LOD interval	Additive effect	PVE (%)	Threshold 0.05	PCA-1	PCA-2
SHN	3	9.2	3.5	8.1–11.5	1.18	12.2	3.4	0.43	0.74
	3	38.8	3.9	37.2–40.9	−1.25	15.1			
	7	18.4	4.9	16.4–19.2	−1.34	17.4			
BRN	7	25.1	5.5	23.3–25.9	2.11	16.6	3.3	−0.41	0.74
IS	4	154.9	3.9	154.3–157.0	−0.18	13.9	3.9	0.03	−0.23
IL	nd							0.16	0.54
IB	1	77.6	5.4	77.1–78.8	0.33	11.5	3.6	−0.06	0.16
	6	5.6	21.3	4.3–7.7	−0.75	61.3			
STT	5	91.0	5.3	90.5–91.6	−0.15	17.6	3.4	−0.06	−0.34
	7	111.9	3.6	110.9–113.5	−0.12	11.6			
SLSP	5	146.8	17.4	146.1–148.1	−0.60	49.7	3.9	−0.01	−0.30
SBSP	5	146.8	23.4	146.1–149.7	−0.81	69.6	3.5	−0.07	−0.46
SDL	1	69.3	4.3	67.9–73.4	−0.15	16.2	3.3	0.63	0.07
	7	10.2	6.3	8.7–12.1	−0.18	23.2			
SDW	8	19.5	5.8	17.7–21.1	−0.07	21.7	3.4	0.52	−0.00
	9	72.3	3.8	71.0–76.3	0.06	14.2			
SDWT	1	26.1	4.1	24.7–26.5	−5.16	17.0	3.4	0.68	−0.17
PL	4	0.4	3.9	0.0–1.9	−0.20	13.0	3.4	−0.35	−0.01
PW	2	134.8	5.9	132.6–137.3	−0.34	21.8	3.5	0.09	−0.01
	4	7.6	4.9	7.4–10.3	−0.30	17.0			
	4	138.5	3.8	137.5–139.2	+0.24	10.9			
SDC	nd							−0.67	0.32
TC	3	9.2	4.8	8.1–9.7	260.21	20.7	3.4	0.26	0.81
	3	38.8	7.6	37.2–40.0	−287.75	27.5			
SDO	3	38.8	4.5	37.2–40.9	−18.98	16.7	3.3	−0.44	0.63
	4	143.3	3.3	141.4–144.2	−15.48	11.4			
	7	105.2	3.6	103.5–106.7	−15.43	11.8			
SUR	6	23.9	4.6	23.3–25.3	−0.15	13.1	3.6	0.07	0.08
	7	77.0	3.4	75.3–78.0	−0.13	9.9			
PCA-QTL							Total variance explained (%)		
PCA1	1	30.8	5.7	29.8–31.6	−0.37	13.4	3.3	18.04	
	5	117.0	3.4	112.3–121.2	0.36	12.7			
	7	18.4	13.8	17.4–19.2	−0.60	35.2			
PCA2	3	38.8	10.6	37.2–41.0	−0.57	29.7	3.4	10.67	
	5	142.3	3.5	140.1–144.5	0.29	8.3			

The additive effect indicates which parental allele causes an increase in the trait value. Positive values indicate that the crop-type (*L. sativa*) allele increases trait values, whereas negative values indicate that the wild-type (*L. serriola*) allele increases trait values. For trait abbreviations, see Table 1; *nd* no QTL detected, *cM* centiMorgan, *LOD* Logarithm of Odds, *1-LOD interval* region on both sides of the QTL peak that corresponds to a decrease of one LOD, *PVE* Percentage Variation Explained, *Threshold* the significance threshold determined by permutation tests

whereas the clustering on LG7 is mainly involved in developmental traits, such as height after 30 days and days to first flower, supported by some intermediate

leaf shape and seed output QTL. PCA-QTL co-localize and confirm these clustering regions. Based on marker comparisons between different linkage

main phenotypic transitions from wild-type to crop. This is comparable with the pattern found in the majority of crops in which phenotypic differentiation is mainly caused by a few genomic regions of major effect (Burger et al. 2008; Hancock 2005). However, the majority of QTL in this study was of intermediate effect, which is more similar to the patterns found in sunflower (Burke et al. 2002; Wills and Burke 2007).

A possible statistical explanation might be that minor QTL went undetected as a result of small sample size (Beavis 1998), biasing the results towards QTL of large effect. However, the large amount of located intermediate QTL suggests that this distribution of QTL is a real feature of the genetic architecture of lettuce. The main phenotypic differentiation between wild and cultivated lettuce seems to be the result of a few genomic regions with major QTL, plus many QTL of intermediate effect. Whether this is the result of the additive effects of individual genes or through epistatic interactions is to be studied (Uwimana 2011).

Clustered domestication QTL

A major trend prevalent across many crop species is that domestication traits are not distributed randomly throughout the genome, but are often found clustered together in specific genomic regions (Burger et al. 2008; Ross-Ibarra 2005). Indeed we identified two such clustering regions. Similarly, Zhang et al. (2007) found several clusters of QTL for shelf life, leaf biophysical, and developmental traits for lettuce. The two potential explanations are that these clustering patterns result from pleiotropic effects of a single gene affecting different traits at once or from physical linkage among multiple genes that were individually selected, or have been integrated in the crop genome due to linkage drag (Burger et al. 2008; Ross-Ibarra 2005).

A QTL spanning 10 cM may contain hundreds of genes (Collard et al. 2005; Mauricio 2001), making distinguishing between both possibilities very difficult and as yet there is no information on the identity of (candidate) genes in these regions. For the different traits in the cluster at LG3 a pleiotropic explanation is not obvious. However, for the second cluster (LG7) the majority of traits seem involved in speed of development, such as height at 30 days and days to first flower. We consider it plausible that at LG7 the

clustering of these traits, together with those for leaf shape, branching, and seed output, is caused by pleiotropic effects from a common major gene for earliness of flowering. Similar results have been found in dry bean (Koinange et al. 1996), where a gene for earliness also affected branching patterns, number of days from flowering to fruiting, and pod number. As a more recent example, in *Arabidopsis* a major flowering gene was found to be involved in germination as well (Chiang et al. 2009).

Direction of effects

QTL research on crop species has revealed that in many instances alleles show opposing effects, meaning a crop allele contributing to a wild phenotype or vice versa (Ross-Ibarra 2005). Our results show that all allelic effects of major QTL were in the expected direction. However, our results show for many QTL with intermediate effects traits were increased by crop and wild alleles alike; such opposing effects have been found in previous lettuce research as well (Argyris et al. 2005; Zhang et al. 2007). We found that crop alleles, for example, induced higher germination rates, and more branches and hence, capitula that might be beneficial in the wild habitat at least under some environmental circumstances (Baack et al. 2008; Kaga et al. 2008).

The genetic architecture of lettuce thus concurs mainly with the main domestication trends observed in other species. The presence of many QTL of intermediate effect in combination with the appearance of opposing effects suggests that domestication of lettuce went readily and could reflect multiple domestication events, as suggested for sunflower (Burke et al. 2002). However, the existence of alleles with opposing effects seems to be a general pattern even for crops that are believed to be domesticated through single selection events (Ross-Ibarra 2005). Gross and Olsen (2010) showed that there are multiple pathways to make domesticated plants and that the inferences made from genetic evidence and one cross is not always straightforward. First, QTL results can be very cultivar specific (Mercer et al. 2006). The formation of a tight dense head and delay in bolting are the result of modern breeding efforts (de Vries 1997), and *L. sativa* ‘Salinas’ is one of the latest bolting and flowering ones (Simonne et al. 2002). Second, these results do need to be verified in the field, as results for controlled

greenhouse and environmentally variable field conditions may differ due to Genotype by Environment interactions (Hails and Morley 2005; Weinig et al. 2002). Moreover, hybrid fitness and selective forces may vary across different environments and in different seasons (Chapman and Burke 2006). Further research on the domestication of lettuce is needed to be able to further disentangle the genetic basis and historic selection pressures.

Implications for crop breeding

The enormous increase in molecular tools, such as genetic maps and markers, is only expected to continue as more and more species are being sequenced (Collard and Mackill 2008). With the introduction of new molecular techniques and analysis tools, crop breeding has entered a new phase and rapid progress is being made in the development of new transgenic cultivars (Vaughan et al. 2007) that have higher yields, increased resistance to diseases and herbicides, or increased tolerance to abiotic stresses, such as salt or drought (Campos et al. 2004; Cuartero et al. 2006; Warwick et al. 2008). Current lettuce breeding is strongly focused on utilization and exploitation of wild lettuce relatives (Lebeda et al. 2009); however, transgenic research is also ongoing (Park et al. 2005). To our knowledge, there are no genetically modified lettuce cultivars in production at the moment. Our results indicate that when they are produced the likelihood of escape will depend on the location of insertion (Stewart et al. 2003).

The results of our study also shed some light on potential mitigation strategies to prevent the escape of transgenes by introgression to wild relatives. A transgene placed in close linkage with a gene or genomic block that causes a lower fitness in the wild habitat is likely to be purged from the wild populations (Gressel 1999; Stewart et al. 2003). Techniques for targeted insertion of transgenes to specific regions in the genome are currently being developed (Lombardo et al. 2011; Nandy and Srivastava 2011; Shukla et al. 2009). Based on these greenhouse results, the clustering regions on LG3 and LG7 may be considered as such possible insertion sites. At these sites, crop alleles invariably contributed to e.g. lower amount of basal side shoots and lower seed output, leading to lower hybrid fitness and chances of introgression. Moreover, if the clustering at LG7 is in fact the result of a major

gene for earliness, crop alleles will induce a delay in bolting which might be deleterious in the wild habitat. Indeed, in a recent field study it was shown that a crop genomic background at the clustering region on LG7 provided a negative fitness effect to hybrid individuals (Hartman et al. 2012). However, no fitness effect was detected for the clustering region on LG3. QTL research, such as ours, will therefore prove as a valuable tool to give a first indication of regions in the crop genome less likely to introgress before any transgene is inserted (Kwit et al. 2011; Stewart et al. 2003), but such results do have to be verified in the field under selective conditions (Hartman et al. 2012).

Acknowledgments We are grateful to the laboratory of R. W. Michelmore (UC Davis) for graciously providing seeds from their RIL collection to us and for discussions. The Compositae Genome Project (<http://compgenomics.ucdavis.edu>) that developed the RILs is supported by the NSF Plant Genome Program award #0820451. We like to thank the assistants and technical staff involved especially Rob Bregman, Louis Lie, Ludek Tikovsly, Thijs Hendrix, and Harold Lemereis. This study is funded by the Netherlands Organization for Scientific Research (NWO) as part of the ERGO program (838.06.042).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Abramoff MD, Magalhaes PJ, Ram SJ (2004) Image processing with image. *J Biophotonics Int* 11:36–42
- Argyris J, Truco MJ, Ochoa O, Knapp SJ, Still DW, Lenssen GM, Schut JW, Michelmore RW, Bradford KJ (2005) Quantitative trait loci associated with seed and seedling traits in *Lactuca*. *Theor Appl Genet* 111:1365–1376
- Baack EJ, Sapir Y, Chapman MA, Burke JM, Rieseberg LH (2008) Selection on domestication traits and quantitative trait loci in crop-wild sunflower hybrids. *Mol Ecol* 17: 666–677
- Beavis W (1998) QTL analyses: power, precision and accuracy. In: Paterson AH (ed) *Molecular dissection of complex traits*. CRC Press, Boca Raton, pp 145–161
- Burger JC, Chapman MA, Burke JM (2008) Molecular insights into the evolution of crop plants. *Am J Bot* 95:113–122
- Burke JM, Tang S, Knapp SJ, Rieseberg LH (2002) Genetic analysis of sunflower domestication. *Genetics* 161: 1257–1267
- Burke JM, Burger JC, Chapman MA (2007) Crop evolution: from genetics to genomics. *Curr Opin Genet Dev* 17: 525–532

- Cai HW, Morishima H (2002) QTL clusters reflect character associations in wild and cultivated rice. *Theor Appl Genet* 104:1217–1228
- Campos H, Cooper A, Habben JE, Edmeades GO, Schussler JR (2004) Improving drought tolerance in maize: a view from industry. *Field Crops Res* 90:19–34
- Chapman MA, Burke JM (2006) Letting the gene out of the bottle: the population genetics of genetically modified crops. *New Phytol* 170:429–443
- Chiang GCK, Barua D, Kramer EM, Amasino RM, Donohue K (2009) Major flowering time gene, flowering locus C, regulates seed germination in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 106:11661–11666
- Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative mapping. *Genetics* 138:963–971
- Collard BCY, Mackill DJ (2008) Marker-assisted selection: an approach for precision plant breeding in the twenty-first century. *Philos Trans R Soc Lond B Biol Sci* 363:557–572
- Collard BCY, Jahufer MZZ, Brouwer JB, Pang ECK (2005) An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica* 142:169–196
- Cuartero J, Bolarin MC, Asins MJ, Moreno V (2006) Increasing salt tolerance in the tomato. *J Exp Bot* 57:1045–1058
- D'Andrea L, Felber F, Guadagnuolo R (2008) Hybridization rates between lettuce (*Lactuca sativa*) and its wild relative (*L. serriola*) under field conditions. *Environ Biosafety Res* 7:61–71
- D'Andrea L, Broennimann O, Kozłowski G, Guisan A, Morin X, Keller-Senften J, Felber F (2009) Climate change, anthropogenic disturbance and the northward range expansion of *Lactuca serriola* (Asteraceae). *J Biogeogr* 36:1573–1587
- de Vries IM (1990) Crossing experiments of Lettuce cultivars and species (*Lactuca* sect. *Lactuca*, Compositae). *Plant Syst Evol* 171:233–248
- de Vries IM (1997) Origin and domestication of *Lactuca sativa* L. *Genet Resour Crop Evol* 44:165–174
- Dechaine JM, Burger JC, Chapman MA, Seiler GJ, Brunick R, Knapp SJ, Burke JM (2009) Fitness effects and genetic architecture of plant-herbivore interactions in sunflower crop-wild hybrids. *New Phytol* 184:828–841
- Dempewolf H, Rieseberg LH, Cronk QC (2008) Crop domestication in the Compositae: a family-wide trait assessment. *Genet Resour Crop Evol* 55:1141–1157
- Doebley J, Stec A (1993) Inheritance of the morphological differences between maize and teosinte—comparison of results for 2 F2 populations. *Genetics* 134:559–570
- Doerge RW, Churchill GA (1996) Permutation tests for multiple loci affecting a quantitative character. *Genetics* 142:285–294
- Doganlar S, Frary A, Daunay MC, Lester RN, Tanksley SD (2002) Conservation of gene function in the Solanaceae as revealed by comparative mapping of domestication traits in eggplant. *Genetics* 161:1713–1726
- Ellstrand NC (2003) Dangerous Liaisons? When cultivated plants mate with their wild relatives. John Hopkins University Press, Baltimore
- Ellstrand NV, Heredia SM, Leak-Garcia JA, Heraty JM, Burger JC, Yao L, Nohzadeh-Malakshah S, Ridley CE (2011) Crops gone wild: evolution of weeds and invasives from domesticated ancestors. *Evol Appl* 3:494–504
- Frary A, Nesbitt TC, Grandillo S, van der Knaap E, Cong B, Liu JP, Meller J, Elber R, Alpert KB, Tanksley SD (2000) fw2.2: a quantitative trait locus key to the evolution of tomato fruit size. *Science* 289:85–88
- Giannino D, Nicolodi C, Testone G, Di Giacomo E, Iannelli MA, Frugis G, Mariotti D (2008) Pollen-mediated trans-gene flow in lettuce (*Lactuca sativa* L.). *Plant Breed* 127:308–314
- Gressel J (1999) Tandem constructs: preventing the rise of superweeds. *Trends Biotechnol* 17:361–366
- Gross BL, Olsen KM (2010) Genetic perspectives on crop domestication. *Trends Plant Sci* 15:529–537
- Hails RS, Morley K (2005) Genes invading new populations: a risk assessment perspective. *Trends Ecol Evol* 20:245–252
- Hammer K (1984) Das Domestikationssyndrom. *Kulturpflanze* 32:11–34
- Hancock JF (2005) Contributions of domesticated plant studies to our understanding of plant evolution. *Ann Bot* 96:953–963
- Harlan JR (1992) Crops and man. American Society of Agronomy, Madison, WI
- Hartman Y (2012) Genomic regions under selection in crop-wild hybrids of Lettuce, implications for crop breeding and environmental risk assessment. PhD thesis, University of Amsterdam, The Netherlands
- Hartman Y, Hooftman DAP, Uwimana B, van de Wiel CCM, Smulders MJM, Visser RGF, van Tienderen PH (2012) Genomic regions in crop-wild hybrids of lettuce are affected differently in different environments: implications for crop breeding. *Evol Appl*. doi:10.1111/j.1752-4571.2012.00240.x
- Hooftman DAP, Oostermeijer JGB, Jacobs MMJ, Den Nijs HCM (2005) Demographic vital rates determine the performance advantage of crop-wild hybrids in lettuce. *J Appl Ecol* 42:1086–1095
- Hooftman DAP, Oostermeijer JGB, den Nijs JCM (2006) Invasive behaviour of *Lactuca serriola* (Asteraceae) in the Netherlands: spatial distribution and ecological amplitude. *Basic Appl Ecol* 7:507–519
- Hooftman DAP, Jong MJD, Oostermeijer JGB, Den Nijs H (2007) Modelling the long-term consequences of crop-wild relative hybridization: a case study using four generations of hybrids. *J Appl Ecol* 44:1035–1045
- Hooftman DAP, Hartman Y, Oostermeijer JGB, Den Nijs HJCM (2009) Existence of vigorous lineages of crop-wild hybrids in Lettuce under field conditions. *Environ Biosafety Res* 8:203–217
- Isemura T, Kaga A, Tomooka N, Shimizu T, Vaughan DA (2010) The genetics of domestication of rice bean, *Vigna umbellata*. *Ann Bot* 106:927–944
- Johnson WC, Jackson LE, Ochoa O, van Wijk R, Peleman J, St Clair DA, Michelmore RW (2000) Lettuce, a shallow-rooted crop, and *Lactuca serriola*, its wild progenitor, differ at QTL determining root architecture and deep soil water exploitation. *Theor Appl Genet* 101:1066–1073
- Kaga A, Isemura T, Tomooka N, Vaughan DA (2008) The genetics of domestication of the azuki bean (*Vigna angularis*). *Genetics* 178:1013–1036

- Koinange EMK, Singh SP, Gepts P (1996) Genetic control of the domestication syndrome in common bean. *Crop Sci* 36:1037–1045
- Kwit C, Moon HS, Warwick SI, Stewart CN (2011) Transgene introgression in crop relatives: molecular evidence and mitigation strategies. *Trends Ecol Evol* 29:284–293
- Lebeda A, Dolezalova I, Kristkova E, Mieslerova B (2001) Biodiversity and ecogeography of wild *Lactuca* spp. in some European countries. *Genet Resour Crop Evol* 48:153–164
- Lebeda A, Dolezalova I, Kristkova E, Kitner M, Petrzalova I, Mieslerova B, Novotna A (2009) Wild *Lactuca* germplasm for lettuce breeding: current status, gaps and challenges. *Euphytica* 170:15–34
- Lindqvist K (1960a) Cytogenetic studies in the *serriola* group of *Lactuca*. *Hereditas* 46:75–151
- Lindqvist K (1960b) Inheritance studies in Lettuce. *Hereditas* 46:387–470
- Lombardo A, Cesana D, Genovese P et al (2011) Site-specific integration and tailoring of cassette design for sustainable gene transfer. *Nat Methods* 8:861–869
- Mauricio R (2001) Mapping quantitative trait loci in plants: uses and caveats for evolutionary biology. *Nat Rev Genet* 2:370–381
- Mercer KL, Wyse DL, Shaw RG (2006) Effects of competition on the fitness of wild and crop-wild hybrid sunflower from a diversity of wild populations and crop lines. *Evolution* 60:2044–2055
- Nandy S, Srivastava V (2011) Site-specific gene integration in rice genome mediated by the FLP-FRT recombination system. *Plant Biotechnol J* 9:713–721
- Park BJ, Liu ZC, Kanno A, Kameya T (2005) Increased tolerance to salt- and water-deficit stress in transgenic lettuce (*Lactuca sativa* L.) by constitutive expression of LEA. *Plant Growth Regul* 45:165–171
- Poncet V, Lamy F, Devos KM, Gale MD, Sarr A, Robert T (2000) Genetic control of domestication traits in pearl millet (*Pennisetum glaucum* L., Poaceae). *Theor Appl Genet* 100:147–159
- Prince SD, Marks MK, Carter RN (1979) Induction of Flowering in Wild Lettuce (*Lactuca serriola* L.). *New Phytol* 81:265–277
- Prudent M, Causse M, Genard M, Tripodi P, Grandillo S, Bertin N (2009) Genetic and physiological analysis of tomato fruit weight and composition: influence of carbon availability on QTL detection. *J Exp Bot* 60:923–937
- Purugganan MD, Fuller DQ (2009) The nature of selection during plant domestication. *Nature* 457:843–848
- Rieseberg LH, Widmer A, Arntz AM, Burke JM (2002) Directional selection is the primary cause of phenotypic diversification. *Proc Natl Acad Sci USA* 99:12242–12245
- Ross-Ibarra J (2005) Quantitative trait loci and the study of plant domestication. *Genetica* 123:197–204
- Shukla VK, Doyon Y, Miller JC et al (2009) Precise genome modification in the crop species *Zea mays* using zinc-finger nucleases. *Nature* 459:437–441
- Simonne A, Simonne E, Eitenmiller R, Coker CH (2002) Bitterness and composition of lettuce varieties grown in the southeastern United States. *HortTechnology* 12:721–726
- SPSS Inc. (2009) PASW Statistics 17.0 Command Syntax Reference. SPSS Inc., Chicago
- Stewart CN, Halfhill MD, Warwick SI (2003) Transgene introgression from genetically modified crops to their wild relatives. *Nat Rev Genet* 4:806–817
- Thompson RC, Whitaker TW, Kosar WF (1941) Interspecific genetic relationships in *Lactuca*. *J Agric Res* 63:91–107
- Uwimana B (2011) A genetic analysis of the introgression process from cultivated lettuce (*Lactuca sativa* L.) to wild prickly lettuce (*L. serriola* L.). PhD thesis, Wageningen University, The Netherlands
- Vaughan DA, Balazs E, Heslop-Harrison JS (2007) From crop domestication to super-domestication. *Ann Bot* 100:893–901
- Visscher PM, Hill WG, Wray NR (2008) Heritability in the genomics era—concepts and misconceptions. *Nat Rev Gen* 9:255–266
- Voorrips RE (2002) MapChart: software for the graphical presentation of linkage maps and QTLs. *J Hered* 93:77–78
- Wang S, Basten CJ, Zeng ZB (2010) Windows QTL Cartographer 2.5. Department of Statistics, North Carolina State University, Raleigh, NC
- Warwick SI, Legere A, Simard MJ, James T (2008) Do escaped transgenes persist in nature? The case of an herbicide resistance transgene in a weedy *Brassica rapa* population. *Mol Ecol* 17:1387–1395
- Weinig C, Ungerer MC, Dorn LA, Kane NC, Toyonaga Y, Halldorsdottir SS, Mackay TFC, Purugganan MD, Schmitt J (2002) Novel loci control variation in reproductive timing in *Arabidopsis thaliana* in natural environments. *Genetics* 162:1875–1884
- Wills DM, Burke JM (2007) Quantitative trait locus analysis of the early domestication of sunflower. *Genetics* 176:2589–2599
- Zhang FZ, Wagstaff C, Rae AM, Sihota AK, Keevil CW, Rothwell SD, Clarkson GJJ, Michelmores RW, Truco MJ, Dixon MS, Taylor G (2007) QTLs for shelf life in lettuce co-locate with those for leaf biophysical properties but not with those for leaf developmental traits. *J Exp Bot* 58:1433–1449