RESEARCH ARTICLE

Analysis of genetic diversity in *Prunus sibirica* L. in inner Mongolia using SCoT molecular markers

Ha Buer · Sa Rula · Zi Yuan Wang · Shu Fang · Yu'e Bai

Received: 24 June 2021 / Accepted: 3 October 2021 / Published online: 3 January 2022
© The Author(s) 2021

Abstract Population genetic diversity contributes to the protection and utilization of germplasm resources, especially via genetic breeding. In the present study, start codon targeted polymorphism (SCoT) molecular markers were used to study the genetic diversity of 278 individuals from 10 *Prunus sibirica* L. populations in Inner Mongolia. A total of 289 polymorphic bands were amplified with 23 SCoT primers, showing a polymorphism percentage of 98.87% and an average of 12.6 polymorphic bands per primer. The SCoT21, SCoT32, and SCoT53 primers amplified up to 17 bands, and the polymorphism percentage was 100%. The minimum number of bands amplified by SCoT25 was 9, and the polymorphism percentage was 90%. Therefore, SCoT molecular markers were shown to be highly polymorphic and suitable for genetic diversity studies of *P. sibirica* in Inner Mongolia. The analysis of molecular variance showed that 39% of the observed genetic differentiation occurred among populations and 61% occurred within populations, indicating that the genetic differentiation within populations was greater than that among populations.

The results of the unweighted pair-group method with an arithmetic cluster analysis, principal coordinate analysis and STRUCTURE analysis were basically the same and divided the 278 individuals from the 10 populations into 2 groups. The results indicated that the efficient SCoT molecular marker-based genetic diversity analysis of *P. sibirica* in Inner Mongolia can provide a reference for *P. sibirica* variety breeding and resource development.

Keyword *Prunus sibirica* · Provenance · SCoT · Molecular marker · Genetic diversity

Introduction

Prunus sibirica L. (Siberian apricot) is a member of the Rosaceae family and is the dominant species on mountain dunes and dry steppes (Yu 1979). *P. sibirica* is an important ecological and economic tree species in China, mainly distributed in Inner Mongolia, Heilongjiang, Jilin, Liaoning, Gansu, Hebei, and Shanxi in China, and the *P. sibirica* forest in Inner Mongolia covers 47.44 hectares.

P. sibirica shows strong cold and drought resistance and high nutritional and medicinal value (Dong et al. 1991). However, the yield differences in *P. sibirica* in China are significant and very unstable because of long-term seed reproduction, delayed breeding, and the influence of climatic factors. In addition, frost can

Ha Buer and Sa Rula These authors contributed equally to this work.

H. Buer · S. Rula · Z. Y. Wang · S. Fang · Y. Bai (✉)
Forestry College, Inner Mongolia Agricultural University,
Xinjian Street 275, Hohhot 010019, Inner Mongolia,
China
e-mail: baiyue@imau.edu.cn

severely reduce the yield of *P. sibirica*, greatly restricting its industrialization (Ma et al. 2007; Yao et al. 2007; Jin et al. 2018). *P. sibirica* shows self-incompatibility, resulting in large genetic differences among individual genetic resources (Li et al. 2011a, b).

Biodiversity refers to all of the variation among living things on earth and is the basis of survival and biological development (Rao et al. 2002). Genetic diversity is an important component of biological diversity and refers to the genetic differentiation of all living organisms (Lu 2018). Dong et al (2018) studied the diversity of 19 quantitative traits of *P. sibirica* from different populations, and the results showed that the coefficient of variation of the economic traits of fruit and kernel were large, and the coefficient of variation of fruit yield was the largest, indicating that *P. sibirica* has rich germplasm resources with great potential for breeding good varieties. Wang et al (2019) studied the seed traits of *P. sibirica* from 19 populations and showed rich genetic variations in seed traits between 19 populations. Li (2014) studied the genetic diversity of *P. sibirica* and *Prunus armeniaca* in North China according to inter-simple sequence repeat (ISSR), sequence-related amplified polymorphism (SRAP), and simple sequence repeat (SSR) analyses. The results showed that the natural population of *P. sibirica* in China exhibits a high level of genetic diversity, showing high genetic differentiation within its populations but low genetic differentiation and moderate gene flow among populations.

The initial SCoT marker was a molecular marker based on a single primer amplification reaction proposed by Collard and Mackill (2009) in rice. It was a novel molecular marker of the target gene. The strategy was to conduct genome amplification with a single primer according to the conservative nature of the ATG translation of the flanking sequences of plant genes, the goal was to reveal the percentage of dominant polymorphic markers in candidate functional gene regions via a procedure with easy operation to identify rich polymorphisms. Compared with random amplified polymorphic DNA (RAPD), amplified fragment length polymorphism (AFLP), SSR, and ISSR markers, SCoT markers have been efficiently produced and linked to traits, making them convenient for use in molecular marker-assisted breeding. SCoT markers have been used in various plant species, such as *Phoenix dactyifera* (Somayeh Saboori et al. 2020),

Mangifera indica (Li Zhou et al. 2020) and *Diospyros kaki*. (Changfei Guana et al. 2020).

Genetic resources for *P. sibirica* are extremely abundant and present great developmental potential, and they can provide an important genetic basis for the improvement of Chinese *P. sibirica* and the breeding of new varieties. In recent years, research on the genetic diversity of *P. sibirica* has mostly been conducted in northeastern and northern China, and *P. sibirica* is mainly distributed in Inner Mongolia. Moreover, few studies (Bao et al. 2016) have been conducted on the genetic diversity of *P. sibirica* in Inner Mongolia. Therefore, we used SCoT molecular markers to analyze the genetic diversity of 278 individuals from 10 populations of *P. sibirica* in Inner Mongolia. The genetic diversity and genetic structure among populations and individuals of *P. sibirica* in Inner Mongolia were revealed. This study provides a scientific basis for the cultivation and exploitation of the abundant genetic resources of *P. sibirica* in Inner Mongolia.

Materials and methods

Plant materials

The materials were obtained from the *P. sibirica* Germplasm Resource Garden of the Inner Mongolia Fine Variety Breeding Center (Fig. 1). A total of 278 individuals were collected from 10 populations (Table 1). Fresh young leaves were placed in a bucket containing liquid nitrogen, brought back to the laboratory, and stored in a -80 °C freezer for further experiments.

DNA extraction

Genomic DNA was extracted from fresh leaves using a Plant DNA Extraction Kit (TIANGEN, China). The extracted DNA was tested for quality and purity with a spectrophotometry at a wavelength of 260/280 nm using a BioPhotometer (Thermo Fisher Scientific, America) and 1% agarose gel electrophoresis (with a nucleic acid stain) and stored in a -20 °C freezer for further experiments.

Fig. 1 Geographic location map **Table 1** Site conditions and sampling information for *P. sibirica* populations

Population Abbreviation	Longitude and Latitude	Elevation (m)	Sample size
KSK	117°54'57"/43°26'49"	1,100	46
WJG	111°02'38"/40°75'73"	1,629	23
KYZ	121°47'65"/45°06'08"	270	26
KZH	122°35'67"/42°93'51"	198	38
ZLA	122°89'96"/46°72'32"	628	22
LC	112°50'39"/40°53'15"	1,731	31
BLY	118°66'51"/43°53'44"	1,050	25
WLS	108°77'63"/40°66'26"	2,322	14
AL	120°06'56"/43°87'22"	900	27
ZLT	120°91'16"/44°55'63"	812	26

SCoT-PCR

The 80 primer sequences were selected by referring to Collard and Mackill (2009) and were synthesized by Shanghai Sangon Biological Engineering Technology and Services. A total of 23 highly polymorphic and repeatable primers were selected from 80 SCoT primers to evaluate the selected *P. sibirica* accessions (Table 2).

A 96-well gradient PCR instrument was used for the amplification reactions. PCR was performed in a 20 μ L volume containing 10 μ L of 2 $\times$ Taq Mix, 1.6 μ L (30 ng/ μ L) of genomic DNA, 0.8 μ L (10 mol·L⁻¹) of each primer, and 7.6 μ L of ddH₂O. SCoT-PCR amplification was conducted with initial denaturation for 5 min at 95 °C, followed by 40 cycles of 45 s at 94 °C, 45 s at 50 to 54 °C and 2 min at 72 °C, with a final extension of 7 min at 72 °C and holding at 4 °C. The

PCR products were separated in a 1.5% agarose gel using 1 $\times$ TBE running buffer, stained, and photographed, and the records were preserved.

Data analysis

SCoT amplicons were scored in a binary matrix as present (1) or absent (0) for each sample based on the corresponding standard size. Vague bands that could not be easily detected were not scored. POPGEN32 software was used to calculate the total number of bands (NPB) and the number of polymorphous bands (PPB) obtained with the SCoT primers to analyze the genetic diversity index of *P. sibirica* and to calculate Nei's gene diversity index (H), Shannon's information index (I), the number of alleles (Na), the effective number of alleles (Ne), total genetic diversity (Ht), population genetic diversity (Hs), the coefficient of

Table 2 Screened SCoT primer sequence information

Code	Primer name	Primer Sequences (5'- 3')	GC%	Annealing Tm (°C)
1	SCoT3	CAACAATGGCTACCACCG	55.60	52
2	SCoT4	CAACAATGGCTACCACCT	50.00	50
3	SCoT10	CAACAATGGCTACCAGCC	55.60	55
4	SCoT12	ACGACATGGCGACCAACG	61.10	54
5	SCoT13	ACGACATGGCGACCATCG	61.10	54
6	SCoT21	ACGACATGGCGACCCACA	61.10	54
7	SCoT23	CACCATGGCTACCACCAG	61.10	54
8	SCoT25	ACCATGGCTACCACCGGG	66.70	53
9	SCoT27	ACCATGGCTACCACCGTG	61.10	54
10	SCoT29	CCATGGCTACCACCGGCC	72.20	51
11	SCoT30	CCATGGCTACCACCGGCG	72.20	51
12	SCoT31	CCATGGCTACCACCGCCT	66.70	53
13	SCoT32	CCATGGCTACCACCGCAC	66.70	53
14	SCoT34	ACCATGGCTACCACCGCA	61.10	54
15	SCoT43	CAATGGCTACCACCGCAG	61.10	54
16	SCoT44	CAATGGCTACCATTAGCC	50.00	50
17	SCoT45	ACAATGGCTACCACTGAC	50.00	53
18	SCoT46	ACAATGGCTACCACTGAG	50.00	50
19	SCoT47	ACAATGGCTACCACTGCC	55.60	52
20	SCoT53	ACAATGGCTACCACCGAC	55.60	52
21	SCoT58	ACAATGGCTACCACTAGG	50.00	50
22	SCoT71	CCATGGCTACCACCGCCG	72.20	51
23	SCoT72	CCATGGCTACCACCGCCC	72.20	51

genetic differentiation (Gst), and gene flow (Nm). Ntsys-2.0 software was used to calculate Nei's genetic distances from the binary (0, 1) matrices obtained, and the unweighted pair-group method with arithmetic means (UPGMA) method performed in the SHAN program was used for the cluster analysis of the 10 populations to construct the cluster relationship tree diagram. MEGA7 software was used for the cluster analysis of the 278 individuals based on Nei's genetic similarity matrix obtained via the arithmetic mean (UPGMA) method. An analysis of molecular variance (AMOVA) and GenAlEx software were used for the analysis of genetic differentiation among and within populations. A principal coordinate analysis (PCoA) was performed according to the binary matrix using GenAlEx software. STRUCTURE software was used to determine the genetic structure of the studied population, and K was tested from 1 to 10 with ten replicates. IBD software was used for Mantel test to analyze the correlation between genetic distance and geographic distance and altitude.

Results

SCoT polymorphism analysis

In the SCoT analysis, 23 SCoT primers were used for marker amplification in 278 *P. sibirica* individuals. A total of 292 bands that could be scored were produced, among which 289 bands were polymorphic, with a mean of 12.6 polymorphic bands per primer. The polymorphism percentage was 98.87%. Among the 23 SCoT primers, SCoT21, SCoT32, and SCoT53 amplified a maximum of 17 bands, and the polymorphism percentage was 100%. The number of bands amplified by the SCoT 25 primer was at least 9, and the polymorphism percentage was 90%. The 23 SCoT primers produced Na values ranging from 1.90 to 2.00, with a mean of 1.99; Ne values ranging from 1.28 to 1.60, with a mean of 1.44; H values ranging from 0.18 to 0.34, with a mean of 0.26; and I values ranging from 0.30 to 0.50, with a mean of 0.41 (Table 3).

Table 3 Polymorphism information obtained from SCoT primer amplification in *P. sibirica*

	Primer	Total loci	NPB	PPB (%)	Na	Ne	H	I
The total number of bands (NPB), the number of polymorphous bands (PPB), the number of alleles (Na), the effective number of alleles (Ne), Nei's gene diversity index (H), Shannon's information index (I)	SCoT3	9	9	100.00	2.00	1.50	0.30	0.47
	SCoT4	11	11	100.00	2.00	1.45	0.29	0.45
	SCoT10	11	11	100.00	2.00	1.51	0.32	0.50
	SCoT12	11	11	100.00	2.00	1.35	0.22	0.35
	SCoT13	16	16	100.00	2.00	1.28	0.18	0.31
	SCoT21	17	17	100.00	2.00	1.40	0.24	0.38
	SCoT23	11	11	100.00	2.00	1.56	0.33	0.49
	SCoT25	10	9	90.00	1.90	1.49	0.28	0.43
	SCoT27	12	12	100.00	2.00	1.51	0.30	0.46
	SCoT29	11	11	100.00	2.00	1.34	0.21	0.33
	SCoT30	12	11	91.67	1.92	1.36	0.22	0.35
	SCoT31	10	10	100.00	2.00	1.60	0.34	0.50
	SCoT32	17	17	100.00	2.00	1.45	0.27	0.43
	SCoT34	14	14	100.00	2.00	1.46	0.28	0.44
	SCoT43	11	10	90.91	1.91	1.29	0.18	0.30
	SCoT44	14	14	100.00	2.00	1.49	0.28	0.42
	SCoT45	15	15	100.00	2.00	1.30	0.20	0.33
	SCoT46	12	12	100.00	2.00	1.42	0.24	0.36
	SCoT47	13	13	100.00	2.00	1.39	0.24	0.38
	SCoT53	17	17	100.00	2.00	1.38	0.23	0.37
	SCoT58	10	10	100.00	2.00	1.56	0.33	0.50
	SCoT71	12	12	100.00	2.00	1.46	0.27	0.42
	SCoT72	16	16	100.00	2.00	1.51	0.31	0.47
	Mean	12.7	12.6	98.87	1.99	1.44	0.26	0.41
	Total	292	289					

Population genetic diversity and genetic differentiation

The results of the genetic diversity and genetic differentiation analyses in the 10 populations of *P. sibirica* in Inner Mongolia showed that NPB ranged from 110 to 180, with a mean of 156.40 bands; PPB ranged from 37.67 to 61.64, with a mean of 53.56 polymorphisms; Na ranged from 1.40 to 1.62, with a mean of 1.54; Ne ranged from 1.20 to 1.31, with a mean of 1.28; H ranged from 0.12 to 0.19, with a mean of 0.17; and I ranged from 0.18 to 0.29, with a mean of 0.26. Among the 278 individuals, the overall PPB, Na, Ne, H, and I total values were 97.94%, 1.98, 1.43, 0.26, and 0.40, respectively (Table 4), implying the presence of considerable genetic diversity among the *P. sibirica* accessions. The genetic diversity was determined to be highest among the KYZ population, with an H value of 0.19 and a PPB of 61.64%. The

second-highest H value of 0.18 was observed for KZH, for which PPB was 60.62%. In contrast, the genetic diversity was found to be lowest in WJG, with an H value of 0.12 and a PPB of 37.67%.

The 278 individuals showed an H_t of 0.26, with a variance of 0.03; H_s of 0.17, with variance of 0.01; G_{st} of 0.36, where the mean showed a high G_{st} value ($G_{st} > 0.25$); and N_m of 0.89 ($N_m < 1$), with the mean showing low gene flow (Table 5). The AMOVA showed 9 degrees of freedom among the populations and 267 degrees of freedom within the populations. It showed that the mean squares of the variance among the populations was 4,622.89, and that within the populations was 7,385.47. The among-population variance was 17.61 on average and the within-population variance was 27.56. Data from the AMOVA molecular detection prove that 39% of total genetic variance occurred among the populations, while 61% occurred for within the population genetic

Table 4 Genetic diversity indices of *P. sibirica* populations

	Population	No. of Samples	NPB	PPB (%)	Na	Ne	H	I
The total number of bands (NPB), the number of polymorphous bands (PPB), the number of alleles (Na), the effective number of alleles (Ne), Nei's gene diversity index (H), Shannon's information index (I)	KSK	46	145	49.65	1.51	1.27	0.16	0.24
	WJG	23	110	37.67	1.40	1.20	0.12	0.18
	KYZ	26	180	61.64	1.62	1.31	0.19	0.29
	KZH	38	177	60.62	1.61	1.31	0.18	0.28
	ZLA	22	162	55.48	1.56	1.29	0.17	0.27
	LC	30	156	53.42	1.54	1.27	0.16	0.25
	BLY	26	169	57.87	1.59	1.31	0.18	0.28
	WLS	14	153	52.39	1.53	1.27	0.16	0.25
	AL	27	156	53.42	1.54	1.27	0.16	0.25
	ZLT	26	156	53.42	1.54	1.28	0.17	0.26
	Mean	27.80	156.40	53.56	1.54	1.28	0.17	0.26
	Total	278	286	97.94	1.98	1.43	0.26	0.40

Table 5 Genetic differentiation within *P. sibirica* populations

	Total	Ht	Hs	Gst	Nm
Mean	278	0.26	0.17	0.36	0.89
StDev		0.03	0.01		

The total genetic diversity (Ht), population genetic diversity (Hs), the coefficient of genetic differentiation (Gst), gene flow (Nm)

variability (Table 6). This result indicates that there is low genetic differentiation among populations.

Analysis of the among-population genetic structure

POPGEN32 was used to analyze the genetic distance and genetic similarity of the 10 *P. sibirica* populations in Inner Mongolia. The analyses of Nei's genetic similarity and Nei's genetic distance showed that the population genetic distance ranged from 0.03 to 0.25

while the genetic similarity ranged from 0.77 to 0.97. The genetic similarity between WJG and KSK was the highest, at 0.97, and their genetic distance was the lowest, at 0.03. The lowest genetic similarity between ZLT and WJG was 0.77, and the greatest genetic distance was 0.25 (Table 7). Based on Nei's genetic distance, the UPGMA method was used for cluster analysis. The results showed that the 10 populations were divided into two major groups at a genetic distance threshold of 0.22 (Fig. 2). Group I included KSK, WJG, LC, KYZ, KZH, and ZLA. Group II included BLY, AL, WLS, and ZLT. When the genetic distance was less than 0.17, group I was divided into subgroup I and subgroup II. Subgroup I included KSK, while subgroup II included WJG, LC, KYZ, KZH, and ZLA. Group II was divided into subgroup III and subgroup IV. Subgroup III included BLY, AL, and WLS, while subgroup IV included ZLT.

Analysis of genetic structure within populations

Based on Nei's genetic similarity, MEGA7 software was used to perform the clustering analysis of 278

Table 6 Analysis of molecular variance (AMOVA) of *P. sibirica* populations

Source of Variance	Degree of freedom	Mean squares	Variance component	Percentage of variance (%)	P value
Among populations	9	4,622.89	17.61	39	< 0.001
Within populations	267	7,385.47	27.56	61	< 0.001
Total	278	12,008.37	45.22		

Table 7 Genetic similarity (above diagonal) and genetic distance (below diagonal) among populations

Population	KSK	WJG	KYZ	KZH	ZLA	LC	BLY	WLS	AL	ZLT
KSK		0.97	0.91	0.89	0.88	0.85	0.87	0.83	0.81	0.80
WJG	0.03		0.90	0.87	0.86	0.83	0.85	0.81	0.78	0.77
KYZ	0.10	0.10		0.95	0.92	0.88	0.88	0.85	0.82	0.81
KZH	0.12	0.14	0.05		0.94	0.91	0.90	0.86	0.84	0.82
ZLA	0.13	0.15	0.08	0.06		0.96	0.93	0.90	0.88	0.87
LC	0.16	0.18	0.13	0.10	0.04		0.95	0.90	0.89	0.89
BLY	0.14	0.16	0.12	0.11	0.08	0.05		0.92	0.89	0.89
WLS	0.18	0.21	0.16	0.15	0.11	0.11	0.08		0.94	0.91
AL	0.21	0.24	0.20	0.17	0.13	0.12	0.12	0.06		0.96
ZLT	0.22	0.25	0.21	0.20	0.14	0.11	0.11	0.09	0.04

Fig. 2 UPGMA dendrogram of *P. sibirica* populations based on Nei's genetic distance

individuals (Fig. 3), with the PCoA performed using GenAlEx software (Fig. 4). The results showed that the 278 individuals were divided into two groups. The UPGMA clustering analysis showed that the 278 individuals were divided into group I and group II when the genetic similarity was 0.17. Group I included 144 individuals, while group II included 134 individuals. Group I included 23 WJG individuals, 46 KSK individuals, 26 KYZ individuals, 38 KZH individuals, and 11 ZLT individuals. Group II included 27 AL individuals, 14 WLS individuals, 26 ZLT individuals, 25 BLY individuals, 31 LC individuals, and 11 ZLA individuals. When the genetic similarity was ≤ 0.14 ,

group I and group II were divided into four subgroups. A total of 23 WJG individuals and 46 KSK individuals were clustered into subgroup I; 26 KYZ individuals, 38 KZH individuals and 11 ZLA individuals were divided into subgroup II; 27 AL individuals, 12 ZLT individuals, and 9 WLS individuals were clustered into subgroup III; and 14 ZLT individuals, 5 WLS individuals, 11 ZLA individuals, 26 BLY individuals, and 30 LC individuals were included in subgroup IV. The PCoA results showed that the 278 *P. sibirica* individuals were divided into groups I and II. These results were relative to those of the UPGMA clustering analysis of similarity. The two groups were divided

Fig. 3 UPGMA dendrogram of *P. sibirica* populations based on the SCoT markers

into four subgroups. Group I included all individuals in the KSK, WJG, KZH, and KYZ populations and some individuals from ZLA. Group II included all individuals from AL, WLS, ZLT, BLY, and LC and some individuals from ZLA.

To further explore the genetic relationships within populations of *P. sibirica*, the population structure of 278 *P. sibirica* individuals was evaluated using

STRUCTURE version 2.3.4 software (Beaumont et al. 2001). ΔK values computed for all classes indicated a strong signal for $K = 2$ (Fig. 5), and $K = 2$ values provided the most rational arrangement of *P. sibirica* in different regions. A total of 278 *P. sibirica* individuals were divided by the STRUCTURE analysis into 2 groups (Fig. 6). Group I included all individuals in the KSK, WJG, KZH, and KYZ

Fig. 4 Principal coordinate analysis (PCoA) of *P. sibirica* populations based on SCoT markers

Fig. 5 Evanno's test based on delta k among *P. sibirica* cultivars studied

Fig. 6 Population structure of the 278 individuals at K = 2. The same color indicates the same group

populations and some individuals from ZLA. Group II included all individuals from AL, WLS, ZLT, BLY, and LC and some individuals from ZLA. This result is similar to the clusters identified using UPGMA and PCoA.

Discussion

In this study, a genetic diversity analysis was performed using SCoT markers to assess phylogenetic relationships among 278 *P. sibirica* individuals from 10 different populations in Inner Mongolia. Twenty-

three SCoT primers were used for the amplification of genomic DNA from *P. sibirica*. A total of 292 clear bands were obtained, 289 of which were polymorphic. The average number of amplified bands per primer was 12.6, with a polymorphism percentage of 98.87%. In a previous study, researchers used SRAP (Ai et al. 2011), AFLP (Wang 2008), RAPD (Lu 2008), and ISSR (Liu et al. 2011, 2007; Duan et al. 2010; Li et al. 2009) molecular markers to analyze the genetic diversity of *P. sibirica* in different regions, and the results showed that the polymorphism ranged from 58 to 90%. In contrast, the SCoT molecular markers were found to be more polymorphic than other molecular markers. Therefore, SCoT markers may be suitable for the study of genetic diversity according to molecular markers in *P. sibirica*. A previous study of the genetic diversity of *Prunus armeniaca* (Li 2014) indicated H , I , and polymorphism percentage values of $H = 0.17$, $I = 0.26$ and $PPB = 50.07\%$; *P. sibirica* (Ma 2014) indicated H , I , and polymorphism percentage values of $H = 0.23$, $I = 0.35$ and $PPB = 76.74\%$. However, the H , I , and polymorphism percentage values obtained in the present study were $H = 0.26$, $I = 0.40$ and $PPB = 97.94\%$, respectively. We showed that the genetic diversity of the *P. sibirica* populations in Inner Mongolia is higher than that of *Prunus armeniaca* and *P. sibirica*. This result was consistent with the conclusions of other scholars that the genetic diversity of outbred species is higher (Hamrick 1989; Zheng et al. 2008).

As a perennial wild resource, *P. sibirica* is widely distributed in Inner Mongolia and exhibits a large distribution area, long-term evolution and considerable diffusion as well as ecological diversity, leading to rich genetic diversity. According to Wright (1972), when the F_{ST} (G_{ST} , genetic differentiation coefficient) value is 0 to 0.05, the genetic differentiation within populations is low; when the F_{ST} value is 0.05 to 0.15, the genetic differentiation within populations is moderate; when the F_{ST} value is 0.15 to 0.25, the genetic differentiation within populations is high; and when the F_{ST} value is greater than 0.25, the genetic differentiation within populations is great. In our research, the G_{ST} was 0.36. The results showed that the genetic differentiation coefficient of *P. sibirica* within populations in Inner Mongolia was relatively high, with among-population genetic differentiation of 39% and a within-population genetic differentiation of 61%. Wang (2019), Liu et al. (2012) and Ma (2013),

who used SSR and ISSR markers for *P. sibirica* and wild apricot genetic diversity analyses, showed that within-population variation dominated, and the among-population variation was much lower than the within-population variation, which is similar to the results of a previous study in *P. sibirica*.

The reasons for the high within-population genetic differentiation of *P. sibirica* in Inner Mongolia may include the following: 1. Gene flow is an important factor affecting genetic differentiation in a population when $N_m < 1$, which can conversely effectively prevent genetic differentiation caused by genetic drift. The N_m value for *P. sibirica* in Inner Mongolia was 0.89, indicating that genetic drift was not the main factor influencing the genetic differentiation of the *P. sibirica* populations in Inner Mongolia. Some gene flow exists among the *P. sibirica* population in Inner Mongolia, but the N_m intensity was relatively low. 2. The main breeding method of *P. sibirica* is outcrossing, self-incompatibility will refuse self-pollination or inbreeding, and its breeding mainly relies on natural pollination or pollination by visiting insects such as bees, flies and a few butterflies (Liu et al. 2010; Liu 2010). Although pollen can be spread by insects and by wind over long distances, the large distribution area and discontinuities in the population distribution of *P. sibirica* are restricted due to habitat fragmentation (Wu et al. 2015), thus weakening the gene flow within populations of *P. sibirica*. 3. Some populations have been destroyed or their habitats have been degraded, resulting in a gradual fragmentation of the population distribution and limiting the gene flow within the populations. Therefore, *P. sibirica* in Inner Mongolia shows high within-population genetic differentiation, and the topography and landforms and human activities could be potential factors influencing the genetic difference.

In the present study, the genetic structure of *P. sibirica* populations in Inner Mongolia was analyzed. From the clustering diagram, principal component analysis, and structure analysis, showed that most of the populations are clustered together with a similar geographic distribution, and the Mantel test showed a not significant correlation between the genetic distance and geographic distance and between the genetic distance and altitude of the *P. sibirica* studied ($r = 0.04$, $P \leq 0.48$; $r = 0.03$, $P \leq 0.43$). Therefore, the genetic difference among *P. sibirica* is not due to their geographical distance and altitude. The clustering

diagram, principal component analysis and structure analysis of the 278 individuals divided the populations into two groups (I and II). KSK, WJG, KYZ, and KZH were clustered in group I; LC, BLY, WLS, AL, and ZLT were clustered in group II; and only the 22 individuals in the ZLA group were divided between the two groups. A certain degree of gene flow exists among the subgroups, and gene flow also affects the population structure of *P. sibirica* (Liu et al 2012). This may occur because of the area of many sandy land in Inner Mongolia, and the topography and landforms are very complex. Thus, weakening the gene flow opportunities among populations. Second, habitat fragmentation is also an important factor that affects the composition and structure of the ecosystem and the genetic structure of a species.

Conclusion

This study revealed that *P. sibirica* shows high genetic diversity in Inner Mongolia. Therefore, improving the preservation and utilization of specific germplasms of different populations is necessary to avoid the disappearance of a large number of valuable genetic resources. Moreover, expanding the collection of resources to implement transfer protection will be conducive to better scientific research and resource protection in *P. sibirica*, increase the gene flow among populations, and improve the genetic diversity of *P. sibirica*. Based on resource collection, genetic diversity analyses, and genetic structure analyses, the breeding of *P. sibirica* varieties with high yields and frost resistance is being actively performed to meet the increasing demand for the industrial production of *P. sibirica* resources.

Acknowledgements This research was supported by the National Science and Technology of China (2013BAD14B0203) and the Inner Mongolia Science and Technology Department of China (2015020553).

Author contribution HBR wrote the manuscript, organized the data and designed the research. SRL provided the funding for the experiment. WZY and SF conducted experiments. BYE conceived and designed the research. All authors read and approved the manuscript.

Data availability Data is contained within the article.

Declarations

Conflict of interest The authors declare that there is no conflict of interests regarding the publication of this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ai PF, Zhen ZJ, Jin ZZ (2011) Genetic diversity and relationships within sweet kernel apricot and related *Armeniaca*, species based on sequence-related amplified polymorphism markers. *Biochem Syst Ecol* 39(4–6):694–699
- Bao WQ, WuYun TN, Wang L, Zhao H, Du HG (2016) Genetic diversity and population genetic structure of siberian apricot in Inner Mongolia. *Acta Botanica Boreali-Occidentalia Sinica* 36(11):2182–2191
- Beaumont M, Gottelli D, Barratt EM, Kitchener AC, Daniels MJ, Pritchard JK, Bruford MW (2001) Genetic diversity and introgression in the Scottish wildcat. *Mol Ecol* 10(2):319–336
- Collard BCY, Mackill DJ (2009) Start codon targeted (SCoT) polymorphism: a simple novel DNA marker technique for generating gene-targeted markers in plants. *Plant Mol Biol Report* 27(1):86–93
- Dong YS, He R, Lin FQ (1991) Study on genetic resources of *Prunus sibirica*. *Northern Fruit Trees* 1:4–8
- Dong SJ, Wang L, Liu LX, Liu MG, Wu YL, Chen JH, Xia ZZ (2018) Study on quantitative characteristics in *Armeniaca sibirica* clones from different provenances. *Nonwood Forest Res* 36(04):1–8
- Duan W, Yan WD, Wu YTN (2010) Identification and analysis of 14 samples in *Prunus Simonii* by ISSR molecular marker. *J Inner Mongolia Agric Univ* 2:147–153
- Guan CF, Chachar S, Zhang PX, Hua CQ, Wang RZ, Yang Y (2020) Inter- and intra-specific genetic diversity in diospyros using SCoT and IRAP markers. *Horticult Plant J* 6(2):71–80
- Hamrick JL (1989) Isozymes and the analysis of genetic structure in plant populations. *Isozymes Plant Biol.* pp 87–105
- Jin L, Liu GQ, Huang SJ, Wu YL, Zhang X (2018) Genetic diversity and fingerprints of 97 *Armeniaca sibirica* clones based on SSR markers. *Scientia Silvae Science* 54(7):51–61
- Li MM, Cai YL, Qian ZQ (2009) Genetic diversity and differentiation in Chinese sour cherry *Prunus pseudocerasus*

- lindl., and its implications for conservation. *Genet Resour Crop Evol* 56(4):455–464
- Li M, Zhao Z, Yang JA (2011a) Classification on genetic resources of *Prunus sibirica* in the Loess Plateau. *J Northwest Forestry Univ* 1:8–12
- Li M, Zhao Z, Yang JA (2011) Genetic diversity analysis on germplasm of *Prunus sibirica* in different counties in Loess Plateau. *J Northwest A&F Univ* 2:143–149+156.
- Li M (2014) Genetic diversity of common apricot and siberian apricot in north China. Northwest A&F University, Yangling
- Liu LM (2010) Primary research on reproductive ecology of *Armeniaca vulgaris* Lam. Xinjiang Agricultural University, Wulumuqi
- Liu W, Liu D, Zhang A (2007) Genetic diversity and phylogenetic relationships among plum genetic resources in China assessed with inter-simple sequence repeat markers. *J Am Soc Hortic Sci* 132(5):619–628
- Liu YX, Chen WZ, Liu WS (2010) Palynological study on the origin and systematic evolution of kernel-using apricots. *Acta Horticulturae Sinica* 9:1377–1387
- Liu MP, Bo DL, Tian M (2011) Analysis on genetic differentiation of *Armeniaca cathayana* F₁ hybrids by ISSR. *J Central South Univ Forestry Technol* 10:100–104
- Liu HB, Wang Z, Liu J (2012) Genetic diversity and genetic structure of *Prunus sibirica* populations in Yan Mountains. *Scientia Silvae Sci* 8:68–74
- Lu NC (2008) Genetic resources on the genetic diversity of siberian apricot in Jilin. Jilin Agricultural University, Changchun
- Lu CY (2018) Genetic diversity based on SSR and its association analysis with phenotypic traits in *Armeniaca vulgaris*. Shenyang Agricultural University, Shenyang
- Ma LJ (2013) Genetic diversity and genetic structure analysis of wild apricot populations in Northeast China. Jilin Agricultural University, Changchun
- Ma YH, Zhao Z, Li KY (2007) Optimization of technology for almond oil extraction by supercritical CO₂. *Trans CSAE* 4:272–275
- Ma LJ, Guo TJ, He DD, Li F, Zhang YB, Ma LX (2014) Genetic diversity analysis of two natural *A. sibirica* populations in northeast China. *J Jilin Agric Univ* 36(03):300–305
- Rao VR, Hodgkin T (2002) Genetic diversity and conservation and utilization of plant genetic resources. *Plant Cell, Tissue Organ Cult* 68(1):1–19
- Saboori S, Noormohammadi Z, Sheidai M, Marashi SS (2020) SCoT molecular markers and genetic fingerprinting of date palm (*Phoenix dactylifera* L.) cultivars. *Genet Resour Crop Evol* 67:73–82
- Wang L (2019) Genetic diversity and population genetic structure of *Armeniaca sibirica* clones. Shenyang Agricultural University, Shenyang
- Wang SX, Hu XY, Fan SQ, Cu YF, Yu HY, Wang LB (2019) Geographical variation analysis on *Armeniaca sibirica* seed characteristics from different provenances. *Nonwood Forest Res* 37(01):74–79+86
- Wang L (2008) Using AFLP markers to analyze genetic diversity in *Prunus Sibirica* (L.) Lam. Hebei Agricultural University, Baoding
- Wright S (1972) Evolution and the genetics of populations. A treatise in four volumes. Volume 4. Variability within and among natural populations. *J Biosoc Sci* 4(2):253–256.
- Wu YT, Amo H, Xu J (2015) Population structure of and conservation strategies for wild pyrus ussuriensis maxim in China. *Plos One* 10(8):e0133686
- Yao ZY, Li KY, Zhao Z (2007) Antioxidant properties of melanin from testae of *Armeniaca vulgaris* var. Ansu. *Scientia Silvae Science* 10:59–63
- Yu DJ (1979) Chinese fruit tree taxonomy. Beijing Agricultural Press. pp 45–90
- Zeng J, Zeng YQ, Zang CH (2008) Genetic diversity in natural populations of *Sorbus Puhuashanensis*. *Biodivers Sci* 6:562–569
- Zhou L, He XH, Yu HX, Chen MY, Fan Y, Zhang XJ, Fang ZB, Luo C (2020) Evaluation of the genetic diversity of mango (*Mangifera indica* L.) seedling germplasm resources and their potential parents with start codon targeted (SCoT) markers. *Genet Resour Crop Evol* 67:41–58

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.