

Erratum to: Influence of hydrophyte abundance on the spatial distribution of zooplankton in selected lakes in Greece

K. Stefanidis · E. Papastergiadou

Published online: 12 January 2017
© Springer International Publishing Switzerland 2017

Erratum to: Hydrobiologia (2010) 656:55–65
DOI 10.1007/s10750-010-0435-0

Due to an unfortunate mistake, the units for ‘Total phosphorus’ and ‘Dissolved inorganic nitrogen’ pre-

sented in Table 2 are incorrectly displayed in the original publication. The correct units must read $\mu\text{g/l}$ instead of mg/l .

The corrected Table 2 is published here and should be treated as definitive.

Table 2 Mean values for physicochemical parameters amongst the four lakes studied

Standard error in brackets

	Mikri Prespa	Kastoria	Vegoritis	Petron
Total phosphorus ($\mu\text{g/l}$)	125.5 (22.7)	133 (18.7)	178.3 (60.7)	142.5 (19.9)
Dissolved inorganic nitrogen ($\mu\text{g/l}$)	644.4 (165.8)	390.5 (76.5)	425.7 (98.5)	415.9 (69.9)
Chlorophyll- <i>a</i> (mg/m^3)	17.4 (2.9)	21.3 (2.2)	6 (0.83)	30.8 (6.73)
pH	7.8 (0.12)	8.1 (0.12)	7.95 (0.16)	7.3 (0.17)
DO (mg/l)	9.8 (0.6)	8.89 (0.46)	5.47 (0.57)	4.67 (0.32)
Conductivity ($\mu\text{S/cm}$)	292.1 (3.93)	334.3 (3)	718.9 (39.6)	852.5 (15.7)
Secchi depth (m)	1.21 (0.24)	0.53 (0.03)	0.88 (0.28)	0.36 (0.04)

The online version of the original article can be found under doi:[10.1007/s10750-010-0435-0](https://doi.org/10.1007/s10750-010-0435-0).

K. Stefanidis (✉) · E. Papastergiadou
Department of Biology, University of Patras,
26500 Patras, Greece
e-mail: kstefani@upatras.gr

E. Papastergiadou
e-mail: evapap@upatras.gr