



Quantum Violation of the Suppes-Zanotti Inequalities and “Contextuality”

Karl Svozil¹

Received: 8 February 2021 / Accepted: 25 May 2021 / Published online: 10 June 2021
© The Author(s) 2021

Abstract

The Suppes-Zanotti inequalities involving the joint expectations of just three binary quantum observables are (re-)derived by the hull computation of the respective correlation polytope. A min-max calculation reveals its maximal quantum violations correspond to a generalized Tsirelson bound. Notions of “contextuality” motivated by such violations are critically reviewed.

Keywords Suppes-Zanotti inequalities · Greenberger-Horne-Zeilinger argument · Kochen-Specker theorem · Born rule · Min-max calculation · Convex polytope · Contextuality

1 Two-Partite Vector-Based Expectations not Satisfying Classical Bounds

Classical bounds on probabilities and expectations can be expected to be “violated” by or “being different” from quantum probabilities and expectations because the latter are based on multi-dimensional vectorial entities whereas the former are based on scalars in (sub)sets of power sets. Exactly how these violations are operationalized and measured has developed from an intuitive, heuristic search in the early days [1–3] into a systematic method [4–8].

The most elementary expression of the classical versus quantum difference quoted earlier is the two-partite correlation function of two dichotomic observables $X, Y \in \{-1, +1\}$. It is empirically collected from a series of N measurements of X and Y and defined by $\langle X, Y \rangle_s \approx \frac{1}{N} \sum_{i=1}^N X_i Y_i$, where the index i refers to the i th measurement, and s refers to a specific (unaltered) state on which these repetitive measurements are performed. It is assumed that, if N increases, the limit exists and is monotonically approached – that is, for “large enough” N , $\langle X, Y \rangle_s$ is a “good approximation” [9].

✉ Karl Svozil
svozil@tuwien.ac.at

¹ Institute for Theoretical Physics, TU Wien, Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria

1.1 Classical Predictions on “singlet-type” States

It is not too difficult to model a classical two-partite state q which shows “singlet-like” characteristics – an example would be the angular momentum in a particular spacial direction of two fragments of a bomb which originally had no angular momentum in any direction [10]. An argument involving equidistribution of angular momenta of the fragments reveals a linear classical correlation on such a state; that is, $\langle X, Y \rangle_c = \mathbf{E}(X, Y) = \mathbf{E}(\theta) = -1 + 2\theta/\pi$, where the angle $0 \leq \theta \leq \pi$ characterizes the “spatial separation” of the directions of these observables X and Y .

1.2 General Classical Predictions

To construct a generic classical situation, a generalized urn model [11] is introduced which can also be phrased in terms of finite-state identification problems of automata allowing complementarity [12]. Its formalization is in terms of set-theoretic partitions [13] and power sets.

In terms of generalized urn models, we consider urns filled with black balls painted with three different colors, one color per observable X, Y , and Z . Since each observable may have two different outcomes we can, for instance, label these outcomes by “+” and “-”, printed on these balls in the respective colors. There are eight such ball types. As the urn is filled with an arbitrary distribution of ball types, it can only be ascertained that they occur with probabilities $0 \leq \lambda_{\pm\pm\pm} \leq 1$, were the indices refer to the respective symbols in the colors associated with our three observables. Since in such a scheme the ball types are mutually exclusive and their enumeration is complete (i.e., exhaustive), we can suppose that $\sum_{i,j,k \in \{+, -\}} \lambda_{ijk} = 1$, and the joint expectations add up accordingly; e.g., $\mathbf{E}(X, Y) = \sum_{k \in \{+, -\}} [\lambda_{++k} + \lambda_{--k} - (\lambda_{+-k} + \lambda_{-+k})]$.

1.3 Quantum Predictions on a Singlet State

The quantum predictions of a single observable in an arbitrary direction characterized by the spherical coordinates $0 \leq \theta \leq \pi$ and $0 \leq \varphi < 2\pi$ is derived from the Pauli spin matrices σ_x, σ_y and σ_z forming the spin operator $\boldsymbol{\sigma}(\theta, \varphi) = \sigma_x \sin \theta \cos \varphi + \sigma_y \sin \theta \sin \varphi + \sigma_z \cos \theta$ and the single particle projection operator $\mathbf{S}_{\pm}(\theta, \varphi) = \frac{1}{2} [\mathbb{I}_2 \pm \boldsymbol{\sigma}(\theta, \varphi)]$ for the states “-” and “+”, respectively. The respective two-partite projection operators are $\mathbf{S}_{\pm_1 \pm_2}(\theta_1, \varphi_1, \theta_2, \varphi_2) = \mathbf{S}_{\pm_1}(\theta_1, \varphi_1) \otimes \mathbf{S}_{\pm_2}(\theta_2, \varphi_2)$. Finally, the operator associated with the two-partite expectations is $\mathbf{F}(X, Y) = \mathbf{F}(\theta_1, \varphi_1, \theta_2, \varphi_2) = \mathbf{S}_{++} + \mathbf{S}_{--} - (\mathbf{S}_{+-} + \mathbf{S}_{-+}) = \boldsymbol{\sigma}(\theta_1, \varphi_1) \otimes \boldsymbol{\sigma}(\theta_2, \varphi_2)$.

Suppose we are interested in the correlation function for a singlet state in the Bell basis $q = |\Psi_{-}\rangle\langle\Psi_{-}|$ with $|\Psi_{-}\rangle = \frac{1}{\sqrt{2}}(0, 1, -1, 0)^T$, then the quantum prediction yields $\langle X, Y \rangle_q = \mathbf{F}(\theta_1, \varphi_1, \theta_2, \varphi_2) = -[\cos \theta_1 \cos \theta_2 + \cos(\varphi_1 - \varphi_2) \sin \theta_1 \sin \theta_2]$. For $\varphi_1 = \varphi_2$ this reduces to the well-known cosine form $\langle X, Y \rangle_q = \mathbf{F}(\theta_1, 0, \theta_2, 0) = \mathbf{F}(\theta_1 - \theta_2) = -\cos(\theta_1 - \theta_2)$, that is, the two-partite correlation for dichotomic observables $X, Y = \pm 1$ of the two-partite singlet state is proportional to the Euclidean scalar product between the vectors associated with X and Y .

The maximal quantum-to-classical violations

$$\max_{\theta \in [0, \pi]} |\mathbf{E}(\theta) - \mathbf{F}(\theta)| = \sqrt{1 - \left(\frac{2}{\pi}\right)^2} - \frac{2}{\pi} \cos^{-1} \frac{2}{\pi} \approx 0.2 \quad (1)$$

resulting from less, as well as more, equal occurrences of the joint observables $++/--$ and $+--/+$, occur at angles $(d/d\theta) [\mathbf{E}(\theta) - \mathbf{F}(\theta)] = 0$, that is, at

$$\begin{aligned} \theta &= \sin^{-1} \frac{2}{\pi} \text{ as well as} \\ \theta &= \pi - \sin^{-1} \frac{2}{\pi}, \text{ respectively.} \end{aligned} \tag{2}$$

1.4 Quantum Predictions on more General Pure States

By a min-max calculation [14] it is not too difficult to compute those quantum states which, given arbitrary angles between the two observables X and Y , yield the minimal and maximal correlations: all that is needed is the eigensystem of $\mathbf{F}(\theta_1, \varphi_1, \theta_2, \varphi_2)$. Rather than enumerating this eigensystem in full generality the special case $\theta_1 = \theta$ and $\theta_2 = \varphi_1 = \varphi_2 = 0$ is posted, resulting in the (decomposable) vectors (modulo normalization)

$$\begin{aligned} |\psi_{1,\min}\rangle &= (0, \cos \theta + 1, 0, \sin \theta)^\top \text{ as well as} \\ |\psi_{2,\min}\rangle &= (\cos \theta - 1, 0, \sin \theta, 0)^\top \end{aligned} \tag{3}$$

for the minimal expectation $\langle X, Y \rangle = -1$; and

$$\begin{aligned} |\psi_{1,\max}\rangle &= (0, \cos \theta - 1, 0, \sin \theta)^\top \text{ as well as} \\ |\psi_{2,\max}\rangle &= (\cos \theta + 1, 0, \sin \theta, 0)^\top \end{aligned} \tag{4}$$

for the maximal expectation $\langle X, Y \rangle = 1$, respectively.

2 The Case of Three Observables

2.1 Classical Bounds

One might as well stop here, contemplate the elementary difference between two forms of probabilities based on scalars and power sets in the classical case, and on vectors and the vector space spanned by them in the quantum case, and leave it at that. However, this is not what happened historically: Bell and others tried to find criteria for non-compliance with classical behavior involving more than just two observables. In particular, Suppes and Zanotti [15–17] presented special cases of what Boole called “conditions of possible experience” [18, 19] involving just three dichotomic observables $X, Y, Z \in \{-1, +1\}$.

The original method of deriving these bounds is rather involved. But with today’s convex polytope techniques [4, 7, 20] it is not too difficult to derive those inequalities: (i) form all possible combinations of joint occurrences by multiplying the respective dichotomic observables – in this case $\mathbf{E}(X, Y) = XY, \mathbf{E}(X, Z) = XZ, \mathbf{E}(Y, Z) = YZ$; (ii) form the 3-tuples (that is, the finite ordered list or sequence) of all three numbers for particular instances of $X, Y, Z \in \{-1, +1\}$ ($\mathbf{E}(X, Y), \mathbf{E}(X, Z), \mathbf{E}(Y, Z)$) = (XY, XZ, YZ) , (iii) pretend these 3-tuples are coordinates (with respect to the Cartesian three-dimensional standard basis) of vertices of a convex polytope, and (iv) according to the Minkowski-Weyl “main” representation theorem [21–23] represent this polytope as its facets obtained by the hull computation [23, 24]. These facet (in)equalities represent Boole-Bell type “conditions of possible (classical) experience”.

With three dichotomic observables, such procedures result in eight three-dimensional row vectors. Four of them are linearly independent. They are interpreted as the vertices of a

correlation polytope. The row vectors, stacked on top of one another, form a 4×3 Travis [25] matrix [26]

$$T_{ij} = \begin{pmatrix} +1 & +1 & +1 \\ +1 & -1 & -1 \\ -1 & +1 & -1 \\ -1 & -1 & +1 \end{pmatrix}. \quad (5)$$

The hull computation (eg, by `pycddlib` [27], a Python wrapper of Fukuda's `cddlib` algorithm [28] implementing the Double Description Method [29]) yields the four Suppes-Zanotti-Brodi inequalities [15, 16]

$$\begin{aligned} -1 &\leq \mathbf{E}(X, Y) + \mathbf{E}(X, Z) + \mathbf{E}(Y, Z), \\ -1 &\leq -\mathbf{E}(X, Y) - \mathbf{E}(X, Z) + \mathbf{E}(Y, Z), \\ -1 &\leq \mathbf{E}(X, Y) - \mathbf{E}(X, Z) - \mathbf{E}(Y, Z), \\ -1 &\leq -\mathbf{E}(X, Y) + \mathbf{E}(X, Z) - \mathbf{E}(Y, Z). \end{aligned} \quad (6)$$

2.2 Quantum Bounds by Min-Max Calculation

The min-max calculations [14] of the associated operators $\mathbf{F}(X, Y) \pm \mathbf{F}(X, Z) \pm \mathbf{F}(Y, Z)$ with the quantum expectation \mathbf{F} as defined earlier amounts to summing up the separate terms and determining the eigensystem of these new observables. It yields quantum bounds allowing ranges bounded by

$$-3 < \mathbf{F}(X, Y) \pm \mathbf{F}(X, Z) \pm \mathbf{F}(Y, Z) < 3 \quad (7)$$

which violate the classical ones (6) by almost the greatest algebraically possible amount.

For the sake of more concrete realizations, we shall set all azimuthal angles to zero and take equidistant polar angles such that the directions of X , Y , and Z in configuration space are 0 , θ , and 2θ , respectively. Then the min-max computation associated with $\mathbf{F}(0, \theta) + \mathbf{F}(0, 2\theta) + \mathbf{F}(\theta, 2\theta)$ exhibits two eigenvalues

$$\mu_1 = -(5 + 4 \cos \theta)^{1/2} \leq -(1 + 2 \cos \theta) = \mu_2 \quad (8)$$

which, in a certain domain of θ , violate the first inequality in (6). The associated pure states are proportional to

$$\begin{aligned} |\mathbf{x}_1\rangle &= (a, b, -b, a)^T, \text{ where} \\ a &= 2(\cos \theta + 1) \sin \theta \text{ and} \\ b &= 2 \cos \theta + \cos(2\theta) + \sqrt{5 + 4 \cos \theta}, \text{ as well as} \\ |\mathbf{x}_2\rangle &= (-\sin \theta, \cos \theta, \cos \theta, \sin \theta)^T, \text{ respectively.} \end{aligned} \quad (9)$$

Note that for $\theta \rightarrow 0$ these two states converge to indecomposable vectors proportional to the Bell basis states $(0, 1, -1, 0)^T$ as well as $(0, 1, 1, 0)^T$. Indeed, for $\theta \rightarrow 0$, the two other eigenstates rendering the two eigenvalues $(5 + 4 \cos \theta)^{1/2}$, $1 + 2 \cos \theta \rightarrow 3$, converge to the remaining states in the Bell basis.

2.3 Composition of Higher-Order Distribution by Lower-Order Ones

For some “practical” application recall Specker’s story about [30] “a wise man from Ninive ... who was ... concerned almost exclusively about his daughter” and an oracle potential suitors had to cope with: “The suitors were led in front of a table on which three boxes were positioned in a row, and they were ordered to indicate which of the boxes contained a gem and which were empty. And now no matter how many times they tried, it seemed to be impossible to solve the task. After their predictions, each of the suitors was ordered to open

two boxes which they had indicated to be both empty or both not empty: it turned out each time that one contained a gem and the other did not, and, to be precise, sometimes the gem was in the first, sometimes in the second of the boxes that were opened. But how can it be possible that from three boxes neither two can be indicated as empty, nor as not empty?"

A similar scheme was mentioned by Garg and Mermin [5]: "if we have three dichotomic variables each of which assumes either the value 1 or -1 with equal probability and all the pair distributions vanish unless the members of the pair have different values . . ."

These scenarios mention three observables and strict anti-correlations between pairs of observable outcomes, such that $\mathbf{E}(X, Y) = \mathbf{E}(X, Z) = \mathbf{E}(Y, Z) = -1$. As can be readily checked by the (maximal) violation of the first Suppes-Zanotti-Brodi inequalities (6) no classical global probability distribution allows this. But quantum mechanics can "almost" provide a realization as it yields "almost perfect" anti-correlations at "almost vanishing" angles $0 < \theta \ll 1$. The "reason" for this is threefold: (i) the quantum expectation function, as mentioned earlier, is $\langle X, Y \rangle_q = \mathbf{F}(\theta_1, 0, \theta_2, 0) = -\cos(\theta_1 - \theta_2)$; (ii) the three expectation functions are complementary and therefore cannot be measured simultaneously – they have no simultaneous value definiteness; and (iii) the quantum resources exploit a four-dimensional Hilbert space with probabilities based on vectors rather than scalars.

It might be worth noting that Greenberger, Horne, and Zeilinger proposed another, adaptive, protocol involving expectations of order three and going beyond stochastic quantum violations of classical predictions [31–33] which could be rewritten as a game "people play" [34–36] in which particular quantum states allow certain players always to win whereas this is not guaranteed classically [37].

3 The Case of Four and more Observables

For completeness, we just mention that the addition of an additional variable yields the well-known Clauser-Horne-Shimony-Holt inequalities [2]. A polytope derivation can be found in Refs. [4, 7, 20]. Its quantum bound $-2\sqrt{2} \leq \mathbf{F}(W, Y) + \mathbf{F}(W, Z) + \mathbf{F}(X, Y) - \mathbf{F}(X, Z) \leq 2\sqrt{2}$ derived by Cirel'son (aka Tsirelson) [38] can be straightforwardly obtained from a min-max calculation [14] of its eigensystem. The quantum states rendering this bound can be represented by the vectors proportional to $(-1, 1, 1, 1)^T$ and $(-1, -1, -1, 1)^T$, respectively.

The polytope method can be straightforwardly scaled to derive Boolean "bounds of classical experience" for over four observables [39–41]. Their respective quantum violations can again be derived by a min-max calculation [14].

4 "Contextuality" in Context

Let me add a cautionary remark on the widely held opinion that violations of classical Boolean criteria such as the Suppes-Zanotti-Brodi inequalities suggest or even imply "contextuality". Presently the term "contextual" [42–52] is often heuristically used as "violation of some inequality that is derived by assuming classical probability distributions" [53, 54]. There are a variety of notions [55] and accompanying measures [45, 56–58] for the term "contextuality".

This "modern" quantitative use of the word can be contrasted with Bohr's synthetic suggestion of a *conditionality of phenomena* by [45, 46, 59] "the impossibility of any sharp separation between the behavior of atomic objects and the interaction with the measuring

instruments which serve to define the conditions under which the phenomena appear.” A related proposition from the realist Bell contends that [60] “the result of an observation may reasonably depend . . . on the complete disposition of the apparatus.”

In this line of thought an experimental outcome—or, in another wording, a phenomenon that should be considered as [61–63] “the comprehension of the effects observed under given experimental conditions”—is composed of contributions from both the measured object as well as from the measurement apparatus. Therefore the entire experimental configuration—effectively the experimental context—needs to be taken into account. As not all experimental contexts can be expected to be physically realizable simultaneously, not all observables can be expected to be jointly measurable. In essence this view suggests that contextuality reduces to what Bohr considers to be complementarity [45, B1–B3]; and also to Heisenberg’s related Principle of Indeterminacy or Uncertainty Principle—contextuality from indeterminacy [46, 47, 64]. (See also Glauber’s concrete quantum amplifier model [65–67], as well as the “Humpty-Dumpty” model of spin measurements [68, 69].)

However, this does not imply – and it may be even misleading to believe – that these conceivable “results of an observation” (aka outcome/event) are “dormant” properties of the object (alone) which become “visible/actuated” by some “complete disposition of the apparatus” (aka context). More precisely, there need not be any functional (in the sense of uniqueness) dependency of the outcome that originates from inherent information, causes or factors residing in or determined by the observed system; no value definite intrinsic property of the object alone. One could understand Bohr and Bell also by their insistence that the value definite properties (characterizing its physical state) of the object become “amalgamated” with (properties of) the measurement apparatus, so that an observation signals the combined information both of the object as well as of the measurement apparatus.

If one prepares a quantized system to be in a pure state formalized by a vector, then it is perfectly value definite for observable properties corresponding to that same preparation (context). But if there is a mismatch between preparation and measurement, the latter environment distorts value definiteness by an “inflow” of information from “outside of” the object. Consequently, it makes no sense to speak of any such measurement result as “being an element of physical reality” associated with the observed system alone – one has to add the (open) environment which “translates” the preparation into the measurement, thereby introducing (external with respect to the object) noise [67, 70].

5 What Propositions Support which Probabilities?

For comparing probabilities and expectations on propositional structures I maintain that in all such considerations two issues need to be distinguished as separate criteria:

- (i) Given some particular type of propositional structure (aka logics); which variety of probability distribution(s) is(are) supported by this propositional structure?
- (ii) Given two or more such varieties of probability distributions, exactly what types of probability distributions should be compared with one another? Is this not a question that needs to be settled for the particular type of systems dealt with?

I am unaware of any systematic way of answering the first question (i). One approach, motivated by Gleason-type theorems [71–76], is in terms of is Cauchy-type functional equations.

For instance, the same propositional structure may, on the one hand, support a classical hidden variable theory based on scalars as well as on (subsets of) a single Boolean algebra, while on the other hand, accommodate a quantum interpretation based on multi-dimensional vector space entities [77]. Take, for example, the Specker bug/cat's cradle [78–80], or the house/pentagon/pentagram [81–83] logics: both have a classical interpretation in terms of partitions of the sets of two-valued measures [13] as well as a faithful orthogonal representation [84, 85] as vectors.

But there are also structures that do not allow any global classical probability distribution yet support a vector coordinatization (aka faithful orthogonal representation). Examples are the Specker bug combo denoted by Γ_3 by Kochen and Specker [86] that has a nonseparable set of two-valued states. In the extreme case there exists no classical truth assignment (relative to admissibility; ie, exclusivity and completeness): take, for example, Γ_2 [86], or the logics introduced in Refs. [87, 88]. One “demarcation criterion” is the separability of the observables by two-valued states, as expressed in Kochen and Specker's Theorem 0 [86].

Conversely, there exists a plethora of propositional structures [77] that allow a partition logic interpretation, and therefore global classical probability distributions; and yet they do not support any faithful orthogonal representation, and therefore no quantization and no quantum probabilities. The simplest such example are three observables which, when depicted in a hypergraph [89–91], form a cyclical triangular structure.

It might not be too unreasonably to state that quantum “contextuality” needs only to show up if the observables satisfy Kochen and Specker's demarcation criterion by forming some propositional structure that has no classical realization and no joint probability distribution [52]. Before that one is talking about “complementary” configurations, which also allow global classical probability distributions – albeit with different probabilistic predictions yielding violations of Boole's “conditions of possible (classical) experience”.

6 Contextuality as Object Constructions

As has been mentioned earlier, most investigations into “contextuality” concentrate on the second criterion (ii) and compare discords between classical versus quantum probabilistic predictions. Thereby a presumption is an insistence that one is only willing to accept classical Boolean propositional structures representable by (power) sets as ontological entities.

This presumption is meshed with what Bell claimed to be true: that “*everything has definite properties*” [92]. That is, there is a common belief in “Omni-definiteness”, that any outcome of some measurement reflects an “inner property” or “element of physical reality” [93] of the “object” one is pretending to “measure”. No doubts are raised about the construction of this “object” which may involve important signal contributions from the measurement apparatus. Pointedly stated: the very notion of “physical object” [94] – rather than an “image of our mind” in the sense of Hertz [95, 96] – may be a naive conception that is inappropriate for situations in which one is dealing with certain types of complementary “observables” and, in particular, that have no simultaneous value definiteness [88, 97]. If, for instance, one would also be willing to contemplate vectors as fundamental ontological entities, then value definiteness ensues as pure states, and arguments based on the scarcity or even absence of classical “non-contextual” truth assignments decay into thin air.

Acknowledgements This research was funded in whole, or in part, by the Austrian Science Fund (FWF), Project No. I 4579-N.

This paper was stimulated by a discussion on “objectification” and a respective draft paper [94] of Noson Yanofsky, as well as by a question raised by Andrei Khrennikov in an email message relating to his recent paper [17].

Funding Open access funding provided by Austrian Science Fund (FWF).

Declarations

Conflict of Interests The author declares no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Bell, J.S.: On the Einstein Podolsky Rosen paradox. *Physics Physique Fizika* **1**, 195–200 (1964)
- Clauser, J.F., Horne, M.A., Shimony, A., Holt, R.A.: Proposed experiment to test local hidden-variable theories. *Phys. Rev. Lett.* **23**, 880–884 (1969)
- Eugene, P.: Wigner, On hidden variables and quantum mechanical probabilities. *Am. J. Phys.* **38**, 1005–1009 (1970)
- Froissart, M.: Constructive generalization of Bell’s inequalities. *Il Nuovo Cimento B* (11, 1971-1996) **64**, 241–251 (1981)
- Garg, A., Mermin, D.N.: Farkas’s lemma and the nature of reality: Statistical implications of quantum correlations. *Found. Phys.* **14**, 1–39 (1984)
- Pitowsky, I.: The range of quantum probability. *J. Math. Phys.* **27**, 1556–1565 (1986)
- Tsirelson, B.S.: Some results and problems on quantum Bell-type inequalities. *Hadronic Journal Supplement* **8**, 329–345 (1993)
- Svozil, K.: What is so special about quantum clicks? *Entropy* **22**, 602 (2020). arXiv:1707.08915
- Uffink, J.: Subjective probability and statistical physics. In: Beisbart, C., Hartmann, S. (eds.) *Probabilities in Physics*, pp. 25–49. Oxford University Press, Oxford (2011)
- Peres, A.: *Quantum Theory: Concepts and Methods*. Kluwer Academic Publishers, Dordrecht (1993)
- Wright, R.: Generalized urn models. *Found. Phys.* **20**, 881–903 (1990)
- Moore, E.F.: Gedanken-experiments on sequential machines. In: Shannon, C.E., McCarthy, J. (eds.) *Automata Studies*. (AM-34), pp. 129–153. Princeton University Press, Princeton (1956)
- Svozil, K.: Logical equivalence between generalized urn models and finite automata. *Int. J. Theoretical Physics* **44**, 745–754 (2005). arXiv:quant-ph/0209136
- Filipp, S., Svozil, K.: Generalizing Tsirelson’s bound on Bell inequalities using a min-max principle. *Physical Rev. Lett.* **93**, 130407 (2004). arXiv:quant-ph/0403175
- Suppes, P., Zanotti, M.: When are probabilistic explanations possible? *Synthese* **48**, 191–199 (1981)
- Brody, T.A.: The Suppes-Zanotti theorem and the Bell inequalities. *Revista Mexicana de Física* **35**, 170–187 (1989)
- Khrennikov, A.: Can there be given any meaning to contextuality without incompatibility? *Int. J. Theoretical Phys.* <https://doi.org/10.1007/s10773-020-04666-z> (2020)
- Boole, G.: On the theory of probabilities. *Philos. Trans. R. Soc. Lond.* **152**, 225–252 (1862)
- Pitowsky, I.: George Boole’s ‘conditions of possible experience’ and the quantum puzzle. *British J. Philosophy Sci.* **45**, 95–125 (1994)

20. Pitowsky, I.: Quantum Probability — Quantum Logic Lecture Notes in Physics, vol. 321. Springer-Verlag, Berlin, Heidelberg (1989)
21. Ziegler, G.M.: Lectures on Polytopes. Springer, New York (1994)
22. Schrijver, A.: Theory of Linear and Integer Programming, Wiley Series in Discrete Mathematics & Optimization. John Wiley & Sons, New York Toronto, London (1998)
23. Fukuda, K.: Frequently asked questions in polyhedral computation. Accessed on July 29th 2017 (2014)
24. Pitowsky, I.: Correlation polytopes their geometry and complexity. *Math. Program.* **50**, 395–414 (1991)
25. Travis, R.D.: The Logic of a Physical Theory, Master's thesis, Wayne State University, Detroit, Michigan, USA (1962), Master's Thesis under the supervision of David J. Foulis
26. Greechie, R.J.: Orthomodular Lattices, Ph.D. thesis, University of Florida, Florida USA (1996)
27. Troffaes, M.: `pycddlib` is a Python wrapper for Komei Fukuda's `cddlib`. Accessed on December 22th, 2020 (2020)
28. Fukuda, K.: `cdd` and `cddplus` homepage, `cddlib` package `cddlib-094h`, (2000,2017), Accessed on July 1st, 2017
29. Motzkin, T.S., Raiffa, H., Thompson, G.L., Thrall, R.M.: The double description method. In: Kuhn, H.W., Tucker, A.W. (eds.) *Contributions to Theory of Games*, vol. 2. Princeton University Press, Princeton (1953)
30. Specker, E.: Die Logik nicht gleichzeitig entscheidbarer Aussagen. *Dialectica* **14**, 239–246 (1960). english traslation at [arXiv:1103.4537](https://arxiv.org/abs/1103.4537)
31. Greenberger, D.M., Horne, M.A., Zeilinger, A.: Going beyond Bell's theorem. In: Kafatos, M. (ed.) *Bell's Theorem, Quantum Theory, and Conceptions of the Universe, Fundamental Theories of Physics*, vol. 37, pp. 69–72 (1989). Kluwer Academic Publishers, Springer Netherlands, Dordrecht. [arXiv:0712.0921](https://arxiv.org/abs/0712.0921)
32. Greenberger, D.M., Horne, M.A., Shimony, A., Zeilinger, A.: Bell's theorem without inequalities. *Am. J. Phys.* **58**, 1131–1143 (1990)
33. Mermin, D.N.: What's wrong with these elements of reality. *Phys. Today* **43**, 9–10 (1990)
34. Bouwmeester, D., Pan, J.-W., Daniell, M., Weinfurter, H., Zeilinger, A.: Observation of three-photon greenberger-horne-zeilinger entanglement. *Phys. Rev. Lett.* **82**, 1345–1349 (1999)
35. Jian-Wei Pan, D., Bouwmeester, M., Weinfurter, D.H., Zeilinger, A.: Experimental test of quantum nonlocality in three-photon Greenberger-Horne-Zeilinger entanglement. *Nature* **403**, 515–519 (2000)
36. Bacon, D.: The GHZgame. Section I of CSE 599d – Quantum Computing Quantum Entanglement and Bell's Theorem lecture notes accessed on January 9th (2021)
37. Svozil, K.: Revisiting the Greenberger-Horne-Zeilinger argument in terms of its logical structure, orthogonality, and probabilities, [arXiv:2006.14623](https://arxiv.org/abs/2006.14623) (2020)
38. Cirel'son (=Tsirel'son), B.S.: Quantum generalizations of Bell's inequality. *Lett. Math. Phys.* **4**, 93–100 (1980)
39. Pitowsky, I., Svozil, K.: New optimal tests of quantum nonlocality. *Phys. Rev. A* **64**, 014102 (2001). [arXiv:quant-ph/0011060](https://arxiv.org/abs/quant-ph/0011060)
40. Sliwa, C.: Symmetries of the Bell correlation inequalities. *Phys. Lett. A* **317**, 165–168 (2003). [arXiv:quant-ph/0305190](https://arxiv.org/abs/quant-ph/0305190)
41. Collins, D., Gisin, N.: A relevant two qbit Bell inequality inequivalent to the CHSH inequality. *J. Phys. A: Math. Gen.* **37**, 1775–1787 (2004). [arXiv:quant-ph/0306129](https://arxiv.org/abs/quant-ph/0306129)
42. Dzhafarov, E.N., Cervantes, V.H., Kujala, J.V.: Contextuality in canonical systems of random variables. *Philos. Trans. R. Soc. A: Math. Phys. Eng. Sci.* **375**, 20160389 (2017). [arXiv:1703.01252](https://arxiv.org/abs/1703.01252)
43. Abramsky, S.: Contextuality: At the borders of paradox. In: Landry, E. (ed.) *Categories for the Working Philosopher*, pp. 262–285. Oxford University Press, Oxford (2018). [arXiv:2011.04899](https://arxiv.org/abs/2011.04899)
44. Grangier, P.: Contextual objectivity: a realistic interpretation of quantum mechanics. *European J. Phys.* **23**, 331–337 (2002). [arXiv:quant-ph/0012122](https://arxiv.org/abs/quant-ph/0012122)
45. Khrennikov, A.: Bohr against Bell: complementarity versus nonlocality. *Open Physics* **15**, 734–738 (2017)
46. Jaeger, G.: Quantum contextuality in the copenhagen approach. *Philos. Trans. R. Soc. A: Math. Phys. Eng. Sci.* **377**, 20190025 (2019)
47. Jaeger, G.: Quantum contextuality and indeterminacy. *Entropy* **22**, 867 (2020)
48. Auffèves, A., Grangier, P.: Extracontextuality and extravalence in quantum mechanics. *Philos. Trans. R. Soc. A, Math. Phys. Eng. Sci.* **376**, 20170311 (2018). [arXiv:1801.01398](https://arxiv.org/abs/1801.01398)
49. Auffèves, A., Grangier, P.: Deriving born's rule from an inference to the best explanation. *Foundations of Physics* **50**, 1781–1793 (2020). [arXiv:1910.13738](https://arxiv.org/abs/1910.13738)
50. Grangier, P.: Completing the quantum formalism in a contextually objective framework, preprint [arXiv:2003.03121](https://arxiv.org/abs/2003.03121) (2020)

51. Budroni, C., Cabello, A., Gühne, O., Kleinmann, M., Larsson, J.Å.: Quantum contextuality, arXiv:2102.13036 [quant-ph] (2021)
52. Svozil, K.: Varieties of contextuality emphasizing (non)embeddability, arXiv:2103.06110 (2021)
53. Cabello, A.: Experimentally testable state-independent quantum contextuality. *Phys. Rev. Lett.* **101**, 210401 (2008). arXiv:0808.2456
54. Cabello, A.: Converting contextuality into nonlocality, arXiv:2011.13790 [quant-ph] (2020)
55. Svozil, K.: Proposed direct test of a certain type of noncontextuality in quantum mechanics. *Phys. Rev. A* **80**, 040102 (2009)
56. Svozil, K.: How much contextuality? *Nat. Comput.* **11**, 261–265 (2012). arXiv:1103.3980
57. Abramsky, S., Barbosa, R.S., Mansfield, S.: Contextual fraction as a measure of contextuality. *Phys. Rev. Lett.* **119**, 050504 (2017)
58. Kujala, J.V., Dzhaferov, E.N.: Measures of contextuality and non-contextuality. *Philos. Trans. R. Soc. A Mathematical, Phys. Eng. Sci.* **377**, 20190149, 16 (2019)
59. Bohr, N.: Discussion with Einstein on epistemological problems in atomic physics. In: Schilpp, P.A. (ed.) *Albert Einstein: Philosopher-Scientist*, pp. 200–241 (1949). *The Library of Living Philosophers*, Evanston, Ill.
60. Bell, J.S.: On the problem of hidden variables in quantum mechanics. *Rev. Mod. Phys.* **38**, 447–452 (1966)
61. Bohr, N.: The causality problem in atomic physics. In: *New Theories in Physics*, International Institute of Intellectual Co-operation, Paris, vol. 11–30 (1939). conference organized in collaboration with the International Union of Physics and the Polish Intellectual Co-operation Committee, Warsaw, May 30th–June 3rd 1938, reprinted in [62]
62. Bohr, N.: The causality problem in atomic physics. In: *Niels Bohr Collected Works*, vol. 7, pp. 299–322. Elsevier (1996)
63. Hilgevoord, J., Uffink, J.: The uncertainty principle. In: Zalta, E.N. (ed.) *The Stanford Encyclopedia of Philosophy* (2016). Metaphysics Research Lab, Stanford University winter 2016 edn.
64. Ozawa, M.: Universally valid reformulation of the Heisenberg uncertainty principle on noise and disturbance in measurement, *Phys. Rev. A* **67**. <https://doi.org/10.1103/physreva.67.042105> (2003)
65. Glauber, R.J. In: Pikes, E.R., Sarkar, S. (eds.): *Amplifiers, attenuators and the quantum theory of measurement*. Adam Hilger, Bristol (1986)
66. Glauber, R.J.: Amplifiers, attenuators and Schrödingers cat. In: *Quantum Theory of Optical Coherence*, Wiley-VCH Verlag GmbH & Co KGaA, pp. 537–576 (2007)
67. Glauber, R.J.: Amplifiers, attenuators, and Schrödinger's cat. *Ann. N. Y. Acad. Sci.* **480**, 336–372 (1986)
68. Englert, B.-G., Schwinger, J., Scully, M.O.: Is spin coherence like Humpty-Dumpty? I. Simplified treatment. *Found. Phys.* **18**, 1045–1056 (1988)
69. Schwinger, J., Scully, M.O., Englert, B.-G.: Is spin coherence like Humpty-Dumpty? II. General theory. *Zeitschrift für Physik D: Atoms, Molecules and Clusters* **10**, 135–144 (1988)
70. Svozil, K.: Quantum information via state partitions and the context translation principle. *J. Modern Optics* **51**, 811–819 (2004). arXiv:quant-ph/0308110
71. Busch, P.: Quantum states and generalized observables: a simple proof of Gleason's theorem. *Phys. Rev. Lett.* **91**, 120403, 4 (2003)
72. Caves, C.M., Fuchs, C.A., Manne, K.K., Renes, J.M.: Gleason-type derivations of the quantum probability rule for generalized measurements. *Foundations of Physics: An International Journal Devoted to the Conceptual Bases and Fundamental Theories of Modern Physics* **34**, 193–209 (2004)
73. Granström, H.: Gleason's theorem. Master's thesis, Stockholm University (2006)
74. Wright, V.J.: Gleason-type theorems and general probabilistic theories. Ph.D. thesis, University of York (2019)
75. Wright, V.J., Weigert, S.: A gleason-type theorem for qubits based on mixtures of projective measurements. *J. Phys. A Math. Theor.* **52**, 055301 (2019)
76. Wright, V.J., Weigert, S.: Gleason-type theorems from Cauchy's functional equation. *Found. Phys.* **49**, 594–606 (2019)
77. Svozil, K.: Faithful orthogonal representations of graphs from partition logics. *Soft Computing* **24**, 10239–10245 (2020). arXiv:1810.10423
78. Kochen, S., Specker, E.P.: Logical structures arising in quantum theory. In: *The Theory of Models*, Proceedings of the 1963 International Symposium at Berkeley, North Holland, Amsterdam, New York, Oxford, pp. 177–189 (1965). reprinted in Ref. [98, pp. 209–221]
79. Pitowsky, I.: Betting on the outcomes of measurements: a bayesian theory of quantum probability. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **34**, 395–414 (2003). quantum Information and Computation, arXiv:quant-ph/0208121

80. Pitowsky, I.: Quantum mechanics as a theory of probability. In: Demopoulos, W., Pitowsky, I. (eds.) *Physical Theory and its Interpretation, The Western Ontario Series in Philosophy of Science*, vol. 72, pp. 213–240. Springer, Netherlands (2006). arXiv:[quant-ph/0510095](https://arxiv.org/abs/quant-ph/0510095)
81. Geller, E.R., Greechie, R.J., Miller, F.R.: Weights on spaces. In: Enz, C.P., Mehra, J. (eds.) *Physical Reality and Mathematical Description*. D. Reidel Publishing Company, Springer Netherlands, Dordrecht, Holland, pp. 167–192 (1974)
82. Wright, R.: The state of the pentagon. A nonclassical example. In: Marlow, A.R. (ed.) *Mathematical Foundations of Quantum Theory*, pp. 255–274. Academic Press, New York (1978)
83. Klyachko, A.A., Can, M.A., Binicioğlu, S., Shumovsky, A.S.: Simple test for hidden variables in spin-1 systems. *Phys. Rev. Lett.* **101**, 020403 (2008). arXiv:[0706.0126](https://arxiv.org/abs/0706.0126)
84. Lovász, L.: On the Shannon capacity of a graph. *IEEE Trans. Inf. Theory* **25**, 1–7 (1979)
85. Lovász, L., Saks, M., Schrijver, A.: Orthogonal representations and connectivity of graphs. *Linear Algebra Appl.* **114–115**, 439–454 (1989). special Issue Dedicated to Alan J. Hoffman
86. Kochen, S., Specker, E.P.: The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics (now Indiana University Mathematics Journal)* **17**, 59–87 (1967)
87. Cabello, A., Estebaranz, J.M., García-Alcaine, G.: Bell-Kochen-Specker theorem: A proof with 18 vectors. *Phys. Lett. A* **212**, 183–187 (1996). arXiv:[quant-ph/9706009](https://arxiv.org/abs/quant-ph/9706009)
88. Abbott, A.A., Calude, C.S., Svozil, K.: A variant of the Kochen-Specker theorem localising value indefiniteness. *J. Math. Phys.* **56**, 102201 (2015). arXiv:[1503.01985](https://arxiv.org/abs/1503.01985)
89. Greechie, R.J.: Orthomodular lattices admitting no states. *Journal of Combinatorial Theory Series A* **10**, 119–132 (1971)
90. Kalmbach, G.: *Orthomodular Lattices* London Mathematical Society Monographs, vol. 18. Academic Press, London and New York (1983)
91. Bretto, A.: *Hypergraph theory, Mathematical Engineering*. Springer, Cham, Heidelberg, New York, Dordrecht, London, pp. xiv+119 (2013)
92. Bertlmann, R.A.: Real or not real that is the question *The European Phys. J. H* **45**, 205–236 (2020). arXiv:[2005.08719](https://arxiv.org/abs/2005.08719)
93. Einstein, A., Podolsky, B., Rosen, N.: Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.* **47**, 777–780 (1935)
94. Yanofsky, N.S.: The mind and the limitations of physics, preprint, accessed on January, 14, 2021 (2019)
95. Hertz, H.: *Prinzipien der Mechanik*, Johann Ambrosius Barth (Arthur Meiner, Leipzig) mit einem Vorwort von H. von Helmholtz (1894)
96. Hertz, H.: *The principles of mechanics presented in a new form*, (MacMillan and Co., Ltd., London and New York) with a foreword by H. von Helmholtz, translated by D. E. Jones and J. T. Walley (1899)
97. Pitowsky, I.: Infinite and finite Gleason’s theorems and the logic of indeterminacy. *J. Math. Phys.* **39**, 218–228 (1998)
98. Specker, E.: *Selecta*. Basel, Birkhäuser (1990)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.