



Universal gates with wires in a row

Ville Salo¹

Received: 31 August 2020 / Accepted: 8 August 2021 / Published online: 24 September 2021
© The Author(s) 2021

Abstract

We give some optimal size generating sets for the group generated by shifts and local permutations on the binary full shift. We show that a single generator, namely the fully asynchronous application of the elementary cellular automaton 57 (or, by symmetry, ECA 99), suffices in addition to the shift. In the terminology of logical gates, we have a single reversible gate whose shifts generate all (finitary) reversible gates on infinitely many binary-valued wires that lie in a row and cannot (a priori) be rearranged. We classify pairs of words u, v such that the gate swapping these two words, together with the shift and the bit flip, generates all local permutations. As a corollary, we obtain analogous results in the case where the wires are arranged on a cycle, confirming a conjecture of Macauley-McCammond-Mortveit and Vielhaber.

Keywords Reversible gates · Groups

1 Introduction

In this paper, we study the subgroup G of the self-homeomorphism group of $\{0, 1\}^{\mathbb{Z}}$ generated by shifts and local permutations that only modify a bounded set of cells (which can be seen as reversible finitary logical gates) on the set of all bi-infinite binary sequences, studied previously in [2] under the name $OB(2, 1)$ (as a subgroup of the larger group $RTM(2, 1)$).

We prove a result about generators for the group G : In Sect. 7, we characterize those pairs of words u, v such that the homeomorphism swapping u and v at the origin, together with the shift and the bit flip, generates all of G . In the process, we identify some natural subgroups of G in Sect. 5, with constraints related to linearity, and to (group-)geometric phenomena such as one-sided information flow.

Research supported by the Academy of Finland grant 2608073211.

✉ Ville Salo
vosalo@utu.fi

¹ University of Turku, Turku, Finland

We say a set of local permutations is a *universal gate set* if it, together with the shift, generates G . As an application of the result about word swaps, we prove there exists a singleton universal gate set by showing that the co-called asynchronous ECA 57 gate generates both a universal word swap and the bit flip. In particular, it follows that G is 2-generated.

In [7,13], it is asked whether asynchronous ECA 57 is universal in the case when wires are arranged on a cycle $\mathbb{Z}/n\mathbb{Z}$. We solve this question by showing that universality on \mathbb{Z} implies universality on $\mathbb{Z}/n\mathbb{Z}$ for all large enough n in Sect. 4. We include copy-pasteable implementations of standard gates in terms of shifts of asynchronous ECA 57 in the appendices.

We find the group G interesting for several reasons: From the perspective of computation and logical gates, G is precisely the computation you can do with reversible gates if your infinitely many wires are arranged in a row. It differs from the groupoid [15] and clone formalisms [3] most concretely in the aspect that the model has infinitely many borrowed bits (in the sense of [14]) which arise naturally from the infiniteness of \mathbb{Z} . Reordering of the wires is not built into the formalism, which strengthens the notion of universality.

The group G seems like a natural simplest group that can be used for the geometric group theory of reversible logic gates.

The author finds G and its variants useful in the context of symbolic dynamics, and has applied G and its local embeddings (in the LEF sense [5]) in several works on automorphism groups, e.g., such groups appear in disguise in [9,10].

2 Definitions

We assume $0 \in \mathbb{N}$ and in $f \circ g = fg$, g is applied first. All intervals $[a, b]$, as well as half-open and open ones, are interpreted inside \mathbb{Z} . Words are 0-indexed by default. For two words $u, v \in \{0, 1\}^n$ write $D(u, v) \subset [0, n - 1]$ for the set of coordinates where u and v differ, and $d(u, v) \in \{0, 1\}^n$ for the characteristic function of $D(u, v)$.

We consider $X = \{0, 1\}^{\mathbb{Z}}$ with the product topology arising from the discrete topology of $\{0, 1\}$. The *shift* is $\sigma : X \rightarrow X$ defined by $\sigma(x)_i = x_{i+1}$. A power σ^n is also informally referred to as a *shift*, which should not cause confusion. For $n \geq 1$, the *n-periodic points* are the points $x \in X$ satisfying $\sigma^n(x) = x$. The set of *n-periodic points* is denoted by X_n . The *shifts* of a function are its conjugates by shifts σ^n .

There is an obvious way to see X as a vector space over the two-element field, and addition and terms such as *linear* and *affine* for maps on X are interpreted with respect to this structure. The subset $X_0 = \{x \in X \mid \exists R : \forall |i| \geq R : x_i = 0\}$ is a subspace. An *affine translation* refers to a function $t : X \rightarrow X$ of the form $x \mapsto x + w$, with $w \in X_0$.

We assume basic knowledge of group theory, such as split extensions and semidirect products, see any standard reference [8]. A locally finite group is one whose finitely generated subgroups are finite. We use the $\langle F_1, \dots, F_n \rangle$ notation to denote the smallest subgroup of a group G (deduced from the context) containing the elements equal to F_i (when $F_i \in G$) or found in F_i (when F_i is a set).

3 G

Definition 1 The group G is defined as the subgroup of the self-homeomorphism group of X generated by the shift σ and homeomorphisms $f : X \rightarrow X$ satisfying

$$\exists r : \forall i : (|i| \geq r \implies f(x)_i = x_i). \tag{1}$$

We make some basic observations about the structure of this group G (see also [2,9]).

We call self-homeomorphisms f of X satisfying Eq. 1 *local permutations, inert¹ elements of G* or *gates*. A set of gates F is *universal* if $G = \langle F, \sigma \rangle$.

For an inert element $f \in G$, the finite set of coordinates that may change under the application of f is called its *support*. Of more importance than the support and the number r is the R in the following lemma, which states that if f is inert in G , then not only does it change only finitely many symbols in a given point of X , but it also *looks at* only finitely many coordinates.

Lemma 1 *An element $f \in G$ is inert if and only if there exists a strong radius R and a local rule $\hat{f} : \{0, 1\}^{[-R, R]} \rightarrow \{0, 1\}^{[-R, R]}$ such that $f(x)_i = x_i$ for all $|i| > R$, and $f(x)_i = \hat{f}(x_{[-R, R]})_i$ for $i \in [-R, R]$.*

Proof Clearly, any R and \hat{f} define an inert element of G .

Suppose then $f \in G$ is inert, and let r be as in the definition. Since f is continuous, there exists R such that $f(x)_{[-r, r]}$ is determined by $x_{[-R, R]}$, where we may assume $R \geq r$. Since $f(x)_i = x_i$ when $|i| \in [-R, -r - 1] \cup [r + 1, R]$, in fact $f(x)_{[-R, R]}$ is also determined by $x_{[-R, R]}$. So we can pick $\hat{f}(w) = f(x^w)_{[-R, R]}$ where $x^w \in \{0, 1\}^{\mathbb{Z}}$ is any point with $x^w_{[-R, R]} = w$. Since f is a homeomorphism, it is in particular a bijection, so \hat{f} is also a bijection. □

Define the *strong support* of inert f as the set of coordinates that can be changed by f , and whose change may affect another coordinate, which is finite by the previous lemma.

The number R can be arbitrarily larger than r , even if $r = 0$.

Remark 1 We note that (by easy counterexamples) the inert elements of G are not the same as self-homeomorphisms of X which have clopen support in the usual sense of homeomorphisms (where the support of a homeomorphism is the closure of the complement of its set of fixed points). Neither are they the same as self-homeomorphisms of X which always modify only finitely many coordinates, or ones that always modify only a bounded number of coordinates. By definition, they are precisely the self-homeomorphisms of X which always modify a bounded *set* of coordinates.

Lemma 2 *Elements of G can be written in a unique way as $\sigma^n \circ f$ with f inert. There is a unique homomorphism $\alpha : G \rightarrow \mathbb{Z}$ satisfying $\alpha(\sigma) = 1$. The kernel of α consists of the inert elements. The group G is (locally finite)-by- \mathbb{Z} , more precisely a semidirect product of \mathbb{Z} acting on the locally finite group of inert elements.*

¹ This word is by analogue with subshift automorphism group terminology [6].

The inert map g in $f = \sigma^n \circ g$ is called the *inert part* of f .

Proof Since the local permutations form a group themselves, and this group is normalized by σ (i.e., conjugation of an inert element by σ gives an inert element), it follows that every element of G is of the form $\sigma^n \circ f$ where f is inert, and clearly n is uniquely determined, after which f is clearly uniquely determined. Any homomorphism to \mathbb{Z} maps any inert f to 0 (since inert elements are of finite order), so by the normal form, $\sigma \mapsto 1$ uniquely determines the homomorphism.

It follows from this that $\ker \alpha$ consists of precisely the inert elements, and inert elements clearly form a locally finite group, since the support of the composition of two inert element is at most the union of the support of the generators. This gives G the structure of a (locally finite)-by- \mathbb{Z} group, and since the homomorphism α is split, G is in fact a semidirect product of \mathbb{Z} acting on the locally finite group of inert elements. \square

We end this section with some general discussion.

The inert elements can be naturally considered reversible gates by Lemma 1, as they perform an operation on finitely many wires of the input independently of the other wires. From this perspective, the shift should be seen as either a technical tool or as an *internal symmetry* of the group—shifts are not necessarily computationally meaningful, but *conjugation by a shift* is very meaningful, as it simply means applying a gate to another contiguous set of wires, and this is precisely what it means to say that G is a semidirect product of \mathbb{Z} acting on the inert elements.

Though we do not do so here, from this point of view of logical gates it would be natural to see G not just as a group but as a group where the shifts are part of the structure. A group H with a fixed epimorphism $\pi : H \rightarrow \mathbb{Z}$ is often called (\mathbb{Z} -) *indicable*, so one could sensibly call a group with a fixed split epimorphism $\pi : H \rightarrow \mathbb{Z}$ *strongly \mathbb{Z} -indicable*, with the understanding that π and the right inverse can be taken to be part of the structure, if needed. In particular, if the \mathbb{Z} -embedding giving the right inverse is taken part of the structure, then substructures of G always contain the shifts.

We mention some standard notions in category theory with are related: One can see a group H with a preferred \mathbb{Z} -subgroup as a pair (H, ψ) where $\psi : \mathbb{Z} \rightarrow H$ is a homomorphism (that is part of the structure). Such pairs form the *under category*, also known as the *coslice category*, $\mathbb{Z} \downarrow \text{Grp}$. This does not enforce the semidirect product structure in the sense that the \mathbb{Z} picked out by ψ can behave very badly in other objects of this category, but this does enforce the property that subobjects of G automatically contain the shifts (note that monomorphisms in Grp are the injections). Dually, one can see G as an indicable group with a fixed $\pi : G \rightarrow \mathbb{Z}$, i.e., an element of the slice category $\text{Grp} \downarrow \mathbb{Z}$, but this does not force subobjects of G to contain the shifts.

4 G_n

Definition 2 For any $n \in \mathbb{N}$, write $G_n = \text{Alt}(\{0, 1\}^n)$.

The notation is meant to imply that G_n can be seen as a modulo n variant of G . In this section, we develop an explicit algebraic connection.

For inert f , the *strong shift-invariant radius* of f is the minimal R among strong radii of conjugates $\sigma^m \circ f \circ \sigma^{-m}$, and for general $f \in G$, the strong shift-invariant radius is the strong shift-invariant radius of the inert part. The importance of the strong shift-invariant radius is the following.

Definition 3 Let $f = \sigma^m \circ g \in G$, with g inert with strong shift-invariant radius R . Suppose $n > 2R + 1$. Then, for $x \in X_n$ define

$$\tilde{f}_n(x) = \sigma^m \circ \lim_{k \rightarrow \infty} \left(\prod_{i=-k}^k \sigma^{-in} \circ g \circ \sigma^{in} \right) (x)$$

For $w \in \{0, 1\}^n$ define $f_n(w) = \tilde{f}_n(\dots w w w . w w w \dots)_{[0, n-1]}$.

In particular, $\sigma_n(aw) = wa$ for $a \in \{0, 1\}$, $w \in \{0, 1\}^{n-1}$.

Since $n > 2R + 1$, the limit is well-defined as the maps $\sigma^{-in} \circ g \circ \sigma^{in}$ commute pairwise (for distinct values of i) and every $j \in \mathbb{Z}$ is in the support of only finitely many of them (in fact only one of them). Observe that \tilde{f}_n commutes with σ^n , and thus preserves the set of n -periodic points. The action of \tilde{f}_n on the n -periodic points of $\{0, 1\}^{\mathbb{Z}}$ fully determines the action of f_n on $\{0, 1\}^n$, and vice versa.

Remark 2 For completeness, we give a formula for f_n . Let $f \in G \setminus \{\text{id}\}$ be inert, let the strong shift-invariant radius of f be R , and let $|m|$ be minimal such that $\sigma^m \circ f \circ \sigma^{-m}$ has strong radius R , and let $\hat{f} : \{0, 1\}^{[-R, R]} \rightarrow \{0, 1\}^{[-R, R]}$ be the local rule for $\sigma^m \circ f \circ \sigma^{-m}$. If $[m - R, m + R] \equiv [a, b] \pmod n$ for some $a, b \in [0, n - 1]$, then define

$$f_n(w) = w_{[0, a]} \cdot \hat{f}(w_{[a, b]}) \cdot w_{[b, n-1]}$$

and if $[m - R, m + R] \equiv [0, a] \cup (n - 1 - b, n - 1] \pmod n$ for some $a, b \in [0, n - 1]$ (where possibly $a = 0$ or $b = 0$) then define

$$f_n(w) = \hat{f}(u)_{[b, 2R+1]} \cdot w_{[a, n-1-b]} \cdot \hat{f}(u)_{[0, b]}$$

where $u = w_{(n-1-b, n-1]} \cdot w_{[0, a]}$. For $f = \sigma^k \circ g$ with g inert, we have $f_n = \sigma_n^k \circ g_n$

Note that if m, R, \hat{f} are as above, then f applies its local rule \hat{f} “at m ”, i.e., in the coordinates $[m - R, m + R]$.

Lemma 3 Let $f = \sigma^k \circ g$ with g inert with strong shift-invariant radius R . If $n > \max(2, 2R + 1)$, then $f_n \in G_n$.

Proof Consider first inert g with strong shift-invariant radius R . Since $n > 2R + 1$, by the formulas of Remark 2, g_n permutes only the contents of some $2R + 1$ coordinates of $[0, n - 1]$, so in fact g_n performs $2^{n-(2R+1)}$ “copies” of the same permutation depending on the values of the other $n - 2R - 1$ coordinates, thus it is a composition

of an even number of permutations with disjoint supports, all having the same parity, thus $g_n \in G_n$.

The shift σ_n is a permutation of the n coordinates of a given word, so it can be written as a product of swaps of adjacent coordinates. When $n \geq 3$, coordinate swaps are even by the argument of the above paragraph, and thus σ_n is even as a composition of even permutations. Thus for $f = \sigma^k \circ g$ with g inert, we have $f_n = \sigma_n^k \circ g_n \in G_n$. \square

The proof shows that $\sigma_n \in G_n$ whenever $n \geq 3$. We mention a combinatorial consequence of this: The parity of a permutation on a set X is the parity of $|X| - c$ where c is the number of cycles in the cycle decomposition. Thus, the sign of σ_n is $2^n - p_n$ where p_n is the number of orbits of words of length n under the cyclic shift. The sequence p_n is well-known, it is sequence A000031 in the OEIS, and a standard formula for it is $p_n = \frac{1}{n} \sum_{d|n} \phi(d) 2^{n/d}$ (where ϕ is the Euler totient function). We are stating precisely that p_n is even for all $n \geq 3$. Since $p_1 = 2$ and $p_2 = 3$, p_2 is the only odd value in the sequence p_1, p_2, p_3, \dots

Lemma 4 *Let g be inert with strong shift-invariant radius R , and suppose $n > \max(2, 2R + 1)$. Then, $g_n \circ \sigma_n^m = \sigma_n^m \circ (\sigma^{-m} \circ g \circ \sigma^m)_n$ for all $m \in \mathbb{Z}$.*

Proof We prove instead the equivalent equality $\bar{g}_n \circ \bar{\sigma}_n^m = \bar{\sigma}_n^m \circ \overline{(\sigma^{-m} \circ g \circ \sigma^m)}_n$ for the actions on n -periodic points. Letting $x \in X_n$, we have

$$\begin{aligned} \bar{g}_n \circ \bar{\sigma}_n^m(x) &= \lim_{k \rightarrow \infty} \left(\prod_{i=-k}^k \sigma^{-in} \circ g \circ \sigma^{in} \right) (\sigma^m(x)) \\ &= \sigma^m \circ \lim_{k \rightarrow \infty} \left(\prod_{i=-k}^k \sigma^{-in} \circ (\sigma^{-m} \circ g \circ \sigma^m) \circ \sigma^{in} \right) (x) \\ &= \bar{\sigma}_n^m \circ \overline{(\sigma^{-m} \circ g \circ \sigma^m)}_n(x) \end{aligned}$$

where σ^m waddles over the finite expressions by using abelianity of \mathbb{Z} and by conjugating g s, and the result stays correct in the limit since σ is continuous. \square

The map $f \mapsto f_n$ does not give a homomorphism from G to G_n (it is not even well-defined for all elements of G), and even if we induce a mapping from G to G_n using a finite generating set of G , it will typically not give a homomorphism (the group G is not residually finite).

As long as information is “not passed around the cycle,” we see the same relations in both G and G_n , through the mapping constructed above. The following lemma states a weak version of this fact:

Lemma 5 *Let $f^1, \dots, f^k \in G$ with $f^i = \sigma^{\ell_i} \circ g^i$, where g^i is inert. Let g^i have strong shift-invariant radius R_i and let $|m_i|$ be minimal such that $\sigma^{-m_i} \circ g^i \circ \sigma^{m_i}$ has strong radius R_i . Let $t = \sum_j |\ell_j|$. Suppose that for some h , $[m_i - t - R_i, m_i + t + R_i] \subset [h, h + n - 1]$ for all i . Then, $(f^1 \circ \dots \circ f^k)_n = f_n^1 \circ \dots \circ f_n^k$.*

Proof First, we reduce to the inert case.

Observe that by iterating Lemma 4 the RHS of the equality is

$$f_n^1 \circ \dots \circ f_n^k = \sigma_n^{\ell_1} \circ g_n^1 \circ \dots \circ \sigma_n^{\ell_k} \circ g_n^k = \sigma_n^{\sum_j \ell_j} \circ h_n^1 \circ \dots \circ h_n^k$$

where $h^i = \sigma^{-t_i} \circ g^i \circ \sigma^{t_i}$ for some $|t_i| \leq t$ (where t_i is some partial sum). In particular, the h^i are inert with strong shift-invariant radius R_i . The LHS on the other hand is

$$(f^1 \circ \dots \circ f^k)_n = (\sigma^{\sum_j \ell_j} \circ h^1 \circ \dots \circ h^k)_n = \sigma_n^{\sum_j \ell_j} \circ (h^1 \circ \dots \circ h^k)_n$$

by definition.

We may thus assume $\ell_i = 0$ for all i , by replacing each m_i with $m_i + t_i$ (after which the corresponding assumption $[m_i - R_i, m_i + R_i] \subset [h, h + n - 1]$ still holds), and by cancelling the common part $\sigma_n^{\sum_j \ell_j}$. We have then reduced the problem to the case where the $f^i = g^i = h^i$ are inert.

Let us now prove the equivalent equality

$$\overline{(f^1 \circ \dots \circ f^k)_n} = \bar{f}_n^1 \circ \dots \circ \bar{f}_n^k.$$

on X_n . For this, observe that the action of the maps f^i on the interval $[h, h + n - 1]$ is precisely the same as the action of the maps \bar{f}_n^i , and the action of the \bar{f}_n^i on any contiguous sequence of n integers determines the action. Thus, the map $f^i \mapsto \bar{f}_n^i$ even induces a homomorphism from $\langle f^1, \dots, f^k \rangle$ to $\langle \bar{f}_n^1, \dots, \bar{f}_n^k \rangle$, in particular we have $(f^1 \circ \dots \circ f^k)_n = \bar{f}_n^1 \circ \dots \circ \bar{f}_n^k$ as desired. \square

A finitely generated group H with word norm $|\cdot|_H$ with respect to some finite generating set is *LEF* (locally embeddable in finite groups) if for all n there exists an injective map $\phi : H \rightarrow F$ where F is a finite group, such that for all $h, h' \in H$ with $|h|_H, |h'|_H \leq n$, we have $\phi(h)\phi(h') = \phi(hh')$. See [5]. The above lemma shows that G is LEF.

5 How not to generate G (subgroups)

We now define some important self-homeomorphisms of X (not all in G) and make simple observations about subgroups they generate. For the purpose of universality, the importance of understanding subgroups is of course that if a set $F \subset G$ is contained in a proper subgroup $F \subset H < G$, then also $\langle F \rangle \leq H < G$, preventing F from generating G .

The *reversal* $R : X \rightarrow X$ defined by $R(x)_i = x_{-i}$ is not an element of G , but normalizes G inside the self-homeomorphism group of X (so G has index 2 in $\langle R, G \rangle$). The *swap* is $s \in G$ defined by $s(x)_0 = x_1, s(x)_1 = x_0, \forall i \notin \{0, 1\} : s(x)_i = x_i$. The *k-CNOT* is $c^k \in G$ defined by $c^k(x)_0 = 1 - x_0 \iff x_{[1,k]} = 1^k$ and $\forall i \neq 0 : c^k(x)_i = i$. The map c^0 inverts the symbol at the origin, and is usually called the *NOT* gate in the theory of logical gates. We also refer to it as the *flip*. The map c^1

is linear, and is called the *CNOT* or *controlled NOT*. It is useful to keep in mind the identity

$$c^1 \circ (\sigma^{-1} \circ (R \circ c^1 \circ R) \circ \sigma) \circ c^1 = s \tag{2}$$

which states the swap in terms of CNOTs. The gate c^2 is called the *Toffoli gate*.

We make some observations about subgroups generated by these elements. Write $\text{Sym}_{<\infty}(\mathbb{Z})$ for the group of finite-support permutations on \mathbb{Z} .

Lemma 6

$$\begin{aligned} \langle s, \sigma \rangle &= \{f \in G \mid \exists \pi \in \text{Sym}_{<\infty}(\mathbb{Z}), n \in \mathbb{Z} : \forall i, x : f(x)_i = x_{\pi(i)+n}\} \\ \langle c^0, \sigma \rangle &= \{f \in G \mid \exists c \in X_0, n \in \mathbb{Z} : \forall x : f(x) = \sigma^n(x) + c \cong \mathbb{Z}_2 \wr \mathbb{Z}\} \\ \langle c^1, Rc^1R, \sigma \rangle &= \{f \in G \mid \forall x, y : f(x+y) = f(x) + f(y)\} \\ \langle c^0, c^1, Rc^1R, \sigma \rangle &= \{f \in G \mid \exists c \in X_0 : \forall x, y : f(x+y) = f(x) + f(y) + c\} \end{aligned}$$

Proof (Proof sketch) In each case, it is straightforward to show that the generators are contained in the RHS, and that the RHS is a group. To see that all of the group is generated, for the first equality we need that every permutation can be decomposed into swaps. The second is clear, and this is a standard realization of the lamplighter group $\mathbb{Z}_2 \wr \mathbb{Z}$. The third follows from (2) and the fact elementary matrices generate matrix groups. The RHS of the fourth equality is equivalent to f being the sum of a linear map and a constant in X_0 , and the subgroup $\langle c^0, \sigma \rangle$ allows addition of arbitrary constants. □

The first group is a standard example of a finitely generated LEF group that is not residually finite [5] (G itself is also such an example).

The following subgroups are important in the classification of universal word swap gates, and we include a more complete treatment.

Definition 4 If $V \leq X_0$ is a shift-invariant vector space, let

$$G_V = \{f \in G \mid \exists n \in \mathbb{Z}, v \in X_0 : \forall x : f(x) \in \sigma^n(x) + v + V\}.$$

Define

$$G_R = \{\sigma^n \circ f \mid f \in \ker(\alpha), n \in \mathbb{Z}, \exists (g_i)_{i \in \mathbb{Z}} : \forall x \in X : f(x)_i = g_i(x_{(-\infty, i]})\}.$$

for some functions $g_i : \{0, 1\}^{(-\infty, i]} \rightarrow \{0, 1\}$. Define $G_L = R \circ G_R \circ R$

Informally, G_V is the subgroup of G for which the cosets of V are a system of blocks of imprimitivity, and these cosets are permuted by an affine translation. The interpretation of G_R is that information can only travel to the right, though literally this only holds in the inert part.

Example 1 For all k , we have $c^k \in G_L$ and $Rc^kR \in G_R$. We have $c^0 \in G_V$ for any V , and if π is the permutation $(00\ 11)(01)(10)$ of $\{0, 1\}^2$, in cycle notation, then the map $f(x) = x_{(-\infty,0)} \cdot \pi(x_{[0,1]})x_{[2,\infty)}$ is in G_V where $V = \{x \in X_0 \mid \sum_i x_i \equiv 0 \pmod 2\}$.

This map $f \in G_V$ (which is $f_{00,11}$ in the notation of Sect. 7), is not linear, but is affine. The map that performs the permutation $(000\ 111)$ is in $G_{V'}$ for a proper subspace $V' < X_0$, but is not even affine. ◦

Lemma 7 *Let $V < X_0$ be a shift-invariant proper subspace. Then, G_V, G_L, G_R are proper subgroups of G .*

Proof First, we show G_V is a subgroup. Let us call the vector v in the definition the *offset* and n the *shift*. Let $f \in G_V$ with offset v and shift n . Then for any x , $f(x+V) \subset \sigma^n(x) + \sigma^n(V) + v + V = \sigma^n(x) + v + V$ by shift-invariance of V . Since f is a homeomorphism, cosets of V partition X , and $\sigma^n(x) + v = \sigma^n(y) + v \iff x - y \in V$, we have $f(x+V) = \sigma^n(x) + v + V$. Thus,

$$f(\sigma^{-n}(x+v) + V) = \sigma^n(\sigma^{-n}(x+v)) + v + V = x + V$$

for any x , thus $f^{-1}(x+V) = \sigma^{-n}(x) + \sigma^{-n}(v) + V$, thus $f^{-1}(x) \in \sigma^{-n}(x) + \sigma^{-n}(v) + V$ for all x , so $f^{-1} \in G_V$ with offset $\sigma^{-n}(v)$ and shift $-n$.

Suppose then $f, g \in G_V$, and let the offsets be u, v , and shifts m, n , respectively. Then, $f \circ g(x) \in f(\sigma^n(x) + v + V) \in \sigma^{m+n}(x) + \sigma^m(v) + u + V$, so $f \circ g$ admits offset $\sigma^m(v) + u$ and shift $m + n$, thus $f \circ g \in G_V$.

To see that G_V is a proper subgroup, observe that since V is shift-invariant and $V \neq X_0$, V does not contain the vector ${}^\omega 0.10^\omega$. We in fact have $c^1 \notin G_V$ for the CNOT c^1 : since $c^1(0^\mathbb{Z}) = 0^\mathbb{Z}$ and c^1 is inert, if $c^1 \in G_V$ then necessarily the offset v in the definition would satisfy $v \in V$. But then, $f({}^\omega 0.010^\omega) = {}^\omega 0.110^\omega \in {}^\omega 0.010^\omega + V$ implies ${}^\omega 0.10^\omega \in V$, a contradiction.

Next, we show G_R is a subgroup. Let us call the sequence $(g_i)_{i \in \mathbb{Z}}$ appearing in the definition the *rule sequence*. Let $\sigma^{-n} \circ f \in G_R$ with rule sequence $(g_i)_{i \in \mathbb{Z}}$. Then, it has inverse $\sigma^{-n} \circ (\sigma^n \circ f^{-1} \circ \sigma^{-n})$, and we only need to find a rule sequence for $\sigma^n \circ f^{-1} \circ \sigma^{-n}$.

First, we show that f^{-1} has a rule sequence. Observe that necessarily $g_i(y) = y_i$ for all but finitely many i (since f is inert). Observe then that we have $g_i(ya) \neq g_i(yb)$ for all $i \in \mathbb{Z}$ and $y \in \{0, 1\}^{(-\infty, i-1]}$, $a \neq b$. There are several ways to see this: it follows from the fact that $f^m = \text{id}$ for some m (since f is inert), or from a direct counting argument using the fact $g_i(y) = y_i$ for large $|i|$.

Now, we can build, by induction, a sequence of functions $h'_i : \{0, 1\}^{(-\infty, i]} \rightarrow \{0, 1\}^{(-\infty, i]}$ such that $h'_i(g_i(y)) = y_i$ for all $y \in \{0, 1\}^{(-\infty, i]}$, and the functions agree in the sense $h'_{i+1}(ya)_{(-\infty, i]} = h'_i(y)$ for $y \in \{0, 1\}^{(-\infty, i]}$, $a \in \{0, 1\}$: Pick $h_i = \text{id}$ for all small enough i , and $h'_{i+1}(ya) = h'_i(y)b$ where b is the unique letter such that $g_{i+1}(h'_i(y)b) = a$. Now, $h_i(y) = h'_i(y)_i$ defines a rule sequence for f^{-1} .

Now,

$$\sigma^n f^{-1} \sigma^{-n}(x)_i = f^{-1} \sigma^{-n}(x)_{i+n} = h_{i+n}(\sigma^{-n}(x)_{(-\infty, i+n]}) = h_{i+n}(x_{(-\infty, i]}).$$

where we shift the domain of h_{i+n} to $h_{i+n} : \{0, 1\}^{(-\infty, i]} \rightarrow \{0, 1\}$ with the obvious interpretation. Thus, $(h_{i+n})_{i \in \mathbb{Z}}$ is a rule sequence for $\sigma^n \circ f^{-1} \circ \sigma^{-n}$.

Suppose then that $\sigma^n \circ f, \sigma^m \circ f' \in G_V$ with f, f' inert. The above paragraph shows that the set of inert elements having a rule sequence is closed under conjugation by the shift, so by $\sigma^m f' \sigma^n f = \sigma^{m+n} (\sigma^{-n} f' \sigma^n) f$, we only need to show that if f' and f have rule sequences, then so does $f' \circ f$. This follows from

$$f' f(x)_i = g'_i(f(x)_{(-\infty, i]}) = g'_i(y)$$

where $y_j = g_j(x_{(-\infty, j]})$ for $j \leq i$. This expression only looks at the values of x in $x_{(-\infty, i]}$, thus defines a rule sequence.

To see that G_R is a proper subgroup, observe that the swap $s \notin G_R$.

The proof for G_L is symmetric. □

6 How to generate G and G_n

The following is essentially folklore in the theory of reversible circuits.

Lemma 8 *Let $n \geq 4$. Letting*

$$S = \{(\sigma^{-m} \circ s \circ \sigma^m)_n \mid m \in [0, n - 2]\},$$

we have $G_n = \langle S, c_n^0, c_n^2 \rangle$. In particular,

$$G_n = \langle s_n, c_n^0, c_n^2, \sigma_n \rangle.$$

Proof The maximal minimal strong shift-invariant radius among gates in $S \cup \{c^0, c^2\}$ is 1 (for c^2), so the expressions are well-defined and we have $S \cup \{c_n^0, c_n^2\} \subset G_n$ by Lemma 3.

It is known that every even permutation on $\{0, 1\}^n$ can be written as a finite composition of applications of the NOT gate and Toffoli gate when we are allowed to apply these gates to arbitrary 3-tuples of wires (this is the folklore part, see e.g., [4, 14]). The swaps in S allow arbitrary permutations of the coordinates, so we can conjugate any 3-tuple of coordinates to the support $\{0, 1, 2\}$ of the Toffoli gate c_n^2 , and similarly we can apply the NOT gate in any coordinate.

For the second claim, observe that the gates in S are conjugate to s_n by powers of σ_n . □

Lemma 9

$$G = \langle c^2, s, c^0, \sigma \rangle = \langle c^2, c^1, Rc^1R, c^0, \sigma \rangle$$

Proof It is enough to show that the local permutations are generated, and by conjugating with powers of σ (which is in both claimed generating sets), it is enough to show that those with strong support contained in $[0, n - 1]$ are generated, for arbitrarily large n .

By the previous lemma, the leftmost generating set can perform every even permutation of the word on the interval $[0, n - 1]$ for $n \geq 4$: none of the generators of G_n in the first generating set in the Lemma 8 send information over the borders of the interval $[0, n - 1]$, so the action of the subgroup generated by

$$\{\sigma^{-m} \circ s \circ \sigma^m \mid m \in [0, n - 2]\} \cup \{c^0, c^2\}$$

precisely simulates the action of G_n of the corresponding generators in the lemma.

As in Lemma 3, we see that every permutation of $\{0, 1\}^n$ is even when seen as a permutation of $\{0, 1\}^{n+1}$ where the rightmost bit is never changed or looked at (i.e., it is a *borrowed bit* in the terminology of [14]), so the leftmost generating set indeed generates all of G .

The second equality follows from (2). □

In logical gates contexts where wire swaps are available for free, a commonly used gate set is $\{c^0, c^1, c^2\}$, known as the NCT library (NOT, Controlled NOT, Toffoli gate), and corresponds to the rightmost generating set $c^2, c^1, Rc^1R, c^0, \sigma$ above (though we need two versions of c^1). The gate c^1 is usually included as it is cheaper than c^2 in physical implementations, but for the purpose of theory, it can simply be expressed in terms of c^0 and c^2 , and thus need not be included when s is included, giving the leftmost generating set c^2, s, c^0, σ .

When the wires are organized into a finite cycle, any universal set of gates for G gives one for the alternating group G_n .

Theorem 1 *Let $F \subset G$ be any finite generating set. Then for large enough n , $\{f_n \mid f \in F\}$ generates G_n .*

Proof It is enough to show that $\{f_n \mid f \in F\}$ generates some generating set of G_n . By Lemma 8, it is enough to generate all of S and the gates c_n^0, c_n^2 . Thus, it is enough to generate the standard generating set $\{s_n, c_n^0, c_n^2, \sigma_n\}$.

The gate s can be written as a finite composition of elements in F , i.e., $s = f^1 \circ \dots \circ f^k$ for some $f^1, \dots, f^k \in F$. By Lemma 5, for n large enough, we then have $s_n = f_n^1 \circ \dots \circ f_n^k$. The same argument applies to the gates in $\{c^0, c^2, \sigma\}$, so for large enough n , we indeed have

$$\{s_n, c_n^0, c_n^2, \sigma_n\} \subset \langle f_n \mid f \in F \rangle,$$

concluding the proof. □

The crucial point in the proof was that G_n has essentially a single finite generating set that does not depend on n , so it is enough to write the formulas generating this generating set, and these formulas then work for large n . The converse of Theorem 1 is not true, i.e., the fact $\{f_n \mid f \in F\}$ and the shift together generate G_n for all large enough n does not imply that F and σ generate G :

Proposition 1 *For all $n \geq 4$, $G_n = \langle c_n^0, c_n^1, c_n^2 \rangle$.*

Proof Write $C^i = \sigma_n^{-i} \circ c_n^1 \circ \sigma_n^i$, and observe that

$$\begin{aligned} &c^1 \circ (\sigma^{-1} \circ (R \circ c^1 \circ R) \circ \sigma) \circ c^1 \\ &= C^{n-2} \circ \dots \circ C^1 \circ \dots \circ C^{n-1} \circ \dots \circ C^1 \circ \dots \circ C^{n-1}. \end{aligned}$$

By (2), we obtain the generating set in Lemma 8. □

Note that $\{c^0, c^1, c^2\} \subset G_L$, so these gates and the shift do not generate G , we are only able to generate the standard generating set for all n by sending information around the cycle.

7 Generating G by word-swapping

For words $u, v \in \{0, 1\}^n$, write $f_{u,v}$ for the map $f_{u,v}(x)_{[0,n-1]} = v$ if $x_{[0,n-1]} = u$, $f_{u,v}(x)_{[0,n-1]} = u$ if $x_{[0,n-1]} = v$, $f_{u,v}(x)_{[0,n-1]} = x_{[0,n-1]}$ otherwise, and $f_{u,v}(x)_i = x_i$ for $i \notin [0, n - 1]$. For example, the Toffoli gate is $c^2 = f_{011,111}$.

We give a classification of the pairs of words u, v such that $f_{u,v}$ is a universal gate together with the shift and the flip, i.e., $\langle f_{u,v}, \sigma, c^0 \rangle = G$. The characterization is that the words differ in exactly one place, which is not at the border.

Theorem 2 *Let $u', v' \in \{0, 1\}^n$ be distinct words and let $G' = \langle c^0, f_{u',v'}, \sigma \rangle$. Then, $G = G'$ if and only if $D(u', v') \in 0^*0100^*$.*

Proof For sufficiency, suppose the RHS holds. Since $c^0, \sigma \in G'$, we can conjugate u' to $u = 0^n$ and v' to $v = D(u', v')$. By applying the gate $f_{0^n,v}$, flipping the i th bit on the tape, applying the gate again, and again flipping the i th bit, we effectively eliminate one of the bits that $f_{0^n,v}$ looks at. In symbols,

$$\sigma c^0 f_{0^n,0v} c^0 f_{0^n,0v} \sigma^{-1} = f_{0^{n-1},v}$$

and

$$\sigma^{-n} c^0 \sigma^n f_{0^n,v0} \sigma^{-n} c^0 \sigma^n f_{0^n,v0} = f_{0^{n-1},v}.$$

By the assumption, v contains the subwords 01 and 10, so c^1 and Rc^1R are in G' by applying the formulas of the previous paragraph, eliminating all but the subword 01 (resp. 10) of v . Then by (2), also $s \in G'$.

Now, use the same trick to eliminate all but the subword 010 from v to obtain $f_{000,010} \in G'$. This can be conjugated to the Toffoli gate c^2 by applying suitable shifts of the swap and the flip. It then follows from Lemma 9 that $G' = G$.

As for necessity, if $D(u', v') \in 0^*$, then $f_{u',v'}$ is the trivial gate. If $D(u', v') \in 0^*1$, then $\{\sigma, c^0, f_{u',v'}\} \subset G_R$. If $D(u', v') \in 10^*$, then $\{\sigma, c^0, f_{u',v'}\} \subset G_L$.

If $D(u', v')$ contains more than one 1, then consider $w = {}^\omega 0.D(u', v')0^\omega$ as a vector in X_0 . Together with its shifts, it generates a shift-invariant vector space $V \leq X_0$. It does not generate all of X_0 : one way to see this is that the support of a sum of distinct

shifts of w has cardinality at least 2, since the leftmost coordinate of the leftmost summand and the rightmost coordinate of the rightmost summand are not cancelled. It is straightforward to show that $\{c^0, f_{u',v'}, \sigma\} \subset G_V$. □

In particular, the previous theorem shows that c^0, c^2, σ do not generate G (as we also observed directly after Proposition 1), since the Toffoli gate $f_{011,111}$ satisfies $D(011, 111) = 100 \notin 0^*0100^*$.

One cannot omit c^0 from the theorem, since we can never have $G = \langle f_{u,v}, \sigma \rangle$ for the simple reason that either $0^{\mathbb{Z}}$ or $1^{\mathbb{Z}}$ is a fixed point for both generators when $D(u, v) \in 0^*0100^*$, and when $D(u, v) \notin 0^*0100^*$, even $\{f_{u,v}, c^0, \sigma\}$ does not generate G by the theorem.

8 Asynchronous application of elementary CA

We now exhibit a very simple inert element that generates G together with the shift, i.e., a *singleton universal gate set*. This is an optimal way to generate G in several ways: we need only one gate, which only modifies one cell at a time, and which has strong radius $r = 1$. This radius is optimal since all permutations of two or fewer bits are affine.

We recall the *Wolfram number* [12] of an elementary cellular automaton. Enumerate words $\{0, 1\}^3$ in reverse lexicographic order as $w_0 = 111, w_1 = 110, w_2 = 101, \dots$. To each $b \in \{0, 1\}^8$, we associate a self-homeomorphism e^b of X defined by $e^b(x)_0 = b_j$ where $w_j = x_{[-1,1]}$, and $e^b(x)_i = x_i$ for $i \neq 0$. This number b is traditionally written in decimal to obtain the Wolfram number. We show that the singleton universal gate sets among the e^n are precisely $e^{57} = e^{00111001_2}$ and $e^{99} = e^{01100011_2}$.

Usually, a cellular automaton applies the local rule to all cells of \mathbb{Z} at once. The definition here is the definition of (fully) asynchronous application of the cellular automaton, as studied in [7,13].

The map e^{57} can be described as follows: $e^{57}(x)_0 = 1 - x_0$ unless $x_{-1} = 0$ and $x_1 = 1$, and no other cell is changed. Thus, $e^{57} = c^0 \circ \sigma \circ f_{001,011} \circ \sigma^{-1}$. The map e^{99} is its mirror image.

Theorem 3 *The function e^n is a universal gate in G if and only if $n \in \{57, 99\}$.*

In particular, $G = \langle e^{57}, \sigma \rangle = \langle e^{99}, \sigma \rangle$ is 2-generated.

Proof Consider general $e^n, n \in \{0, 1, \dots, 255\}$. For e^n to be in G , the binary representation of n must be of the form

$$n = (ab(1 - a)(1 - b)cd(1 - c)(1 - d))_2.$$

If $d = 1$, then $0^{\mathbb{Z}}$ is fixed, so we must have $d = 0$. Similarly we must have $a = 0$, so for universality we must have $n = (0b1(1 - b)c0(1 - c)1)_2$, and the remaining choices are among $00110011, 00111001, 01100011, 01101001$. These are $e^{51}, e^{57}, e^{99}, e^{105}$, respectively. We have $e^{51} = c^0$, so it is not a singleton universal gate set by Lemma 6. The gate e^{105} sums the bits at x_{-1}, x_1 to the bit at x_0 , and then flips the value at x_0 .

n . Without some additional argument, universal gate sets only form a Σ_2^0 set in this context.

Note also that e_{57} is the composition of a word swap and the flip. We do not have a characterization of word swaps whose composition with the flip is a singleton universal gate set. A related question is whether a random inert gate in $\text{Sym}(\{0, 1\}^n)$, together with the shift, generates G with high probability (as $n \rightarrow \infty$).

We asked a related question also in [10], namely whether such a gate could even be universal as a gate on $[0, n - 1]$, when we are not allowed to apply it over the borders. We do not know any gates that are universal in this stronger sense; the gate e^{57} is not universal in this sense.

We do not know whether G is finitely presented. A presentation for the groupoid of reversible gates is also not known [11].

In [1] the analog of Post's lattice for all reversible bit operations is fully described. Since they allow ancilla bits (bits containing some prescribed value, and which need not return to their initial value in the end), it is not immediately clear that this classification implies something in our framework. A full description of the lattice of subgroups of G containing the shift and the swap would amount to classifying the lattice of reversible transformations implementable using *borrowed bits*, in the terminology of [14]; we do not know if there is a known classification. We have not checked whether the varied Toffoli gate of [14] is a universal singleton gate set in our sense.

We note that, unlike the lattice of [1] and Post's lattice, the lattice of subgroups of G containing the shift (but not necessarily the swap) is uncountable—it is easy to embed, for example, the infinite direct sum $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}_2$, which has uncountably many subgroups.

There are obvious generalizations of G which seem interesting. It is a standard direction of generalization in the theory of reversible gates to change the binary alphabet to a higher-arity one, and sometimes the qualitative properties change, e.g., the parity of the finitely generated part of the group of reversible gates depends on the parity of the alphabet [3, 4, 11]. Here, the fact we have a geometry for the arrangement of the wires allows us to do much more: we can change the geometry to an arbitrary group, and instead of just increasing the size of the alphabet, we can replace the set of legal configurations by a subshift. This leads to interesting issues, which we plan to study in a follow-up paper.

Acknowledgements We thank Ilkka Törmä for useful suggestions.

Funding Open access funding provided by University of Turku (UTU) including Turku University Central Hospital.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Straight-line grammar for the standard generators

We give a context-free grammar, obtained directly from the proof, that describes the standard generators for the swap s , the flip c^0 , the CNOTs c^1 , Rc^1R , and the Toffoli gate c^2 , more precisely the grammar generates a single string over the alphabet $\{1, 2, \dots, 6\}$ whose letters indicate the cell of $[0, n]$ ($n \geq 7$) where the e^{57} gate should be applied. The grammar generates the standard gates in cell $4 \in [0, n]$ when the start symbol is varied.

In the grammar we use an extension of the NCT naming scheme: N_i applies c^0 at i (i.e., it applies $\sigma^{-i} \circ c^0 \circ \sigma^i$), C_i corresponds to $\sigma^{-i} \circ c^1 \circ \sigma^i$ and T_i corresponds to $\sigma^{-i} \circ c^2 \circ \sigma^i$. The swap of cells i and $i + 1$ is S_i , and $\sigma^{-i} \circ R \circ c^1 \circ R \circ \sigma^i$ is D_i . The terminal symbol $i \in \{1, \dots, 6\}$ is interpreted as $\sigma^{-i} \circ e^{57} \circ \sigma^i$.

The grammar is $G = (V, \Sigma, R, S')$, where the start symbol is the element of $\{N_3, C_3, T_3, D_3, S_3\}$ we want to generate,

$$V = \{N_3, C_3, T_3, D_3, S_3\} \cup \{N_2, N_4, N_5, D_4, \bar{E}_3, \bar{E}_4\},$$

$$\Sigma = \{1, 2, 3, 4, 5, 6\},$$

and R contains the following rewrite rules:

$$T_3 \rightarrow S_3 N_3 \bar{E}_4 N_3 S_3$$

$$S_3 \rightarrow C_3 D_4 C_3$$

$$C_3 \rightarrow \bar{E}_3 N_2 \bar{E}_3 N_2$$

$$D_3 \rightarrow N_2 \bar{E}_3 N_4 \bar{E}_3 N_2 N_4$$

$$D_4 \rightarrow N_3 \bar{E}_4 N_5 \bar{E}_4 N_3 N_5$$

$$\bar{E}_3 \rightarrow N_3 3$$

$$\bar{E}_4 \rightarrow N_4 4$$

$$N_2 \rightarrow 12312321212132121212323121321232323231232321213232$$

$$N_3 \rightarrow 23423432323243232323434232432343434342343432324343$$

$$N_4 \rightarrow 345345434343543434345453435434545454545454543435454$$

$$N_5 \rightarrow 45645654545465454545656454654565656564565654546565$$

The standard gates

We list the strings generated by the straight-line grammar from Sect. 1 from start symbols N_3, C_3, T_3, D_3, S_3 , i.e., implementations of standard gates as shifts of e^{57} .

$$N_3 \rightarrow 23423432323243232323434232432343434342343432324343$$

$$C_3 \rightarrow 123123212121321212123231213212323232312323212132323234$$

$$2343232324323232343423243234343423434323243431231232$$

$$121213212121232312132123232323123232121323232342343232$$

$$32432323234342324323434342343432324343$$

$$T_3 \rightarrow 12312321212132121212323121321232323231232321213232323423$$

4323232432323234342324323434342343432324343123123212121
321212123231213212323232312323212132323234234323232432323
234342324323434343423434323243434564565454565454565645
465456565656456565454656523423432323243232323434232432343
4343423434323243434345345434343543434345453435434545453
454543435454456456545454654545456564546545656565645656545
465654345345434343543434345453435434545454534545434354542
34234323232432323234342324323434342343432324343123123212
1213212121232312132123232323123232121323232342343232324323
2323434232432343434342343432324343123123212121321212123231
2132123232323123232121323232342343232324323232343423243234
34343423434323243432342343232324323232343423243234343423
4343232434343453454343435434343454534354345454545345454343
545423423432324323232343423243234343423434323243431231
2321212132121212323121321232323231232321213232323423432323
2432323234342324323434343423434323243431231232121213212121
2323121321232323231232321213232323423432323243232323434232
4323434343423434323243434564565454546545454565645465456565
6564565654546565234234323232432323234342324323234343423434
323243434345345434343543434345453435434545454534545434343545
4456456545454654545456564546545656564565654546565434354345
4343435434343454534354345454545454343545423423432323243
2323234342324323434343423434323243431231232121213212121232
3121321232323231232321213232323423432323243232323434232432
3434343423434323243431231232121213212121232312132123232323
123232121323232434323232432323234342324323434342343432
324343

$D_3 \rightarrow$ 3453454343435434343454534354345454545345454343545412312321
2121321212123231213212323232312323212132323234234323232432
323234342324323434343423434323243433453454343435434343434545
3435434545454534545434354543234234323232432323234342324323
4343434234343232434312312321212132121212323121321232323231
232321213232

$S_3 \rightarrow$ 1231232121213212121232312132123232323123232121323232342343
2323243232323434232432343434342343432324343123123212121321
2121232312132123232323123232121323232342343232324323232343
4232432343434342343432324343456456545454654545456564546545
65656564565654546565234234323232432323234342324323434342
34343232434343453454343435434343454534354345454534545434
35454456456545454654545456564546545656564565654546565434
5345434343543434345453435434545454534545434354542342343232
32432323234342324323434342343432324343123123212121321212
1232312132123232323123232121323232342343232324323232343423
2432343434342343432324343123123212121321212123231213212323
23231232321213232323423432323243232323434232432343434234
3432324343

Checking the identity in Theorem 3

```

def a(l):
    if l[0] == 0 and l[2] == 1:
        return l
    return [l[0], 1-l[1], l[2], l[3], l[4]]

def b(l):
    if l[1] == 0 and l[3] == 1:
        return l
    return [l[0], l[1], 1-l[2], l[3], l[4]]

def c(l):
    if l[2] == 0 and l[4] == 1:
        return l
    return [l[0], l[1], l[2], 1-l[3], l[4]]

def center_flipped(l, s):
    orl = l
    for i in s:
        if i == "a":
            l = a(l)
        if i == "b":
            l = b(l)
        if i == "c":
            l = c(l)
    center_is_flipped = orl[2] != l[2]
    others_are_not = orl[:2] + orl[3:] == l[:2] + l[3:]
    return center_is_flipped and others_are_not

def binw(n):
    if n == 0:
        yield []
    else:
        for i in [0,1]:
            for b in binw(n-1):
                yield [i] + b

s = "abcabcbababacbabababcbcabacbabcbcbcbcbcbababcbcb"

for l in binw(5):
    if not center_flipped(l, reversed(s)):
        print ("It is not the flip!")
        break
else:
    print ("It is the flip.")

```

References

1. Aaronson, S., Grier, D., Schaeffer, L.: The classification of reversible bit operations. In 8th Innovations in Theoretical Computer Science Conference, vol. 67. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern (2017)
2. Barbieri, S., Kari, J., Salo, V.: The Group of Reversible Turing Machines, pp. 49–62. Springer International Publishing, Cham (2016)
3. Boykett, T.: Closed systems of invertible maps. *J. Mult.-Valued Logic Soft Comput.* **32**(5-6), 565–605
4. Boykett, T., Kari, J., Salo, V.: Strongly Universal Reversible Gate Sets, pp. 239–254. Springer International Publishing, Cham (2016)
5. Ceccherini-Silberstein, T., Coornaert, M.: Cellular Automata. In: Cellular Automata and Groups. Springer Monographs in Mathematics. Springer, Berlin, Heidelberg (2010)
6. Kim, K.H., Roush, F.W.: Solution of two conjectures in symbolic dynamics. *Proc. Am. Math. Soc.* **112**(4), 1163–1168 (1991)
7. Macauley, M., Jon, M., Mortveit, H.S.: Dynamics groups of asynchronous cellular automata. *J. Algebraic Comb.* **33**(1), 11–35 (2011)
8. Robinson, D.: A Course in the Theory of Groups. Graduate Texts in Mathematics. Springer, New York (1996)
9. Salo, V.: Transitive action on finite points of a full shift and a finitary Ryan’s theorem. *Erg. Theory Dyn. Syst.*, pp. 1–31, (2017)
10. Salo, V.: Universal groups of cellular automata. ArXiv e-prints, August 2018. Available at <https://arxiv.org/abs/1808.08697>
11. Selinger, P.: Reversible k-valued logic circuits are finitely generated for odd k. ArXiv e-prints, April 2016. Available at <https://arxiv.org/abs/1604.01646>
12. Stephen, W.: Statistical mechanics of cellular automata. *Rev. Mod. Phys.* **55**(3), 601–644 (1983)
13. Vielhaber, M.: Computing by temporal order: Asynchronous cellular automata. In Formenti, E. (ed.) Proceedings 18th International Workshop on Cellular Automata and Discrete Complex Systems and 3rd International Symposium Journées Automates Cellulaires (AUTOMATA&JAC 2012), La Marana, Corsica, vol. 90, pp. 166–176 (2012)
14. Xu, S.: Reversible Logic Synthesis with Minimal Usage of Ancilla Bits. ArXiv e-prints, June 2015. Available at <https://arxiv.org/abs/1506.03777>
15. Yves, L.: Towards an algebraic theory of boolean circuits. *J. Pure Appl. Algebra* **184**, 2003 (2003)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.