Correction to: Assessment of Autism Spectrum Disorder in Deaf Adults with Intellectual Disability: Feasibility and Psychometric Properties of an Adapted Version of the Autism Diagnostic Observation Schedule (ADOS-2)

D. Holzinger^{1,2,3} · C. Weber^{1,4} · S. Bölte^{5,6,7} · J. Fellingner^{1,2,8} · J. Hofer^{1,2,9}

Published online: 17 August 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to:
Journal of Autism and Developmental Disorders
<https://doi.org/10.1007/s10803-021-05203-5>

Correction regarding Table 1 and the corresponding text passage under “Participants and Study Setting”. Corrections are highlighted bold.

Participants and Study Setting (3rd paragraph)

The sample had a mean age of 44.6 years (range 17 to 75, SD = 18.7) and a majority was male (65%). The level of intellectual functioning varied. **While 50% had a cognitive level of functioning above an age of reference (AOR) of 6 years, in 50% the AOR was below 6 years.** The degree of hearing loss was....

The original article can be found online at <https://doi.org/10.1007/s10803-021-05203-5>.

✉ J. Hofer
johannes.hofer@bblinz.at

- ¹ Forschungsinstitut für Entwicklungsmedizin, Johannes Kepler Universität Linz, Linz, Austria
- ² Institut für Sinnes- und Sprachneurologie, Konventhospital Barmherzige Brüder, Seilerstätte 2, 4021 Linz, Austria
- ³ Institut für Sprachwissenschaft, Karl-Franzens-Universität Graz, Graz, Austria
- ⁴ Institut für Inklusive Pädagogik, Pädagogische Hochschule OÖ, Linz, Austria
- ⁵ Department of Women’s and Children’s Health, Center of Neurodevelopmental Disorders (KIND), Centre

for Psychiatry Research, Karolinska Institutet & Stockholm Health Care Services, Region Stockholm, Stockholm, Sweden

- ⁶ Child and Adolescent Psychiatry, Stockholm Health Care Services, Region Stockholm, Stockholm, Sweden
- ⁷ Curtin Autism Research Group, Curtin School of Allied Health, Curtin University, Perth, WA, Australia
- ⁸ Abteilung für Sozialpsychiatrie der Universitätsklinik für Psychiatrie und Psychotherapie, Medizinische Universität Wien, Vienna, Austria
- ⁹ Abteilung für Pädiatrie I, Medizinische Universität Innsbruck, Innsbruck, Austria

Table 1 Participant characteristics

	Modul 1 (n = 8)	Modul 2 (n = 16)	Modul 3 (n = 32)
Age M (SD)	41.25 (19.91)	37.00 (18.52)	49.00 (17.48)
Sex male n (%)	4 (50.0)	14 (87.5)	19 (59.4)
Cognitive functioning AOR ^a M (SD)	3.28 (1.58)	4.80 (1.39)	6.77 (1.73)
Cognitive functioning AOR^a n (%)			
≥ 9 years	0 (0.0)	1 (6.3)	4 (12.5)
< 9 to 6 years	1 (12.5)	1 (6.3)	21 (65.6)
< 6 to 3 years	4 (50.0)	14 (87.5)	7 (21.9)
< 3 years	3 (37.5)	0 (0.0)	0 (0.0)
Hearing loss n (%)			
Moderate	0 (0.0)	0 (0.0)	1 (3.1)
Severe	0 (0.0)	1 (6.3)	5 (15.6)
Profound	8 (100.0)	15 (93.8)	26 (81.3)
Cerebral Palsy n (%)	4 (50.0)	2 (14.3)	7 (21.9)
Epilepsy ^a n (%)	5 (62.5)	6 (42.9)	4 (12.5)
Psychiatric Diagnosis n (%)	1 (12.5)	3 (21.4)	10 (32.3)

^asignificant differences ($p < 0.05$) between modules (Non-parametric One-way Anova for continuous variables and exact Fisher tests for categorical variables)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.