



Correction to: Brief Report: Development and Validation of the Autism Spectrum Knowledge Scale General Population Version: Preliminary Analyses

Maryellen Brunson McClain¹ · Bryn Harris² · Sarah E. Schwartz¹ · Kandice J. Benallie¹ · Megan E. Golson¹ · Chandler M. Benney¹

Published online: 21 December 2021
© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to:
Journal of Autism and Developmental Disorders
(2019) 49:3007–3015
<https://doi.org/10.1007/s10803-019-04019-8>

Please note the following corrections to this article:

In Table 2 the correct response to Item 12, *Some individuals with autism spectrum disorder may be uncoordinated or clumsy*, should be True. The corrected Table 2 is shown below.

The original article has been revised.

The original article can be found online at <https://doi.org/10.1007/s10803-019-04019-8>.

✉ Maryellen Brunson McClain
maryellen.mcclainverdoes@usu.edu

¹ Utah State University, 2810 Old Main Hill, Logan, UT 84322, USA

² University of Colorado Denver, Denver, CO, USA

Table 2 ASKSG Items and Response Choices, with both the percentage of sample responding correctly and estimates of item difficulty from the Rasch Model

Item #	Item text	Correct response	Observed % correct	Item difficulty parameter (Eta)	
				Est.	SE
13	Many individuals with autism spectrum disorder have difficulties expressing themselves	True	88.68	−1.494	0.178
10	Individuals with autism spectrum disorder may have strict routines or rituals	True	87.42	−1.368	0.170
5	Autism spectrum disorder is caused by a lack of motherly warmth	False	85.85	−1.225	0.162
11	Individuals with autism spectrum disorder have difficulties interacting socially or others	True	83.96	−1.068	0.155
31	Many children with autism spectrum disorder are at risk for academic difficulties	True	83.65	−1.044	0.153
24	Social skills training is an effective treatment for some individuals with autism spectrum disorder	True	83.02	−0.995	0.151
9	Children with autism spectrum disorder may not play with toys the way they are intended	True	81.13	−0.858	0.146
26	Most individuals with autism spectrum disorder will never learn to speak	False	80.50	−0.815	0.144
17	If a teacher believes a student has autism spectrum disorder, they can make a diagnosis	False	79.56	−0.751	0.142
8	All individuals with autism spectrum disorder have low intellectual quotients (i.e., IQs)	False	78.30	−0.670	0.139
12	Some individuals with autism spectrum disorder may be uncoordinated or clumsy	True	75.16	−0.479	0.133
22	There are no beneficial treatments available for individuals with autism spectrum disorder	False	75.16	−0.479	0.133
28	Autism spectrum disorder only affects children	False	74.84	−0.460	0.133
27	Symptoms of autism spectrum disorder do not change throughout an individual's life	False	71.07	−0.252	0.128
2	Vaccines can cause autism spectrum disorder	False	70.44	−0.218	0.127
15	Diagnosis of autism spectrum disorder is primarily based on behavioral observations and parent interviews	True	67.92	−0.089	0.124
30	Up to 70% of individuals with autism spectrum disorder also have an additional mental health diagnosis (e.g., anxiety)	True	66.35	−0.010	0.123
29	Many individuals with autism spectrum disorder have difficulties living and working independently in adulthood	True	65.09	0.051	0.122
3	Boys are four times as likely than girls to have autism spectrum disorder	True	64.47	0.081	0.122
1	Less than 2% of people in the US have autism spectrum disorder	True	58.81	0.346	0.119
25	Intellectual quotient (i.e., IQ) and age affect treatment success for children with autism spectrum disorder	True	58.81	0.346	0.119
14	Symptoms of autism spectrum disorder do not appear before the age of 2 years	False	56.29	0.460	0.118
16	Autism spectrum disorder can only be diagnosed after the age of 4 years	False	54.09	0.559	0.117
23	Restricting certain foods (e.g., gluten) is an effective treatment for autism spectrum disorder.	False	48.11	0.824	0.177
20	It is possible for autism spectrum disorder to develop into adulthood	False	42.14	1.091	0.118
4	Children who have a brother or sister with autism spectrum disorder are more likely to develop the disorder	True	40.57	1.163	0.119
18	Autism spectrum disorder can be diagnosed with brain imaging	False	38.68	1.249	0.120
19	For a diagnosis of autism spectrum disorder, symptoms must be present from early childhood	True	38.68	1.249	0.120
6	Advanced paternal (father) age is a risk factor for autism spectrum disorder	True	38.36	1.264	0.120
7	There are no differences in the identification rates of autism spectrum disorder across racial and ethnic groups	False	36.16	1.367	0.121
21	A diagnosis of autism spectrum disorder can only be made by a medical doctor	False	20.44	2.222	0.142

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.