




Correction to: Does cleavage stage morphology increase the discriminatory power of prediction in blastocyst transfer outcome?

Zhenfang Liu¹ · Jiali Cai^{1,2} · Lanlan Liu^{1,2} · Ling Ouyang³ · Jinghua Chen¹ · Chao Yang¹ · Kaijie Chen¹ · Xiaolian Yang¹ · Jianzhi Ren¹ · Xiaoming Jiang^{1,2} 

Published online: 18 December 2023
© Springer Science+Business Media, LLC, part of Springer Nature 2023

Correction to: Journal of Assisted Reproduction and Genetics
<https://doi.org/10.1007/s10815-023-02997-4>

In this article the Figs. 1, 2, and 3 captions had been interchanged; the figure captions should have appeared as shown below.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10815-023-02997-4>.

✉ Xiaoming Jiang
keaishmily@126.com

¹ Reproductive Medicine Center, Xiamen University Affiliated Chenggong Hospital, Xiamen 361003, Fujian, China

² School of Medicine, Xiamen University, Xiamen 361005, Fujian, China

³ Medical Quality Management Department, Xiamen University Affiliated Chenggong Hospital, Xiamen 361003, Fujian, China




Fig. 1 The AUCs of ROC curves of the models. a The AUCs of ROC curves of XGboost, b the AUCs of ROC curves of LASSO, and c the AUCs of ROC curves of GLM. Model 1, all-in model with D3 mor-

phology. Model 2, all-in model without D3 morphology. Model 3, embryo quality only model with D3 morphology. Model 4, embryo quality only model without D3 morphology




Fig. 2 Calibration curves for predicting models




Fig. 3 The importance of each variable in different models. A: XGboost; B: LASSO; C: GLM