Correction to: On a superalgebraically converging, numerically stable solving strategy for electromagnetic scattering by impedance cylinders

Emrah Sever¹ · Yury A. Tuchkin¹ · Fatih Dikmen¹

Published online: 31 January 2022

© Springer Science+Business Media, LLC, part of Springer Nature 2022

Correction to: J Comput Electron (2018) 17:427–435
<https://doi.org/10.1007/s10825-017-1073-9>

The original version of this article unfortunately contained a typographical error, and it has been corrected with this erratum. The first sub-section “Posing of the boundary value problem” intends to proceed with the introduction of the following integro-differential operators **R** and **D**

$$R(\xi(q)) = \int_{\Gamma} \xi(p) \frac{\partial \mathcal{G}_2(q, p)}{\partial n'} dl',$$

$$D(\xi(q)) = \int_{\Gamma} \xi(p) \frac{\partial^2 \mathcal{G}_2(q, p)}{\partial n \partial n'} dl'$$

in addition to the correctly presented **S** and **V**.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10825-017-1073-9>.

✉ Fatih Dikmen
dikmen@gtu.edu.tr

Emrah Sever
emrahsever@gtu.edu.tr

Yury A. Tuchkin
yury.tu@gmail.com

¹ Department of Electronics Engineering, Gebze Technical University, Kocaeli 41400, Turkey