



# Household Social Needs, Emotional Functioning, and Stress in Low-Income Latinx Children and their Mothers

Victoria F. Keeton<sup>1,2</sup> · Janice F. Bell<sup>2</sup> · Christiana Drake<sup>2,3</sup> · Erik O. Fernandez y. Garcia<sup>2,4</sup> · Matthew Pantell<sup>5</sup> · Danielle Hessler<sup>6</sup> · Holly Wing<sup>7</sup> · Patricia P. Silveira<sup>8,9</sup> · Kieran J. O' Donnell<sup>8,9,10</sup> · Euclides José de Mendonça Filho<sup>8,9</sup> · Michael J. Meaney<sup>8,9,11</sup> · Laura M. Gottlieb<sup>6</sup>

Accepted: 3 January 2023 / Published online: 19 January 2023  
© The Author(s) 2023

## Abstract

Latinx families may be particularly vulnerable to emotional dysfunction, due to higher rates of economic hardship and complex social influences in this population. Little is known about the impact of environmental stressors such as unmet social needs and maternal stress on the emotional health of Latinx children from low-income families. We conducted secondary analyses using survey and biomarker data from 432 Latinx children and mothers collected in a separate study. We used binomial and multinomial logistic regression to test if household social needs, or maternal perceived stress or hair cortisol concentration (HCC), predicted child measures of emotional functioning or child HCC, independent of relevant sociodemographic factors. Approximately 40% of children in the sample had symptoms consistent with emotional dysfunction, and over 37% of households reported five or more social needs. High perceived maternal stress predicted higher odds of child emotional dysfunction (OR = 2.15; 95% CI [1.14, 4.04];  $p = 0.01$ ), and high maternal HCC was positively associated with high child HCC (OR = 10.60; 95% CI [4.20, 26.74];  $p < 0.01$ ). Most individual household social needs, as well as the level of household social need, were not independently associated with child emotional dysfunction or child HCC. Our findings begin to define a framework for understanding emotional health, stress, and resilience when caring for Latinx children and mothers living with high levels of social need, and the need for integrated mental health and social needs screening and interventions in settings that serve this population.

**Keywords** Child emotional health · Maternal stress · Social needs · Hair cortisol · Latinx

✉ Victoria F. Keeton  
victoria.keeton@ucsf.edu

<sup>1</sup> University of California, San Francisco, Department of Obstetrics, Gynecology, & Reproductive Sciences, Box 2930, San Francisco, CA 94143, USA

<sup>2</sup> University of California, Davis, Betty Irene Moore School of Nursing, 2450 48th St., Sacramento, CA 95817, USA

<sup>3</sup> University of California, Davis, Department of Statistics, 4101 Mathematical Sciences Bldg., Davis, CA 95616, USA

<sup>4</sup> University of California, Davis, Department of Pediatrics, 2521 Stockton Blvd, Suite 2200, Sacramento, CA 95817, USA

<sup>5</sup> University of California, San Francisco, Department of Pediatrics, 3333 California Street, Box 0848, San Francisco, CA 94143, USA

<sup>6</sup> University of California, San Francisco, Department of Family and Community Medicine, 500 Parnassus Ave, Box 0900, San Francisco, CA 94143, USA

<sup>7</sup> University of California, San Francisco, Center for Health and Community, 3333 California St., Box 0844, San Francisco, CA 94143, USA

<sup>8</sup> Douglas Mental Health University Institute, Douglas Research Center, McGill University, 6875 Boulevard LaSalle, Montreal, QC CA H4H1R3, Canada

<sup>9</sup> Ludmer Centre for Neuroinformatics and Mental Health and Department of Psychiatry, McGill University, Montreal, QC, Canada

<sup>10</sup> Yale Child Study Center & Department of Obstetrics, Gynecology & Reproductive Sciences, Yale School of Medicine, 230 South Frontage Rd., New Haven, CT 06519, USA

<sup>11</sup> Singapore Institute for Clinical Sciences, Agency for Science, Technology and Research (A\*STAR), 1 Fusionopolis Way, #20-10, Singapore 138632, Singapore

## Highlights

- Analysis of stress and household social needs in over 400 pairs of Latinx children and mothers.
- Unique measures, including 18 different social needs, and hair cortisol concentration as a measure of physiologic stress.
- Latinx households with low incomes show a high level of social need, with over one-third reporting having at least five needs.
- High maternal hair cortisol concentration is associated with greater risk for high child hair cortisol.
- Implications for the integration of mental health and social needs screening and intervention in the care of Latinx families.

Poor child emotional health can influence child development, interfere with daily functioning, and impact well-being into adulthood (Ghandour et al., 2019; Ogundele, 2018). Over the last two decades, the prevalence of emotional disorders such as anxiety and depression in children in the U.S. has steadily increased (Bethell et al., 2022; Bitsko et al., 2018; Ghandour et al., 2019), though it still may be underestimated secondary to well-studied barriers to accessing mental health diagnosis and treatment services in disadvantaged communities (Alegria et al., 2010; Ghandour et al., 2019; Marrast et al., 2016). At almost \$11 billion annually, the costs associated with child mental health conditions are higher than for any other child health disorder (Davis, 2014; Ghandour et al., 2019).

## Child Emotional Dysfunction and Unmet Social Needs

Given the known associations between poverty and economic disadvantage and multiple child physical health outcomes, it is not surprising that low household income is associated with higher rates of child emotional dysfunction (Council on Community Pediatrics et al., 2016; Ghandour et al., 2019; Harris & Santos, 2020; Ogundele, 2018). Evidence suggests that associations between poverty and child health are mediated by poverty-related social risk factors and social needs (Beck et al., 2016; Bethell et al., 2022; Fierman et al., 2016; Rodems & Shaefer, 2020). *Social risk factors* include specific adverse social conditions associated with poor health (Alderwick & Gottlieb, 2019; Bethell et al., 2022). While some ambiguity still exists in identifying the most salient social risk factors for child health, multiple health professionals and policy organizations have delineated and recommended screening for common domains of risk that include food, housing, employment, transportation, financial strain, safety, and access to health care (Kreuter et al., 2021). The construct of *social needs* extends the concept of social risk to include individual preferences, emphasizing the patient's shared role in identifying and prioritizing social interventions based on their perception of their most pressing needs (Alderwick & Gottlieb, 2019; Kreuter et al., 2021). Studies have distinguished between the concepts of income poverty

and unmet social needs, highlighting that families may experience persistent social needs even if they only endure short-term or transient episodes of income poverty, and that unmet social needs may contribute more to stress than income poverty (Neckerman et al., 2016; Rodems & Shaefer, 2020; Zilanawala & Pilkauskas, 2012).

Over 50% of Latinx households report having any social needs, as compared to 34% of White, non-Latinx households, regardless of income poverty level (Karpman et al., 2018), yet the relationship between unmet social needs and health outcomes in this population is understudied. Global economic hardship is associated with an increased risk for depression in Latinx parents (Ayon et al., 2010; Harris & Santos, 2020), and food insecurity is independently associated with serious psychological distress in Latinx adult populations (Becerra et al., 2015). Mothers that experience economic hardship are at higher risk for both depression (Arroyo-Borrell et al., 2017; Ayon et al., 2010; Harris & Santos, 2020; Zilanawala & Pilkauskas, 2012) and parenting stress (Duran et al., 2018; Luecken et al., 2013), and maternal depression and stress have each been independently linked to child emotional dysfunction (Arroyo-Borrell et al., 2017; Duran et al., 2018; Larson et al., 2008). Poor maternal mental health may negatively impact child emotional health by exposing the child to, or limiting the parent's ability to protect the child from, chronic stress in the environment (Shonkoff & Garner, 2012). A strong and compelling body of evidence demonstrates that structural determinants are the root causes of maternal health inequities experienced by Latinx and BIPOC populations in the U.S., and supports the movement away from interventions that unjustly hold individuals rather than systems responsible for health outcomes (Crear-Perry et al., 2020). This research suggests that advancing upstream solutions to reduce or mitigate social needs is critical to support maternal and child mental health.

Despite this preliminary work, the research on maternal-child emotional health specific to unmet social needs in U.S. Latinx populations is sparse. The paucity of evidence may be due in part to barriers in participant recruitment and data collection unique to this group (Alegria et al., 2010; Hopwood et al., 2009). Much of the current data that exists on these topics is gathered from large survey-based population studies that fail to include sizeable numbers of Latinx families (Hopwood et al., 2009). Evidence on emotional

disorders in Latinx populations is additionally limited because many studies fail to account for cultural influences on survey item endorsement and social desirability bias – participants' tendency to provide responses that they consider to be more favorable or acceptable (Hopwood et al., 2009). The sensitive nature of social needs, particularly around the use of public benefits, also has raised concern for social needs underreporting (Meyer et al., (2015)).

## Biomarkers of Stress

The limitations of survey-based research related to emotional health and social needs in Latinx populations have spurred interest in physiologic measures of stress that can be used as proxies for emotional health. The rationale for using physiologic biomarkers of child emotional dysfunction is that chronic stress exposure activates certain inflammatory and hormonal processes at persistent and eventually harmful levels, and in some cases this chronic stress alters gene regulation or expression, or brain structure and function, which impacts emotional health (Bates et al., 2017; Ogundele, 2018; Shonkoff & Garner, 2012). Release of the glucocorticoid hormone, cortisol, produces diverse genomic, metabolic, and physiological changes in response to stress (Khoury et al., 2019) and is now widely used as a biomarker for stress in both adults and children (Bates et al., 2017; Khoury et al., 2019). Studies have found elevated cortisol levels to correlate with poverty (Evans & English, 2002; Luecken et al., 2013), anxiety or depression symptoms (Pervanidou et al., 2013), food insecurity (Ling et al., 2019), and reports of racial discrimination (Berger & Saranyai, 2015), among other outcomes. Some studies also have explored the concordance between cortisol levels of mothers and their children in response to stress, and the factors that may moderate this relationship (Braren et al., 2019; Bryson et al., 2020; Dauegaard et al., 2020; Doan et al., 2020; Hollenbach et al., 2019; Johnson et al., 2018; Ling et al., 2020; Ludmer Nofech-Mozes et al., 2020). However, little research on biomarkers of stress has focused on Latinx mothers and children.

Rates of child emotional dysfunction and stress in the U.S. are high. Reducing these rates will require recognizing the multifactorial etiology of these disorders and developing interventions across the many relevant domains of child and family well-being (Bethell et al., 2022). A small but growing body of literature suggests that unmet social needs contribute to child emotional dysfunction and stress. This work is particularly relevant to Latinx people, who make up over 18% of the U.S. population (Office of Minority Health, 2021) and experience a disproportionate share of both poor mental health outcomes (Macias Gil et al., 2020; Marrast et al., 2016; Paz & Massey, 2016) and social risk (Rodems & Shaefer, 2020), yet

have been underrepresented in the existing literature on this topic. A better understanding of associations between unmet social needs, stress biomarkers, and mental health outcomes in Latinx families can be used to inform meaningful clinical or policy interventions that address mental health disparities in this high risk and marginalized population.

## The Current Study

The primary aim of the study was to use survey and biomarker data from Latinx mothers and their children to examine associations between household social needs, maternal perceived and physiologic stress, and child: a) emotional functioning or b) physiologic stress. We hypothesized that high level of household social needs, maternal perceived stress score, and maternal hair cortisol concentration (HCC) would independently and collectively predict increased child emotional dysfunction and child HCC. Our secondary aim was to explore whether specific social needs were associated with child: 2a) emotional functioning or 2b) child HCC. To accomplish these aims, we leveraged data collected previously in a separate clinical trial (Gottlieb et al., 2020), and performed a secondary analysis of select survey and biomarker data from the enrollment phase of data collection.

## Methods

### Study Design, Setting, and Sample

This study is a secondary analysis of cross-sectional survey and biomarker data obtained from children and their caregivers during the enrollment phase of the Health Advocates Study II (HASII) clinical trial, conducted between July 2016 and June 2018 in the children's urgent care of a pediatric primary care center nested in a large county hospital in San Francisco, California. Patients aged 0 through 17 years old presenting for a visit with their caregiver were recruited for the trial by convenience sampling; the details of their enrollment and inclusion criteria were published previously (Gottlieb et al., 2020). The current study limited the sample to one child per family and to children whose mothers were the enrolled caregiver ( $n = 549$ ), self-reported "Hispanic or Latino" origin or descent ( $n = 455$ ), and for whom there was complete maternal hair cortisol data ( $n = 432$ ).

### Protection of Human Subjects

HASII was approved by University of California, San Francisco Institutional Review Board (IRB), and only participants who consented according to the IRB-approved

protocol were included in this study sample. These secondary analyses of deidentified data were deemed exempt from full review by the Committee for the Protection of Human Subjects at the University of California, Davis.

## Study Measures and Variables

The child was the unit of analysis; measures collected about the child's mother were included in the analytic models of child health outcomes. At enrollment in HASII, a trained bilingual research assistant administered a survey (in English or Spanish) which caregivers completed about themselves and their children, including questions designed to capture their demographic characteristics, social situation and needs, physical and mental health status, and experiences with adverse events and stress. Caregivers were given the option to respond using written surveys (via electronic tablet), or in the case of literacy concerns, provide verbal responses to the research assistant. The research assistant also obtained a hair sample of approximately 3 cm measured from the scalp of each child and mother with sufficient hair length. More details related to the data collection and analysis protocol for the original clinical trial are available and published elsewhere (Gottlieb et al., 2020).

## Outcome Variables

Two outcomes were examined: a) child emotional functioning and b) child HCC. We initially performed analyses with each outcome as a continuous variable, however, two issues emerged that made us question the appropriateness of the models. First, several models exhibited aspects of natural binning such that certain values occurred very frequently and values in between, rarely or never. Second, even if we ignored the binning issue, linearity of associations was also questionable in all models. We then converted each outcome to a categorical variable and developed categories based on published reference ranges for emotional functioning and even percentile distributions for the hair cortisol (as explained below). The overall results did not differ substantially from the linear models, and thus are presented here for the better fitting categorical analyses.

Child emotional functioning was measured based on caregiver survey report using the parent-proxy version of the Pediatric Quality of Life Measurement Model (PedsQL™) (Varni et al., 2007b). The PedsQL™ 4.0 generic core scales measure the perception of health-related quality of life for children and adolescents in the areas of physical, emotional, social, and cognitive or school functioning, using developmentally appropriate language to adjust questions for different age groups (Varni et al., 2001). The PedsQL™ generic core scales have been demonstrated to be reliable and valid in a number of studies

(Varni et al., 2007b; Varni et al., 2001), including those specifically focused on children with mental health conditions (Bastiaansen et al., 2004) and from Spanish-speaking families (Roizen et al., 2008; Seid & Varni, 2005). In those studies, internal consistency reliabilities exceeded the minimum reliability standard of 0.70 required for group comparisons; total scale scores approached or exceeded the recommended alpha criterion of 0.90; construct validity was demonstrated using a known groups approach (Varni et al., 2007b; Varni et al., 2001). Agreement between parent proxy-report and child self-report of the PedsQL™ generic core scales was also assessed and found to be good, especially among samples of healthy children (Varni et al., 2007a, 2007b). In the HASII study, mothers completed the parent-proxy version of the PedsQL™ generic core scales about their children, responding to the questions designed for their child's age group. In the emotional functioning core scale (PedsQL™-EF), parents are asked to assess whether particular symptoms (e.g., sadness, anger, worry) have been a problem for the child over the past month. Items are scored on a five-point Likert scale ranging from 0 (never a problem) to 4 (almost always a problem) and converted to a reverse 100-point scale (where 0 = 100 and 4 = 0) with higher scores indicating better functioning (Varni et al., 2001). We dichotomized responses as "high functioning" (score  $\geq 64$ , coded as the reference group) vs. "low functioning" (score  $< 64$ ), based on recommended PedsQL™-EF cutoff scores for risk of low child emotional functioning in a large, ethnically diverse (61% Latinx) population study of healthy children (Varni et al., 2003).

Child HCC was measured using an established liquid chromatography-mass spectrometry (LC-MS) protocol (Gao et al., 2016); the full assay protocol is available in a previous publication (Gottlieb et al., 2020). HCC results are given in units of picogram per milligram (pg/mg) (Noppe et al., 2014). The subsample with complete child HCC data ( $N = 312$ ) was used in all child HCC outcome analyses. While some studies have attempted to establish normal reference ranges for hair cortisol concentration in healthy children (de Kruijff et al., 2020; Noppe et al., 2014; Prado-Gasco et al., 2019), robust evidence in this area is still lacking, particularly in ethnically diverse samples. Although there has not been agreement on a specific reference range for HCC values, it is widely accepted in the literature that higher levels indicate prolonged activation of the hypothalamic-pituitary-adrenal (HPA) axis, a marker for a chronic physiologic stress response (Bates et al., 2017; Khoury et al., 2019). HCC values were divided into tertiles for analysis so that each contained a third of the sample, and categories were defined by the values reflected in each: "lowest tertile" (0.22–3.99 pg/mg; coded as the reference group), "middle tertile" (4.00–7.82 pg/mg) and "highest tertile" (7.84–1998.88 pg/mg).

## Independent variables

Three primary predictor variables were examined: (1) household social needs, (2) maternal perceived stress score, and (3) maternal HCC. Despite categorizing the outcome variables in our preliminary analyses, running the models with continuous predictors still demonstrated a lack of linearity of associations, even on the logit scale. We therefore converted all three predictor variables to categorical, using evenly divided categories, to achieve the best model fit.

Household social needs were measured in the original study by mothers' survey responses of "yes" or "no" to current concerns about 18 possible issues for either themselves or members of their household (no=reference, yes). The social needs survey was created by the original study investigators and has been described in an earlier publication (Gottlieb et al., 2020). Survey items had been adapted from social risk screening questionnaires used in earlier studies (Gottlieb et al., 2014; Gottlieb et al., 2016), and used validated domain-specific measures when available (Brcic et al., 2011; Johnson et al., 2009; Keller et al., 2008; Kleinman et al., 2007). Despite previous research involving the use of these questionnaire items, currently there is no gold standard for social risk screening instruments, and very little psychometric evidence to evaluate their validity and reliability (Henrikson et al., 2019).

Household social needs variables were constructed in two ways: (1) categorical level of household needs (primary aim), and (2) indication of an individual household need (secondary aim). Household level of social needs was assigned based on a summative count of needs and then divided into four categorical groups for analysis. The first group included respondents who reported no needs, and the remainder of the sample was divided evenly across three additional groups: "none" (0 needs=ref), "1-2 needs", "3-4 needs", and "5 or more needs". This approach using the number of needs reported has been used in other published studies (Bethell et al., 2022; Rodems & Shaefer, 2020). Options for individual social needs included: *financial* (problems paying utility or medical bills, denied other income support, no health insurance), *housing* (difficulty finding housing, or habitability concerns), *food* (running out of food before having money to buy more), *employment* (difficulty finding, or problems with, a job, disability interfering with work, difficulty obtaining unemployment benefits), *transportation* (difficulty affording transportation or disability paratransit), *legal* (deportation, child support, family law issues), and *other* (no primary care provider, difficulty finding childcare or after-school activities, bullying, or household mental health concerns).

Maternal perceived stress was measured with the Perceived Stress Scale-4 (PSS-4), a shortened version of the full instrument (Baik et al., 2017) that measures the degree to which adults perceive stress in their control over or ability to handle events in their lives over the past month

(Warttig et al., 2013). The four items are scored on a five-point Likert scale ranging from 0 (never) to 4 (very often), where summative higher scores reflect higher perceived stress (Warttig et al., 2013). The PSS-4 tool is shown to be reliable and valid in studies of English-speaking (Warttig et al., 2013) and Spanish-speaking (Vallejo et al., 2018) participants. In these studies, internal consistency reliabilities for the instrument exceeded the minimum reliability standard of 0.70 for both English and Spanish, and alphas were adequate for both language groups (Vallejo et al., 2018; Warttig et al., 2013). Due to a lack of established cutoff scores, responses on the 20-point PSS-4 scale were divided evenly into tertiles – each containing a third of responses – and each tertile category was defined by the scores reflected in that group: "lowest tertile" (score: 0–4; coded as the reference group), "middle tertile" (score: 5–7), and "highest tertile" (score: 8 and higher).

Maternal physiologic stress was measured with HCC as described above for child HCC, and also divided categorically into tertiles and defined by the values in each: "lowest tertile" (0.22–3.99 pg/mg; coded as the reference group), "middle tertile" (4.00–7.82 pg/mg) and "highest tertile" (7.84–1998.88 pg/mg).

## Covariates

Sociodemographic and other physical and mental health characteristics theoretically associated with perceived and physiologic stress (Bates et al., 2017; Shonkoff & Garner, 2012), emotional health (Ogundele, 2018), or social needs (Kreuter et al., 2021) were included as covariates. In all models, sociodemographic covariates included mother's age (18–24 years=reference, 25–34 years, 35–44 years, 45 years and older), child's age divided by preschool, school age, and teen (0–5 years=reference, 6–12 years, 13–17 years), child's gender (male=reference, female), preferred language of the mother (English=reference, Spanish), mother's highest level of education (less than high school=reference, high school or general educational development [GED], some college or technical school, college/graduate degree) and annual household income (less than \$20,000=reference, \$20,000–50,000, more than \$50,000, don't know/decline to state).

Child health characteristics included health status, reported by the mother's responses on a 5-point scale ranging from poor to excellent to the statement, "In general, would you say your child's health is" (recoded into three groups as poor/fair=reference, good, very good/excellent); and child's physical functioning, rated on the PedsQL™ physical functioning 100-point scale and dichotomized as "high functioning" (score ≥64, coded as the reference group) vs. "low functioning" (score <64) according to evidence-based cutoffs for this tool (Vami et al., 2003). Maternal health characteristics included symptoms of depression, measured as a summative score on the Patient



Health Questionnaire-8 (PHQ-8) instrument and dichotomized according to evidence-based cutoffs (Kroenke et al., 2010) (score 0–5: “mild”=reference, score >5: “moderate/severe”); and physical health rating on the two-item Patient-Reported Outcomes Measurement Information System (PROMIS®) global physical health scale (Hays et al., 2017), measured by responses on a 5-point scale ranging from poor to excellent to the question, “In general, how would you rate your physical health?” (recoded into three groups as poor/fair=reference, good, very good/excellent).

## Data Analysis

All data were analyzed with Stata, version 16 statistical software (College Station, TX, USA). Descriptive statistics summarized the overall sociodemographic and health characteristics of the study sample and the dependent and independent variables.

We used multiple binomial logistic regression in our analyses of the first outcome to model child emotional functioning (high functioning=reference group) as a function of household social needs, maternal perceived stress, maternal HCC, and the sociodemographic and health covariates. In the first set of models, each social need was analyzed separately as an independent variable, controlling for maternal perceived stress, maternal HCC, and all covariates. In the second model, we used categorical level of social needs as the independent variable. Results from the binomial models are odds ratios (OR) comparing the independent variable categories to the reference group.

In our analyses of the second outcome, we used multinomial logistic regression to model child HCC (measured as lowest=reference group, middle, and highest HCC tertiles) as a function of household social needs, maternal perceived stress, maternal HCC, and the sociodemographic and health covariates. We ran the multinomial logistic regressions first using individual, and second using categorical level, to represent social needs. Due to the known potential associations between age or sex and HCC (Gray et al., 2018), we initially tested for effect modification by age or gender in each model, but did not find any meaningful significant interactions to support stratification by either variable in our final models. Results from the multinomial models are odds ratios (OR).

## Results

### Descriptive Findings

#### Sociodemographic and Health Characteristics

Table 1 summarizes frequencies for sociodemographic and health characteristics for the study sample. Most mothers

were between the ages of 25 and 44 years, and over half the children were less than 6 years old. The sample of children was relatively evenly divided between males and females. Approximately 86% of mothers in the sample reported Spanish as their preferred language, 57% reported less than a high school diploma as the highest level of education completed, and 53% of the sample reported an annual household income of less than \$20,000. Most mothers in the sample rated their child’s overall health as good or very good/excellent and reported their children had high physical functioning. Over one-third of mothers rated their own physical health as poor or fair, and almost half scored in the moderate to severe range for depression risk.

#### Household Social Needs

About 90% of mothers reported at least one current social need in their household, and well over one-third reported having more than five social needs. Descriptive statistics for reported social needs are presented in Table 2. Problems paying bills (39%), housing instability (38%), and food insecurity (36%) were the most reported social needs. Almost one-third of participant households reported job, habitability, transportation, or legal needs, or difficulty finding after-school activities for their children.

#### Maternal Stress

The median score on the PSS-4 was 6 (range: 0, 15; mean: 5.49, SD: 3.21). Median maternal HCC for the full sample was 5.47 pg/mg (range: 0.22, 1998.88; mean: 35.24, SD: 142.83). After verifying with our laboratory that the unusually high HCC values in the sample were not due to measurement or processing error, we also tested the models after removing very large observations – participants whose HCC was greater than 3 SD outside of the mean (Aguinis et al., 2013). Overall significance of findings was unchanged when these observations were removed so results presented here include the full sample.

## Regression Models

### Outcome a): Child Emotional Functioning

In the full sample, 41% of children scored at risk for emotional dysfunction on the PedsQL™-EF. In all fully adjusted binomial regression models (Table 3), the middle and highest tertiles of maternal perceived stress each positively predicted child emotional dysfunction compared to lowest tertile of maternal perceived stress (middle: OR range 1.92–2.73; highest: OR range 1.97–3.43;  $p < 0.05$ ; not all results shown in table). Neither the maternal HCC nor the level of social needs was significantly associated with

**Table 1** Sociodemographic and Health Characteristics ( $N = 432$ )

Covariates	<i>n</i> (%)	Dependent and Independent Variables	<i>n</i> (%)
Maternal Age in Years		Child PedsQL-EF Score	
18–24	55 (12.73)	High Functioning	256 (59.26)
25–34	179 (41.44)	Low Functioning (<64)	176 (40.74)
35–44	167 (38.66)	Child Hair Cortisol Concentration <sup>a</sup>	
45 or older	31 (7.18)	Lowest Tertile=ref	104 (33.33)
Age of Child in Years		Middle Tertile	104 (33.33)
0–5	252 (58.33)	Highest Tertile ( $\geq 37.39$ pg/mg)	104 (33.33)
6–12	126 (29.17)	Maternal PSS-4 Score	
13–17	54 (12.50)	Lowest Tertile	165 (38.19)
Gender of Child		Middle Tertile	131 (30.32)
Female	221 (51.16)	Highest Tertile ( $\geq 8$ )	136 (31.48)
Maternal Preferred Language		Maternal Hair Cortisol Concentration	
English	60 (13.89)	Lowest Tertile=ref	144 (33.33)
Spanish	372 (86.11)	Middle Tertile	144 (33.33)
Highest Education Level Completed		Highest Tertile ( $\geq 7.41$ pg/mg)	144 (33.33)
Less than High School	247 (57.18)		
High school/GED	121 (28.01)		
Technical school/Some College	40 (9.26)		
College Graduate/Graduate school	24 (5.56)		
Annual Household Income			
Less than \$20,000	231 (53.47)		
\$20,000–50,000	142 (32.87)		
More than \$50,000	10 (2.31)		
Don't know/decline to state	49 (11.34)		
Overall Rating of Child Health			
Poor/Fair	48 (11.11)		
Good	191 (44.21)		
Very Good/Excellent	193 (44.68)		
Child PQL Physical Functioning Score			
High Functioning	364 (84.26)		
Low Functioning (<64)	68 (15.74)		
Maternal PHQ-8 Depression Risk Score			
Mild	223 (51.62)		
Moderate/Severe (>5)	209 (48.38)		
Maternal PROMIS Physical Health Rating			
Poor/fair	161 (37.27)		
Good	219 (50.69)		
Very Good/Excellent	52 (12.04)		

<sup>a</sup>Due to missing child hair cortisol data, for this variable  $N = 312$

child emotional dysfunction in any of the models. For our secondary aim, almost none of the individual social needs were independently linked with child emotional dysfunction, except for finding after-school activities or opportunities (OR = 2.09; 95% CI [1.24, 3.52];  $p < 0.01$ ).

#### Outcome b): child HCC

Our analyses for this outcome included only the 312 children with complete HCC data. Chi-square analyses of the

differences in sociodemographic and health characteristics between children with and without child HCC revealed that the group with child HCC data included significantly more mothers who were over 34 years old (50% vs. 35%;  $p = 0.02$ ), more children who were female (57% vs. 35%;  $p < 0.01$ ), fewer children ages 0 to 5 years (49% vs. 80%;  $p < 0.01$ ), and more children with low physical functioning (18% vs. 8%;  $p < 0.01$ ), as compared to the group missing HCC data (results not shown in tables). Median child HCC was 18.69 (range: 0.71, 11,965.91; mean: 382.82, SD:

**Table 2** Current Household Social Needs ( $N = 432$ )<sup>a</sup>

	<i>n</i> (%)
<i>Financial</i>	
Problems paying bills, like electric, gas, water, or phone bills	169 (39.12)
Receiving medical or pharmacy bills you cannot afford	74 (17.13)
Getting cut off from or denied from programs that provide income support, like Cal Fresh (food stamps), CalWorks, etc	63 (14.58)
Having no health insurance	105 (24.31)
<i>Housing</i>	
Unstable housing including eviction, foreclosure, homelessness or staying with friends/family	168 (38.89)
Housing problems, like mold, insects rats or mice	128 (29.63)
<i>Food</i>	
Running out of food before having enough money or food stamps to buy more	156 (36.11)
<i>Employment</i>	
Difficulty finding a job	136 (31.48)
Problems with a current or former job, like unpaid wages, workers comp, discrimination or harassment	32 (7.41)
A disability interfering with the ability to work	48 (11.11)
Difficulty obtaining unemployment insurance	41 (9.49)
<i>Transportation</i>	
Difficulty affording transportation or ADA paratransit	136 (31.48)
<i>Legal</i>	
Other legal issues not mentioned above, including deportation concerns, child support or family law issues or violence	128 (29.63)
<i>Other</i>	
Having no primary care provider for your child or other household member ( $n = 367$ ) <sup>b</sup>	61 (16.62)
Difficulty finding after-school activities or opportunities for you or your child	122 (28.24)
Difficulty finding childcare ( $n = 367$ ) <sup>b</sup>	92 (25.07)
Bullying	48 (11.11)
Concerns about your or another adults' mental or behavioral health in your household	61 (14.12)
<i>Household Level of Social Needs</i>	
None	42 (9.72)
1–2	114 (26.39)
3–4	113 (26.16)
5 or more	163 (37.73)

<sup>a</sup>Reported needs are not mutually exclusive (participants reported up to 18 needs)

<sup>b</sup>This item was added after initial study initiation

1313.84). We again found no statistical differences in findings between models with and without the largest HCC values and thus included all participants in the subsample.

In all fully adjusted multinomial models, neither household level of social needs nor maternal perceived stress

score tertiles predicted middle or highest tertile of child HCC compared to lowest tertile of child HCC (Table 4; results not all shown in table). Being in the highest tertile of maternal HCC significantly predicted both middle (OR: 1.86–2.51;  $p < 0.05$ ) and highest tertiles of child HCC (OR: 9.80–11.00;  $p < 0.01$ ) in comparison to lowest tertile of child HCC across all models (results not all shown in table). For our secondary aim, individual social needs were not independently associated with middle or highest tertile of child HCC in any of the adjusted models.

## Discussion

To our knowledge, this is the first study in the U.S. to examine household social needs and maternal stress as predictors of child emotional dysfunction and child HCC in a sample of low-income and majority Spanish-speaking Latinx families. Over 40% of the children in the sample had emotional dysfunction, a staggering finding given the evidence that low income and Latinx youth are less likely to be diagnosed with and receive treatment for emotional disorders than those from higher income or non-Latinx households (Alegría et al., 2010; Ghandour et al., 2019; Marrast et al., 2016). Similar to previous findings in non-Latinx (Bryson et al., 2020; Daegaard et al., 2020; Doan et al., 2020; Schloß et al., 2019) and Latinx (Hollenbach et al., 2019) populations, maternal HCC was strongly associated with child HCC. This finding may indicate heritability of HCC and/or shared environmental risk factors.

The level of social needs in this sample was also striking. Approximately 90% of mothers reported having at least one social need, and over one-third reported having at least five needs. This far exceeds numbers for Latinx households reported in national survey research (Karpman et al., 2018). Our study is unique in that we asked about a large number of possible social needs, whereas much of the existing evidence in this area focuses on only a few domains of economic hardship (Bethell et al., 2022; Neckerman et al., 2016; Rodems & Shaefer, 2020; Zilanawala & Pilkauskas, 2012). Accordingly, we acknowledge that our broader assessment may explain the higher summative levels identified in our study, though there were other indicators that this is indeed a high needs sample. For instance, problems paying bills and housing instability were the most frequent concerns for our participants, consistent with a previous report that these are the most common social needs in Latinx families with children (Schmeer, 2012). Difficulty finding after-school activities for children was also a common concern and was the only individual need that significantly predicted higher odds of child emotional dysfunction.



**Table 3** Logistic Regression of Low Child Emotional Functioning on Household Social Needs, Maternal Perceived Stress, and Maternal Hair Cortisol ( $N = 432$ )

	Model 1 <sup>a</sup>			Model 2 <sup>b</sup>		
	OR	(95% CI)	<i>p</i>	OR	(95% CI)	<i>p</i>
<i>Household Social Needs (no=ref)</i>						
Problems paying bills	1.21	(0.74, 1.96)	0.43	---	---	---
Medical or pharmacy bills you can't afford	0.84	(0.45, 1.56)	0.59	---	---	---
Getting cut off from or denied income support	1.10	(0.56, 2.17)	0.76	---	---	---
Having no health insurance	0.96	(0.55, 1.67)	0.90	---	---	---
Unstable housing	0.80	(0.49, 1.30)	0.38	---	---	---
Housing problems (e.g., habitability)	1.29	(0.77, 2.16)	0.31	---	---	---
Running out of food	1.54	(0.94, 2.51)	0.08	---	---	---
Difficulty finding a job	1.08	(0.65, 1.79)	0.75	---	---	---
Problems with a current or former job	1.83	(0.76, 4.39)	0.17	---	---	---
Disability interfering with work	0.78	(0.35, 1.71)	0.54	---	---	---
Difficulty obtaining unemployment	1.95	(0.90, 4.24)	0.09	---	---	---
Difficulty affording transportation	1.29	(0.77, 2.17)	0.31	---	---	---
Other legal issues	1.21	(0.73, 2.02)	0.45	---	---	---
No PCP for household member ( $n = 367$ ) <sup>c</sup>	1.48	(0.75, 2.91)	0.25	---	---	---
Difficulty finding after-school activities	2.09	(1.24, 3.52)	<0.01*	---	---	---
Difficulty finding childcare ( $n = 367$ ) <sup>c</sup>	1.56	(0.85, 2.86)	0.14	---	---	---
Bullying	0.70	(0.32, 1.50)	0.36	---	---	---
Concerns about household adult mental health	1.04	(0.52, 2.08)	0.90	---	---	---
<i>Household Level of Social Need</i>						
None=ref	---	---	---	---	---	---
1–2 needs	---	---	---	0.95	(0.37, 2.39)	0.92
3–4 needs	---	---	---	0.90	(0.35, 2.28)	0.83
5 or more needs	---	---	---	1.49	(0.59, 3.72)	0.39
<i>Maternal PSS-4 Score</i>						
Lowest Tertile=ref	---	---	---	---	---	---
Middle Tertile	---	---	---	1.98	(1.07, 3.64)	0.02*
Highest Tertile	---	---	---	2.15	(1.14, 4.04)	0.01*
<i>Maternal Hair Cortisol</i>						
Lowest Tertile=ref	---	---	---	---	---	---
Middle Tertile	---	---	---	0.77	(0.42, 1.38)	0.38
Highest Tertile	---	---	---	0.96	(0.54, 1.69)	0.90

ref = reference group; All models included maternal PSS, maternal HCC and all covariates (results not all shown); \*Denotes significant findings at  $p < 0.05$

<sup>a</sup>Model 1 included individual needs as the social need variable; results reported are for 18 separate models of the outcome on each social need. Maternal PSS predicted emotional dysfunction across all above models (moderate: OR range 1.92–2.73; high: OR range 1.97–3.43;  $p < 0.05$ ; not all shown). Maternal HCC not significant in any model

<sup>b</sup>Model 2 included level of need as the social need variable

<sup>c</sup>This item added after original study began; analyses for this item includes participants with complete data only

While we did not have information about participants' immigration status, it is plausible that immigration influenced the study's urban, predominantly Spanish-speaking population. Threats of impending changes to policies affecting or affected by immigration status that occurred during the HASII study's enrollment period, such as those expected to the "public charge rule,"

negatively impacted enrollment in public assistance programs, including immigrant communities' enrollment in Medicaid (Bustamante et al., 2022; Miller et al., 2022; Wang et al., 2022). It is possible that fear associated with the public charge rule or other programs – e.g. programs that require documentation of immigration status or that depend on English fluency – may partially explain the

**Table 4** Multinomial Logistic Regression of Highest Tertile of Child Hair Cortisol on Household Social Needs, Maternal Perceived Stress, and Maternal Hair Cortisol ( $N = 312$ )

	Model 1 <sup>a</sup>			Model 2 <sup>b</sup>		
	OR	(95% CI)	<i>p</i>	OR	(95% CI)	<i>p</i>
<i>Household Social Needs (no=ref)</i>						
Problems paying bills	1.06	(0.49, 2.31)	0.86	---	---	---
Medical or pharmacy bills you can't afford	0.94	(0.36, 2.42)	0.90	---	---	---
Getting cut off from or denied income support	0.51	(0.16, 1.58)	0.24	---	---	---
Having no health insurance	0.69	(0.28, 1.67)	0.41	---	---	---
Unstable housing	1.42	(0.67, 3.02)	0.35	---	---	---
Housing problems (e.g., habitability)	0.95	(0.42, 2.16)	0.91	---	---	---
Running out of food	0.53	(0.24, 1.16)	0.11	---	---	---
Difficulty finding a job	1.42	(0.65, 3.09)	0.37	---	---	---
Problems with a current or former job	0.76	(0.19, 3.05)	0.70	---	---	---
Disability interfering with work	1.51	(0.47, 4.86)	0.48	---	---	---
Difficulty obtaining unemployment	0.69	(0.21, 2.31)	0.55	---	---	---
Difficulty affording transportation	0.87	(0.37, 2.01)	0.74	---	---	---
Other legal issues	0.65	(0.29, 1.44)	0.29	---	---	---
No PCP for household member (n = 254) <sup>c</sup>	0.42	(0.12, 1.38)	0.15	---	---	---
Difficulty finding after-school activities	1.27	(0.57, 2.81)	0.54	---	---	---
Difficulty finding childcare (n = 254) <sup>c</sup>	1.32	(0.52, 3.34)	0.55	---	---	---
Bullying	0.90	(0.25, 3.20)	0.87	---	---	---
Concerns about household adult mental health	0.95	(0.34, 2.66)	0.92	---	---	---
<i>Household Level of Social Need</i>						
None=ref	---	---	---	---	---	---
1–2 needs	---	---	---	0.52	(0.13, 2.07)	0.35
3–4 needs	---	---	---	1.08	(0.26, 4.39)	0.91
5 or more needs	---	---	---	0.44	(0.10, 1.78)	0.25
<i>Maternal PSS-4 Score</i>						
Lowest Tertile=ref	---	---	---	---	---	---
Middle Tertile	---	---	---	0.66	(0.25, 1.72)	0.40
Highest Tertile	---	---	---	0.60	(0.22, 1.67)	0.33
<i>Maternal Hair Cortisol</i>						
Lowest Tertile=ref	---	---	---	---	---	---
Middle Tertile	---	---	---	1.52	(0.60, 3.83)	0.36
Highest Tertile	---	---	---	10.60	(4.20, 26.74)	<0.01 <sup>*</sup>

ref = reference group; All models included maternal PSS, maternal HCC and all covariates (results not all shown); <sup>\*</sup>Denotes significant findings at  $p < 0.05$

<sup>a</sup>Model 1 included individual needs as the social need variable; results are for 18 separate models of the outcome on each need and shown for highest tertile of child HCC (compared to low). High maternal HCC predicted high child HCC across all models (OR range 9.80–11.00;  $p < 0.01$ ; not all shown). Maternal PSS not significant in any model

<sup>b</sup>Model 2 included level of need as the social need variable; results shown for highest tertile of child HCC (compared to low)

<sup>c</sup>This item added after original study began; analyses for this item includes participants with complete data only

high rates of social needs in our sample. Despite the recent reversal of the public charge rule, evidence suggests that its impact on immigrant communities' trust in public programs will persist (Bustamante et al., 2022), further underscoring the lasting harm of discriminatory policymaking.

Counter to our original hypotheses, level of social needs did not independently predict greater risk of parent-reported child emotional dysfunction. One theory that explains this lack of association is that a mother's awareness of household social needs does not necessarily correlate with her child's experience. For example, parents may struggle with

financial concerns such as paying bills or food insecurity without their child's knowledge, or families in households with food insecurity may prioritize feeding their children first (or instead) so that the child never goes hungry. Conversely, difficulty finding afterschool care or activities is a need that may be much more apparent to a child, which could explain why it was associated with greater emotional dysfunction in our sample. Alternatively, the child's emotional response to high household social needs may somehow be mitigated by other supportive factors known to exist in Latinx populations. The HASII study did not include measures of resilience for us to explore this concept further, but previous studies have documented characteristics of resilience in Latinx families facing different forms of social adversity (Linton et al., 2016; Perreira et al., 2019). "Familismo" is the concept of strong family ties and values that is central to most Latinx cultures (Ayon et al., 2010; Lawton et al., 2014), and is thought to serve as a buffer against many risk factors that can contribute to physical or mental health problems in this population (Ayon et al., 2010; Fillion et al., 2018; Potochnick & Perreira, 2010; Ruiz et al., 2018). For example, in multiple studies of Latinx families, strength of the parent-youth relationship has been shown to be a buffer between parental stress and child emotional problems (Ayon et al., 2010; Frasilheiro et al., 2016; Lorenzo-Blanco et al., 2017; Palermo et al., 2018; Perreira et al., 2019). It is also important to consider that the high overall level of needs and emotional dysfunction in our sample may have limited our ability to detect statistically significant associations between these two variables.

Level of social needs did not independently predict greater risk of being in the highest tertile of child HCC, again in contrast to our hypothesis that social needs would be positively associated with child stress. Our secondary exploration of individual needs also revealed no significant associations between specific social needs and child HCC. This is consistent with a recent study of non-Latinx infants and toddlers that found neither cumulative adversity nor multiple individual adversity indicators were significantly associated with HCC (Bryson et al., 2019). These findings contribute to an already conflicting research base on the associations between social needs and HCC in Latinx children. For example, one recent study showed higher child HCC was associated with greater food insecurity (Ling et al., 2019) while another showed no association between these variables (Distel et al., 2019). In addition to the theories related to child emotional experience discussed above, another theory that could explain the lack of association with child HCC is that prolonged stress exposure can result in a blunted cortisol response, although this has only been shown in studies examining HCC in non-Latinx populations (Dowd et al., 2009; Koumantarou Malisiova et al., 2020; Ouellette et al., 2015; Raffington et al., 2018;

Solarikova et al., 2020). According to this theory, a family's chronic stress from the inability to meet household social needs results in a dampening of the child's ability to mount a physiologic stress response. There is still much to learn about how biomarkers can contribute to our understanding of stress and emotional functioning, particularly in ethnically diverse pediatric populations (Stalder et al., 2017).

Mothers in this relatively large local Latinx sample seeking healthcare primarily preferred Spanish, and had low education and very low incomes, which is consistent with the demographic of Latinx populations living in poverty in the U.S. (Fontenot et al. (2018)). Language is often used as a measure of acculturation (Torres et al., 2012), indicating that our sample may have included a large percentage of immigrants with lower levels of acculturation. Our findings contradict previous evidence related to the "Hispanic health paradox", which proposes that less acculturated Latinx immigrants experience genetic, lifestyle, or cultural protective factors over their U.S. born or more acculturated Latinx counterparts that contribute to better health outcomes (Perreira et al., 2019; Ruiz et al., 2018; Teruya & Bazargan-Hejazi, 2013). The significant level of reported child emotional dysfunction and high HCC in mothers and children in our less acculturated sample could provide support for another body of literature that refutes the paradox theory, although we lack a substantial comparison group to test these assumptions. The paradox theory has been criticized for relying on methodology that favors healthier research participants and likely under-reports morbidity and mortality in immigrant populations (Ceballos & Palloni, 2010; Teruya & Bazargan-Hejazi, 2013); more research on this topic is warranted.

## Limitations

Our study findings should be interpreted with the consideration of five key limitations. First, our outcomes should be considered with acknowledgement of their limitations to generalizability. Convenience sampling was used in the original trial and sampling probabilities for all participants are unknown. Our study sample was restricted to low-income families seeking pediatric healthcare services in a single county hospital setting, as well as to families who were chosen for and consented to participate in a clinical trial and had time to complete study activities. Second, the cross-sectional study design of this baseline data analysis means we cannot infer the directionality of any statistical associations. Third, our findings should be considered in the context of limited research on social screening instrument validity. The screening questions used in the original study were developed de novo prior to the development of multi-domain social screening tools now being used more commonly in practice settings, though none of these has been

tested using gold standard psychometric and pragmatic validity testing procedures (Henrikson et al., 2019). Further, our assessment of child emotional functioning relied on parent proxy-report only and did not include self-report or observational data that could have reduced the potential for response bias. Fourth, in this study, 30% of children were missing hair samples because they were either not obtained or samples were insufficient for analysis. This may have resulted in an over-estimation of some of the associations we report, though many of the measured demographic and health characteristics between those included and excluded from this sub-analysis were similar. Finally, data may also be subject to unmeasured influence by other variables that are associated with child emotional functioning and stress (e.g. immigration status, acculturation stress, parenting style) but were not collected as part of the HASII study.

## Conclusion

This study contributes to a growing body of evidence linking maternal stress and social needs to child emotional dysfunction and stress. It is one of only a small number of published studies examining the physiologic impact of stress and social needs specifically in Latinx populations and adds new evidence on the use of HCC as a measure of stress in Latinx children and mothers. Our study findings have multiple implications for health policy, clinical practice, and research. We found that household social needs in low-income Latinx families with children are high, particularly related to financial and housing insecurity, calling attention to the need for policies that expand access to public assistance programs and remove barriers related to immigration status. Our findings provide further evidence that stress in Latinx mothers increases the risk for poor child emotional health (Arroyo-Borrell et al., 2017; Larson et al., 2008). Interventions that promote equitable access to conditions for optimal health – e.g., increased access to health insurance, mental health services, and respectful and culturally appropriate care (Crear-Perry et al., 2020) – are therefore critical to supporting Latinx mothers. The evaluation of maternal perceived stress may also be an important component of early identification and intervention related to child emotional health. Integrated caregiver and child behavioral health services in healthcare settings that serve low-income Latinx families may improve early diagnosis and intervention for youth at risk for stress and emotional dysfunction. Future research should explore the utility of HCC as a measure of stress in Latinx populations, as well as resilience and other supportive factors that may mitigate the impact of stress and social needs on emotional health in these vulnerable families.

## Data Availability

This paper uses data from a previous RCT. After publication, deidentified participant data is available by request to researchers whose proposed use of the data has been approved. Requests can be made to holly.wing@ucsf.edu, and approval is at the discretion of the primary RCT research team, with signed data use agreement.

## Code Availability

Code for data cleaning and analysis associated with current submission are available from the corresponding author on reasonable request.

**Author Contributions** Study concept/design; Acquisition, analysis, interpretation of data; Critical revision of the manuscript: all authors. Drafting of the manuscript: V.F.K., J.F.B., L.M.G., C.D., and F.y.G. Statistical analysis: V.F.K., J.F.B., C.D., M.P., D.H. Administrative, technical, or material support: H.W., L.M.G. Study supervision: V.F.K., J.F.B., L.M.G.

**Funding** This research is funded in part by the Gordon and Betty Moore Foundation through GBMF4294 to the University of California, Davis Betty Irene Moore School of Nursing. The primary RCT whose data we analyzed was financially supported by the Lisa and John Pritzker Family Fund, and the JPB Foundation of New York through a grant to the JPB Research Network on Toxic Stress: A Project of the Center on the Developing Child at Harvard University. V.F.K. is supported by a University of California, San Francisco, Preterm Birth Initiative transdisciplinary post-doctoral fellowship, funded by Marc and Lynne Benioff and a T32 training grant (1T32HD098057) from the National Institute of Child Health and Human Development entitled “Transdisciplinary Research Training to Reduce Disparities in Preterm Birth and Improve Maternal and Neonatal Outcomes”.

## Compliance with ethical standards

**Conflict of Interest** The authors declare no competing interests.

**Ethics Approval** This study was based on a secondary analysis of deidentified data and was therefore deemed exempt from full review by the Committee for the Protection of Human Subjects at the University of California, Davis.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright

holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Aguinis, H., Gottfredson, R. K., & Joo, H. (2013). Best-practice recommendations for defining, identifying, and handling outliers. *Organizational Research Methods*, *16*(2), 270–301. <https://doi.org/10.1177/1094428112470848>.
- Alderwick, H., & Gottlieb, L. M. (2019). Meanings and misunderstandings: A social determinants of health lexicon for health care systems. *Milbank Quarterly*, *97*(2), 407–419. <https://doi.org/10.1111/1468-0009.12390>.
- Alegria, M., Vallas, M., & Pumariega, A. J. (2010). Racial and ethnic disparities in pediatric mental health. *Child and Adolescent Psychiatric Clinics of North America*, *19*(4), 759–774. <https://doi.org/10.1016/j.chc.2010.07.001>.
- Arroyo-Borrell, E., Renart, G., Saurina, C., & Saez, M. (2017). Influence maternal background has on children's mental health. *International Journal for Equity in Health*, *16*(1), 63. <https://doi.org/10.1186/s12939-017-0559-1>.
- Ayon, C., Marsiglia, F. F., & Bermudez-Parsai, M. (2010). Latino family mental health: Exploring the role of discrimination and familismo. *Journal of Community Psychology*, *38*(6), 742–756. <https://doi.org/10.1002/jcop.20392>.
- Baik, S. H., Fox, R. S., Mills, S. D., Roesch, S. C., Sadler, G. R., Klonoff, E. A., & Malcarne, V. L. (2017). Reliability and validity of the Perceived Stress Scale-10 in Hispanic Americans with English or Spanish language preference. *Journal of Health Psychology*, *1359105316684938*. <https://doi.org/10.1177/1359105316684938>.
- Bastiaansen, D., Koot, H. M., Bongers, I. L., Varni, J. W., & Verhulst, F. C. (2004). Measuring quality of life in children referred for psychiatric problems: psychometric properties of the PedsQL 4.0 generic core scales. *Quality of Life Research*, *13*(2), 489–495. <https://doi.org/10.1023/B:QURE.0000018483.01526.ab>.
- Bates, R., Salsberry, P., & Ford, J. (2017). Measuring stress in young children using hair cortisol: The state of the science. *Biological Research for Nursing*, *19*(5), 499–510. <https://doi.org/10.1177/1099800417711583>.
- Becerra, B. J., Sis-Medina, R. C., Reyes, A., & Becerra, M. B. (2015). Association between food insecurity and serious psychological distress among Hispanic adults living in poverty. *Preventing Chronic Disease*, *12*, E206. <https://doi.org/10.5888/pcd12.150334>.
- Beck, A. F., Tschudy, M. M., Coker, T. R., Mistry, K. B., Cox, J. E., Gitterman, B. A., Chamberlain, L. J., Grace, A. M., Hole, M. K., Klass, P. E., Lobach, K. S., Ma, C. T., Navsaria, D., Northrip, K. D., Sadof, M. D., Shah, A. N., & Fierman, A. H. (2016). Determinants of health and pediatric primary care practices. *Pediatrics*, *137*(3). <https://doi.org/10.1542/peds.2015-3673>.
- Berger, M., & Sarnyai, Z. (2015). “More than skin deep”: Stress neurobiology and mental health consequences of racial discrimination. *Stress*, *18*(1), 1–10. <https://doi.org/10.3109/10253890.2014.989204>.
- Bethell, C. D., Garner, A. S., Gombojav, N., Blackwell, C., Heller, L., & Mendelson, T. (2022). Social and relational health risks and common mental health problems among US children: The mitigating role of family resilience and connection to promote positive socioemotional and school-related outcomes. *Child and Adolescent Psychiatric Clinics of North America*, *31*(1), 45–70. <https://doi.org/10.1016/j.chc.2021.08.001>.
- Bitsko, R. H., Holbrook, J. R., Ghandour, R. M., Blumberg, S. J., Visser, S. N., Perou, R., & Walkup, J. T. (2018). Epidemiology and impact of health care provider-diagnosed anxiety and depression among US children. *Journal of Developmental and Behavioral Pediatrics*, *39*(5), 395–403. <https://doi.org/10.1097/dbp.0000000000000571>.
- Braren, S. H., Perry, R. E., Ursache, A., & Blair, C. (2019). Socioeconomic risk moderates the association between caregiver cortisol levels and infant cortisol reactivity to emotion induction at 24 months. *Developmental Psychobiology*, *61*(4), 573–591. <https://doi.org/10.1002/dev.21832>.
- Brcic, V., Eberdt, C., & Kaczorowski, J. (2011). Development of a tool to identify poverty in a family practice setting: a pilot study. *International Journal of Family Medicine*, *2011*, 812182. <https://doi.org/10.1155/2011/812182>.
- Bryson, H. E., Goldfeld, S., Price, A. M. H., & Mensah, F. (2019). Hair cortisol as a measure of the stress response to social adversity in young children. *Developmental Psychobiology*, *61*(4), 525–542. <https://doi.org/10.1002/dev.21840>.
- Bryson, H. E., Mensah, F., Goldfeld, S., Price, A. M. H., & Giallo, R. (2020). Hair cortisol in mother-child dyads: examining the roles of maternal parenting and stress in the context of early childhood adversity. *European Child & Adolescent Psychiatry*. <https://doi.org/10.1007/s00787-020-01537-0>.
- Bustamante, A. V., Félix-Beltrán, L., Nwadiuko, J., & Ortega, A. N. (2022). Avoiding Medicaid enrollment after the reversal of the changes in the public charge rule among Latino and Asian immigrants. *Health Services Research*, *57*(Suppl 2), 195–203. <https://doi.org/10.1111/1475-6773.14020>.
- Ceballos, M., & Palloni, A. (2010). Maternal and infant health of Mexican immigrants in the USA: The effects of acculturation, duration, and selective return migration. *Ethnicity and Health*, *15*(4), 377–396. <https://doi.org/10.1080/13557858.2010.481329>.
- Council on Community Pediatrics/Gitterman, B. A., Flanagan, P. J., Cotton, W. H., Dilley, K. J., Duffee, J. H., Green, A. E., Keane, V. A., Krugman, S. D., Linton, J. M., McKelvey, C. D., & Nelson, J. L. (2016). Poverty and child health in the United States. *Pediatrics*, *137*(4). <https://doi.org/10.1542/peds.2016-0339>.
- Crear-Perry, J., Correa-de-Araujo, R., Lewis Johnson, T., McLemore, M. R., Neilson, E., & Wallace, M. (2020). Social and Structural Determinants of Health Inequities in Maternal Health. *Journal of Women's Health*, *30*(2), 230–235. <https://doi.org/10.1089/jwh.2020.8882>.
- Dauegaard, S., Olsen, N. J., Heitmann, B. L., & Larsen, S. C. (2020). Familial associations in hair cortisol concentration: A cross-sectional analysis based on the Healthy Start study. *Psychoneuroendocrinology*, *121*, 104836. <https://doi.org/10.1016/j.psyneuen.2020.104836>.
- Davis, K. E. (2014). *Expenditures for treatment of mental health disorders among children, ages 5–17, 2009–2011: Estimates for the u.s. civilian noninstitutionalized population*. Agency for Healthcare Research and Quality. <https://www.ncbi.nlm.nih.gov/books/NBK476259/>.
- de Kruijff, I., Noppe, G., Kieviet, N., Choenni, V., Lambregtse-van den Berg, M. P., Begijn, D. G. A., Tromp, E., Dorst, K., van Rossum, E. F. C., de Rijke, Y. B., & van den Akker, E. L. T. (2020). LC-MS/MS-based reference intervals for hair cortisol in healthy children. *Psychoneuroendocrinology*, *112*, 104539. <https://doi.org/10.1016/j.psyneuen.2019.104539>.
- Distel, L. M. L., Egbert, A. H., Bohnert, A. M., & Santiago, C. D. (2019). Chronic stress and food insecurity: examining key environmental family factors related to body mass index among low-income Mexican-origin youth. *Family and Community Health*, *42*(3), 213–220. <https://doi.org/10.1097/fch.0000000000000228>.
- Doan, S. N., Venkatesh, S., Predroza, M., Tarullo, A., & Meyer, J. S. (2020). Maternal expressive suppression moderates the relations between maternal and child hair cortisol. *Developmental Psychobiology*. <https://doi.org/10.1002/dev.21983>.



- Dowd, J. B., Simanek, A. M., & Aiello, A. E. (2009). Socio-economic status, cortisol and allostatic load: A review of the literature. *International Journal of Epidemiology*, 38(5), 1297–1309. <https://doi.org/10.1093/ije/dyp277>.
- Duran, C. A. K., Cottone, E., Ruzek, E. A., Mashburn, A. J., & Grissmer, D. W. (2018). Family stress processes and children's self-regulation. *Child Development*. <https://doi.org/10.1111/cdev.13202>.
- Evens, G. W., & English, K. (2002). The environment of poverty: Multiple stressor exposure, psychophysiological stress, and socioemotional adjustment. *Child Development*, 73(4), 1238–1248.
- Fierman, A. H., Beck, A. F., Chung, E. K., Tschudy, M. M., Coker, T. R., Mistry, K. B., Siegel, B., Chamberlain, L. J., Conroy, K., Federico, S. G., Flanagan, P. J., Garg, A., Gitterman, B. A., Grace, A. M., Gross, R. S., Hole, M. K., Klass, P., Kraft, C., Kuo, A., & Cox, J. (2016). Redesigning health care practices to address childhood poverty. *Academic Pediatrics*, 16(3 Suppl), S136–S146. <https://doi.org/10.1016/j.acap.2016.01.004>.
- Filion, N., Fenelon, A., & Boudreaux, M. (2018). Immigration, citizenship, and the mental health of adolescents. *PLoS One*, 13(5), e0196859 <https://doi.org/10.1371/journal.pone.0196859>.
- Fontenot, K., Semega, J., & Kollar, M. (2018). *Income and poverty in the United States: 2017*. U. S. Census Bureau. <https://census.gov/content/dam/Census/library/publications/2018/demo/p60-263.pdf>.
- Frasquilho, D., de Matos, M. G., Marques, A., Neville, F. G., Gaspar, T., & Caldas-de-Almeida, J. M. (2016). Unemployment, parental distress and youth emotional well-being: The moderation roles of parent-youth relationship and financial deprivation. *Child Psychiatry & Human Development*, 47(5), 751–758. <https://doi.org/10.1007/s10578-015-0610-7>.
- Gao, W., Kirschbaum, C., Grass, J., & Stalder, T. (2016). LC-MS based analysis of endogenous steroid hormones in human hair. *Journal of Steroid Biochemistry & Molecular Biology*, 162, 92–99. <https://doi.org/10.1016/j.jsbmb.2015.12.022>.
- Ghandour, R. M., Sherman, L. J., Vladutiu, C. J., Ali, M. M., Lynch, S. E., Bitsko, R. H., & Blumberg, S. J. (2019). Prevalence and treatment of depression, anxiety, and conduct problems in US children. *Journal of Pediatrics*, 206(256-267), e253 <https://doi.org/10.1016/j.jpeds.2018.09.021>.
- Gottlieb, L., Hessler, D., Long, D., Amaya, A., & Adler, N. (2014). A randomized trial on screening for social determinants of health: The iScreen study. *Pediatrics*, 134(6), e1611–e1618. <https://doi.org/10.1542/peds.2014-1439>.
- Gottlieb, L. M., Adler, N. E., Wing, H., Velazquez, D., Keeton, V., Romero, A., Hernandez, M., Munoz Vera, A., Urrutia Caceres, E., Arevalo, C., Herrera, P., Bernal Suarez, M., & Hessler, D. (2020). Effects of in-person assistance vs personalized written resources about social services on household social risks and child and caregiver health: A randomized clinical trial. *JAMA Network Open*, 3(3), e200701 <https://doi.org/10.1001/jamanetworkopen.2020.0701>.
- Gottlieb, L. M., Hessler, D., Long, D., Laves, E., Burns, A. R., Amaya, A., Sweeney, P., Schudel, C., & Adler, N. E. (2016). Effects of social needs screening and in-person service navigation on child health: A randomized clinical trial. *JAMA Pediatrics*, 170(11), e162521 <https://doi.org/10.1001/jamapediatrics.2016.2521>.
- Gray, N. A., Dhana, A., Van Der Vyver, L., Van Wyk, J., Khumalo, N. P., & Stein, D. J. (2018). Determinants of hair cortisol concentration in children: A systematic review. *Psychoneuroendocrinology*, 87, 204–214. <https://doi.org/10.1016/j.psyneuen.2017.10.022>.
- Harris, R. A., & Santos, Jr., H. P. (2020). Maternal depression in Latinas and child socioemotional development: A systematic review. *PLoS One*, 15(3), e0230256 <https://doi.org/10.1371/journal.pone.0230256>.
- Hays, R. D., Schalet, B. D., Spritzer, K. L., & Cella, D. (2017). Two-item PROMIS® global physical and mental health scales. *Journal of Patient-Reported Outcomes*, 1(1), 2–2. <https://doi.org/10.1186/s41687-017-0003-8>.
- Henrikson, N. B., Blasi, P. R., Dorsey, C. N., Mettert, K. D., Nguyen, M. B., Walsh-Bailey, C., Macuiba, J., Gottlieb, L. M., & Lewis, C. C. (2019). Psychometric and pragmatic properties of social risk screening tools: A systematic review. *American Journal of Preventive Medicine*, 57(6 Suppl 1), S13–s24. <https://doi.org/10.1016/j.amepre.2019.07.012>.
- Hollenbach, J. P., Kuo, C. L., Mu, J., Gerrard, M., Gherlone, N., Sylvester, F., Ojukwu, M., & Cloutier, M. M. (2019). Hair cortisol, perceived stress, and social support in mother-child dyads living in an urban neighborhood. *Stress*, 22(6), 632–639. <https://doi.org/10.1080/10253890.2019.1604667>.
- Hopwood, C. J., Flato, C. G., Ambwani, S., Garland, B. H., & Morey, L. C. (2009). A comparison of Latino and Anglo socially desirable responding. *Journal of Clinical Psychology*, 65(7), 769–780. <https://doi.org/10.1002/jclp.20584>.
- Johnson, A. B., Mliner, S. B., Depasquale, C. E., Troy, M., & Gunnar, M. R. (2018). Attachment security buffers the HPA axis of toddlers growing up in poverty or near poverty: Assessment during pediatric well-child exams with inoculations. *Psychoneuroendocrinology*, 95, 120–127. <https://doi.org/10.1016/j.psyneuen.2018.05.030>.
- Johnson, S. L., Solomon, B. S., Shields, W. C., McDonald, E. M., McKenzie, L. B., & Gielen, A. C. (2009). Neighborhood violence and its association with mothers' health: assessing the relative importance of perceived safety and exposure to violence. *Journal of Urban Health*, 86(4), 538–550. <https://doi.org/10.1007/s11524-009-9345-8>.
- Karpman, M., Zuckerman, S., & Gonzalez, D. (2018). *Material hardship among nonelderly adults and their families in 2017*. Urban Institute. <https://www.urban.org/research/publication/material-hardship-among-nonelderly-adults-and-their-families-in-2017>.
- Keller, D., Jones, N., Savageau, J. A., & Cashman, S. B. (2008). Development of a brief questionnaire to identify families in need of legal advocacy to improve child health. *Ambulatory Pediatrics*, 8(4), 266–269. <https://doi.org/10.1016/j.ambp.2008.04.004>.
- Khoury, J. E., Bosquet Enlow, M., Plamondon, A., & Lyons-Ruth, K. (2019). The association between adversity and hair cortisol levels in humans: A meta-analysis. *Psychoneuroendocrinology*, 103, 104–117. <https://doi.org/10.1016/j.psyneuen.2019.01.009>.
- Kleinman, R. E., Murphy, J. M., Wieneke, K. M., Desmond, M. S., Schiff, A., & Gapinski, J. A. (2007). Use of a single-question screening tool to detect hunger in families attending a neighborhood health center. *Ambulatory Pediatrics*, 7(4), 278–284. <https://doi.org/10.1016/j.ambp.2007.03.005>.
- Koumantarou Malisiova, E., Mourikis, I., Chalimourdas, T., Nianiakas, N., Michou, M., Mantzou, A., Darviri, C., Vaidakis, N., Zervas, I. M., Chrousos, G. P., & Papageorgiou, C. C. (2020). Low hair cortisol concentrations in obsessive compulsive disorder: A cross-sectional study. *Journal of Psychiatric Research*, 131, 187–193. <https://doi.org/10.1016/j.jpsychires.2020.09.014>.
- Kreuter, M. W., Thompson, T., McQueen, A., & Garg, R. (2021). Addressing social needs in health care settings: Evidence, challenges, and opportunities for public health. *Annual Review of Public Health*, 42, 329–344. <https://doi.org/10.1146/annurev-publichealth-090419-102204>.
- Kroenke, K., Spitzer, R. L., Williams, J. B., & Lowe, B. (2010). The Patient Health Questionnaire somatic, anxiety, and depressive symptom scales: A systematic review. *General Hospital*

- Psychiatry*, 32(4), 345–359. <https://doi.org/10.1016/j.genhosppsy.2010.03.006>.
- Larson, K., Russ, S. A., Crall, J. J., & Halfon, N. (2008). Influence of multiple social risks on children's health. *Pediatrics*, 121(2), 337–344. <https://doi.org/10.1542/peds.2007-0447>.
- Lawton, K. E., Gerdes, A. C., Haack, L. M., & Schneider, B. (2014). Acculturation, cultural values, and Latino parental beliefs about the etiology of ADHD. *Administration and Policy in Mental Health*, 41(2), 189–204. <https://doi.org/10.1007/s10488-012-0447-3>.
- Ling, J., Robbins, L. B., & Xu, D. (2019). Food security status and hair cortisol among low-income mother-child dyads. *Western Journal of Nursing Research*, 41(12), 1813–1828. <https://doi.org/10.1177/0193945919867112>.
- Ling, J., Xu, D., Robbins, L. B., & Meyer, J. S. (2020). Does hair cortisol really reflect perceived stress? Findings from low-income mother-preschooler dyads. *Psychoneuroendocrinology*, 111, 104478. <https://doi.org/10.1016/j.psyneuen.2019.104478>.
- Linton, J. M., Choi, R., & Mendoza, F. (2016). Caring for Children in Immigrant Families: Vulnerabilities, Resilience, and Opportunities [Review]. *Pediatric Clinics of North America*, 63(1), 115–130. <https://doi.org/10.1016/j.pcl.2015.08.006>.
- Lorenzo-Blanco, E. I., Meca, A., Unger, J. B., Romero, A., Szapocznik, J., Pina-Watson, B., Cano, M. A., Zamboanga, B. L., Baezconde-Garbanati, L., Des Rosiers, S. E., Soto, D. W., Villamar, J. A., Lizzi, K. M., Pattarroyo, M., & Schwartz, S. J. (2017). Longitudinal effects of latino parent cultural stress, depressive symptoms, and family functioning on youth emotional well-being and health risk behaviors. *Family Process*, 56(4), 981–996. <https://doi.org/10.1111/famp.12258>.
- Ludmer Nofech-Mozes, J. A., Jamieson, B., Gonzalez, A., & Atkinson, L. (2020). Mother-infant cortisol attunement: Associations with mother-infant attachment disorganization. *Developmental Psychopathology*, 32(1), 43–55. <https://doi.org/10.1017/s0954579418001396>.
- Luecken, L. J., Lin, B., Coburn, S. S., MacKinnon, D. P., Gonzales, N. A., & Crnic, K. A. (2013). Prenatal stress, partner support, and infant cortisol reactivity in low-income Mexican American families. *Psychoneuroendocrinology*, 38(12), 3092–3101. <https://doi.org/10.1016/j.psyneuen.2013.09.006>.
- Macias Gil, R., Marcelin, J. R., Zuniga-Blanco, B., Marquez, C., Mathew, T., & Piggott, D. A. (2020). COVID-19 pandemic: Disparate health impact on the Hispanic/Latinx population in the United States. *Journal of Infectious Disease*, 222(10), 1592–1595. <https://doi.org/10.1093/infdis/jiaa474>.
- Marrast, L., Himmelstein, D. U., & Woolhandler, S. (2016). Racial and ethnic disparities in mental health care for children and young adults: A national study. *International Journal of Health Services*, 46(4), 810–824. <https://doi.org/10.1177/0020731416662736>.
- Meyer, B. D., Mok, W. K. C., & Sullivan, J. (2015). *Household surveys in crisis*. National Bureau of Economic Research. <https://www.nber.org/papers/w21399>.
- Miller, D. P., John, R. S., Yao, M., & Morris, M. (2022). The 2016 Presidential Election, the Public Charge Rule, and Food and Nutrition Assistance Among Immigrant Households. *Am J Public Health*, 112(12), 1738–1746. <https://doi.org/10.2105/ajph.2022.307011>.
- Neckerman, K. M., Garfinkel, I., Teitler, J. O., Waldfogel, J., & Wimer, C. (2016). Beyond income poverty: Measuring disadvantage in terms of material hardship and health. *Academic Pediatrics*, 16(3 Suppl), S52–S59. <https://doi.org/10.1016/j.acap.2016.01.015>.
- Noppe, G., Van Rossum, E. F., Koper, J. W., Manenschijn, L., Bruining, G. J., de Rijke, Y. B., & van den Akker, E. L. (2014). Validation and reference ranges of hair cortisol measurement in healthy children. *Hormone Research in Paediatrics*, 82(2), 97–102. <https://doi.org/10.1159/000362519>.
- Office of Minority Health. (2021, October 12). *Profile: Hispanic/Latino Americans*. U.S. Department of Health and Human Services. <https://minorityhealth.hhs.gov/omh/browse.aspx?lvl=3&lvlid=64>.
- Ogundele, M. O. (2018). Behavioural and emotional disorders in childhood: A brief overview for paediatricians. *World Journal of Clinical Pediatrics*, 7(1), 9–26. <https://doi.org/10.5409/wjcp.v7.i1.9>.
- Ouellette, S. J., Russell, E., Kryski, K. R., Sheikh, H. I., Singh, S. M., Koren, G., & Hayden, E. P. (2015). Hair cortisol concentrations in higher- and lower-stress mother-daughter dyads: A pilot study of associations and moderators. *Developmental Psychobiology*, 57(5), 519–534. <https://doi.org/10.1002/dev.21302>.
- Palermo, F., Ispa, J. M., Carlo, G., & Streit, C. (2018). Economic hardship during infancy and U.S. Latino preschoolers' socio-behavioral health and academic readiness. *Developmental Psychology*, 54(5), 890–902. <https://doi.org/10.1037/dev0000476>.
- Paz, K., & Massey, K. P. (2016). Health disparity among Latina women: Comparison with non-Latina women. *Clinical Medicine Insights: Women's Health*, 9(Suppl 1), 71–74. <https://doi.org/10.4137/cmwh.S38488>.
- Perreira, K. M., Marchante, A. N., Schwartz, S. J., Isasi, C. R., Carnethon, M. R., Corliss, H. L., Kaplan, R. C., Santisteban, D. A., Vidot, D. C., Van Horn, L., & Delamater, A. M. (2019). Stress and resilience: Key correlates of mental health and substance use in the Hispanic community health study of Latino youth. *Journal of Immigrant & Minority Health*, 21(1), 4–13. <https://doi.org/10.1007/s10903-018-0724-7>.
- Pervanidou, P., Bastaki, D., Chouliaras, G., Papanikolaou, K., Laios, E., Kanaka-Gantenbein, C., & Chrousos, G. P. (2013). Circadian cortisol profiles, anxiety and depressive symptomatology, and body mass index in a clinical population of obese children. *Stress*, 16(1), 34–43. <https://doi.org/10.3109/10253890.2012.689040>.
- Potochnick, S. R., & Perreira, K. M. (2010). Depression and anxiety among first-generation immigrant latino youth: Key correlates and implications for future research [Article]. *Journal of Nervous and Mental Disease*, 198(7), 470–477. <https://doi.org/10.1097/NMD.0b013e3181e4ce24>.
- Prado-Gasco, V., de la Barrera, U., Sancho-Castillo, S., de la Rubia-Orti, J. E., & Montoya-Castilla, I. (2019). Perceived stress and reference ranges of hair cortisol in healthy adolescents. *PLoS One*, 14(4), e0214856. <https://doi.org/10.1371/journal.pone.0214856>.
- Raffington, L., Prindle, J., Keresztes, A., Binder, J., Heim, C., & Shing, Y. L. (2018). Blunted cortisol stress reactivity in low-income children relates to lower memory function. *Psychoneuroendocrinology*, 90, 110–121. <https://doi.org/10.1016/j.psyneuen.2018.02.002>.
- Rodem, R., & Shaefer, H. L. (2020). Many of the kids are not alright: Material hardship among children in the United States. *Children and Youth Services Review*, 112, 104767. <https://doi.org/10.1016/j.chilyouth.2020.104767>.
- Roizen, M., Rodriguez, S., Bauer, G., Medin, G., Bevilacqua, S., Varni, J. W., & Dussel, V. (2008). Initial validation of the Argentinean Spanish version of the PedsQL 4.0 Generic Core Scales in children and adolescents with chronic diseases: acceptability and comprehensibility in low-income settings. *Health and Quality of Life Outcomes*, 6, 59. <https://doi.org/10.1186/1477-7525-6-59>.
- Ruiz, J. M., Sbarra, D., & Steffen, P. R. (2018). Hispanic ethnicity, stress psychophysiology and paradoxical health outcomes: A review with conceptual considerations and a call for research. *International Journal of Psychophysiology*, 131, 24–29. <https://doi.org/10.1016/j.ijpsycho.2018.04.001>.

- Schloß, S., Müller, V., Becker, K., Skoluda, N., Nater, U. M., & Pauli-Pott, U. (2019). Hair cortisol concentration in mothers and their children: Roles of maternal sensitivity and child symptoms of attention-deficit/hyperactivity disorder. *Journal of Neural Transmission*, 126(9), 1135–1144. <https://doi.org/10.1007/s00702-018-1944-7>.
- Schmeer, K. K. (2012). Early childhood economic disadvantage and the health of Hispanic children. *Social Science & Medicine*, 75(8), 1523–1530. <https://doi.org/10.1016/j.socscimed.2012.05.031>.
- Seid, M., & Varni, J. W. (2005). Measuring primary care for children of Latino farmworkers: Reliability and validity of the parent's perceptions of primary care measure (P3C). *Maternal Child Health Journal*, 9(1), 49–57.
- Shonkoff, J. P., & Garner, A. S. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, 129(1), e232–e246. <https://doi.org/10.1542/peds.2011-2663>.
- Solarikova, P., Karailieva, L., Rajcani, J., Brezina, I., & Jezova, D. (2020). Cumulative cortisol concentrations in hair of patients with atopy are lower than in healthy subjects and are not related to their perceived stress experience. *Stress*, 1-4. <https://doi.org/10.1080/10253890.2020.1825673>.
- Stalder, T., Steudte-Schmiedgen, S., Alexander, N., Klucken, T., Vater, A., Wichmann, S., Kirschbaum, C., & Miller, R. (2017). Stress-related and basic determinants of hair cortisol in humans: A meta-analysis. *Psychoneuroendocrinology*, 77, 261–274. <https://doi.org/10.1016/j.psyneuen.2016.12.017>.
- Teruya, S. A., & Bazargan-Hejazi, S. (2013). The immigrant and Hispanic paradoxes: A systematic review of their predictions and effects. *Hispanic Journal of Behavioral Science*, 35(4), 486–509. <https://doi.org/10.1177/0739986313499004>.
- Torres, L., Driscoll, M. W., & Voell, M. (2012). Discrimination, acculturation, acculturative stress, and Latino psychological distress: A moderated mediational model. *Cultural Diversity & Ethnic Minority Psychology*, 18(1), 17–25. <https://doi.org/10.1037/a0026710>.
- Vallejo, M. A., Vallejo-Slocker, L., Fernández-Abascal, E. G., & Mañanes, G. (2018). Determining factors for stress perception assessed with the Perceived Stress Scale (PSS-4) in Spanish and other European samples. *Frontiers in Psychology*, 9, 37. <https://doi.org/10.3389/fpsyg.2018.00037>.
- Varni, J. W., Burwinkle, T. M., Seid, M., & Skarr, D. (2003). The PedsQL 4.0 as a pediatric population health measure: Feasibility, reliability, and validity. *Ambulatory Pediatrics*, 3(6), 329–341. [https://doi.org/10.1367/1539-4409\(2003\)003<0329:tpaapp>2.0.co;2](https://doi.org/10.1367/1539-4409(2003)003<0329:tpaapp>2.0.co;2).
- Varni, J. W., Limbers, C. A., & Burwinkle, T. M. (2007a). How young can children reliably and validly self-report their health-related quality of life? An analysis of 8,591 children across age subgroups with the PedsQL 4.0 Generic Core Scales. *Health & Quality of Life Outcomes*, 5, 1. <https://doi.org/10.1186/1477-7525-5-1>.
- Varni, J. W., Limbers, C. A., & Burwinkle, T. M. (2007b). Parent proxy-report of their children's health-related quality of life: An analysis of 13,878 parents' reliability and validity across age subgroups using the PedsQL 4.0 Generic Core Scales. *Health & Quality of Life Outcomes*, 5, 2. <https://doi.org/10.1186/1477-7525-5-2>.
- Varni, J. W., Seid, M., & Kurtin, P. S. (2001). PedsQL 4.0: reliability and validity of the Pediatric Quality of Life Inventory version 4.0 generic core scales in healthy and patient populations. *Medical Care*, 39(8), 800–812. <https://www.ncbi.nlm.nih.gov/pubmed/11468499>.
- Wang, S. S., Glied, S., Babcock, C., & Chaudry, A. (2022). Changes in the Public Charge Rule and Health of Mothers and Infants Enrolled in New York State's Medicaid Program, 2014–2019. *American Journal of Public Health*, 112(12), 1747–1756. <https://doi.org/10.2105/ajph.2022.307066>.
- Warttig, S. L., Forshaw, M. J., South, J., & White, A. K. (2013). New, normative, English-sample data for the ShortForm Perceived Stress Scale (PSS-4). *Journal of Health Psychology*, 18(12), 1617–1628.
- Zilanawala, A., & Pilkauskas, N. V. (2012). Material hardship and child socioemotional behaviors: Differences by types of hardship, timing, and duration. *Children and Youth Services Review*, 34(4), 814–825. <https://doi.org/10.1016/j.childyouth.2012.01.008>.