









Correction to: Precursor reactivity differentiation for single-step preparation of Ag₂Se@Ag₂S core–shell nanocrystals with distinct absorption and emission properties enabling sensitive near-infrared photodetection

Shilin Tang¹ , Chuansheng He¹ , Dong Li¹ , Wenhao Cai¹ , Louzhen Fan¹ , and Yunchao Li^{1,*} 

¹Department of Chemistry, Beijing Normal University, Beijing 100875, China

Published online:

27 May 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

J Mater Sci (2018) 53:11355
<https://doi.org/10.1007/s10853-018-2465-3>

has been updated to include the correct supplemental figure.

In the original article Fig. 3 and supplemental Figure s3 were incorrect. The correct version of Figure 3 follows and the Electronic Supplementary Material

The original article can be found online at <https://doi.org/10.1007/s10853-018-2465-3>.

Address correspondence to E-mail: liyc@bnu.edu.cn




Fig. 3 TEM, STEM and HRTEM images of the **a, b** Ag_2Se nanocrystals and **c–j** $\text{Ag}_2\text{Se}@Ag_2\text{S}$ nanocrystals prepared by using the single-step co-injection protocol with adopting various S to Se

precursor ratios: **c, d** S/Se = 1:3 (CS-1), **e, f** S/Se = 1:1 (CS-2), **g, h** S/Se = 3:1 (CS-3), **i, j** S/Se = 7:1 (CS-4). The scale bars of the STEM images inserted in **a, c, e, g, i** are 2.0 nm.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.