

---

## Errata

### Additional Flux Arising from Unresolved Scales in Eddy Ocean Models

MIKITOSHI HIRABARA, HIROSHI ISHIZAKI, GORO YAMANAKA,  
HIROYUKI TSUJINO and ICHIRO ISHIKAWA

Journal of Oceanography, Vol. 66, pp. 633–647, 2010

p. 638, Equation;

$$F_{Cx} = \frac{1}{\Delta_y} \int_{-\Delta_y/2}^{\Delta_y/2} \left[ (\langle u \rangle + \langle u_y \rangle x) (\langle \tau \rangle + \langle \tau_x \rangle x) - \langle v \rangle \langle \tau \rangle \right] dx$$

should be read as

$$F_{Cx} = \frac{1}{\Delta_y} \int_{-\Delta_y/2}^{\Delta_y/2} \left[ (\langle u \rangle + \langle u_y \rangle y) (\langle \tau \rangle + \langle \tau_y \rangle y) - \langle u \rangle \langle \tau \rangle \right] dy$$

p. 638, Equation;

$$F_{Cy} = \frac{1}{\Delta_x} \int_{-\Delta_y/2}^{\Delta_y/2} \left[ (\langle v \rangle + \langle v_x \rangle x) (\langle \tau \rangle + \langle \tau_x \rangle x) - \langle v \rangle \langle \tau \rangle \right] dx$$

should be read as

$$F_{Cy} = \frac{1}{\Delta_x} \int_{-\Delta_x/2}^{\Delta_x/2} \left[ (\langle v \rangle + \langle v_x \rangle x) (\langle \tau \rangle + \langle \tau_x \rangle x) - \langle v \rangle \langle \tau \rangle \right] dx$$