The Effect of Blockade of Tumor Necrosis Factor α on VLA-1+T-Cells in Rheumatoid Arthritis Patients

**Ilan Bank • Shomron Ben-Horin • Itamar Goldstein •
Alexander Koltakov • Pnina Langevitz •
Michael Ehrenfeld • Esther Rosenthal • Hanan Gur**

Published online: 2 November 2007
© Springer Science + Business Media, LLC 2007

The order of authors has been corrected as follows: Shomron Ben-Horin, Itamar Goldstein, Alexander Koltakov, Pnina Langevitz, Michael Ehrenfeld, Esther Rosenthal, Hanan Gur and Ilan Bank.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10875-007-9119-6>.

I. Bank (✉)
Medicine and Laboratory for Immunoregulation,
Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel
e-mail: ibank@post.tau.ac.il

I. Bank • P. Langevitz • M. Ehrenfeld • H. Gur
Sackler School of Medicine, Tel Aviv University,
Tel Aviv, Israel

S. Ben-Horin
Gastroenterology, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

I. Goldstein
Sheba Cancer Research Center, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

A. Koltakov
Laboratory for Immunoregulation, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

P. Langevitz
Medicine and Rheumatology, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

M. Ehrenfeld
Medicine, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

E. Rosenthal
Hematology, Chaim Sheba Medical Center,
Tel Hashomer 52621, Israel

H. Gur
Medicine, Souraski Medical Center,
6, Weizman Street,
64239 Tel Aviv, Israel