CORRECTION

Correction to: Changes in transcranial motor evoked potentials during hemorrhage are associated with increased serum propofol concentrations

Jeremy A. Lieberman^{1,3} · John Feiner¹ · Mark Rollins¹ · Russ Lyon² · Paul Jasiukaitis⁴

Published online: 8 November 2017
© Springer Science+Business Media B.V. 2017

Correction to: J Clin Monit Comput
DOI 10.1007/s10877-017-0057-4

In the original publication of the article, the corresponding author inadvertently omitted one of the co-authors in the author group. The corrected author group is given in this erratum.

The online version of the original article can be found under doi:[10.1007/s10877-017-0057-4](https://doi.org/10.1007/s10877-017-0057-4).

✉ Jeremy A. Lieberman
lieberman@anesthesia.ucsf.edu

Paul Jasiukaitis
cochransci@sbcglobal.net

¹ Department of Anesthesia & Perioperative Care, UCSF, San Francisco, CA, USA

² Division of Operating Rooms, UCSF, San Francisco, CA, USA

³ Division of Spine Anesthesia, Department of Anesthesia & Perioperative Care, University of California, San Francisco, 4th Floor, Millberry Union, East Tower Room MUE-413, 500 Parnassus Avenue, Box 0648, San Francisco, CA 94143-0648, USA

⁴ Division of Operating Rooms, University of California, San Francisco, San Francisco, CA, USA