

Erratum to: Synergism between Enantiomers Creates Species-Specific Pheromone Blends and Minimizes Cross-Attraction for Two Species of Cerambycid Beetles

Linnea R. Meier¹ · Yunfan Zou² · Jocelyn G. Millar² · Judith A. Mongold-Diers¹ · Lawrence M. Hanks¹

Published online: 26 November 2016
© Springer Science+Business Media New York 2016

Erratum to: J Chem Ecol
DOI 10.1007/s10886-016-0782-z

In the original version of this article, a line on the fourth page, third paragraph, has the retention times for (*R*)- and (*S*)-fuscumol acetate reversed. The correct sentence should read, “Structures were confirmed by coinjection of an aliquot of aeration extract with the mixture of synthetic stereoisomers (Millar et al. 2009); (*S*) - and (*R*)-fuscumol acetate (retention times 51.33 and 51.72 min, respectively) and (*Z*) - and (*E*)-geranylacetone were resolved to baseline (retention times 14.53 and 14.74 min, respectively).” The authors regret this error.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10886-016-0782-z>

✉ Lawrence M. Hanks
hanks@life.illinois.edu

¹ Department of Entomology, University of Illinois at Urbana-Champaign, 320 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801, USA

² Department of Entomology, University of California, Riverside, CA 92521, USA