Author Correction: Temporal Variation of White Rhino Dung Odours

Courtney Marneweck¹ • Andreas Jürgens² • Adrian M. Shrader^{1,3}

Published online: 8 March 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Author Correction: Journal of Chemical Ecology (2017) 43:955–965
<https://doi.org/10.1007/s10886-017-0890-4>

The original version of this article unfortunately contained some mistakes. Firstly, Fig. 5 listed a contaminant. The correct version of Fig. 5 can be found here. Second, an incorrect version of the Supplementary Table A1 was mistakenly included. A corrected and updated version of Supplementary Table A1 can be found here as well, pertaining specifically to the important compounds highlighted in Fig. 5 and with additional chemical information included.

The authors apologize for this oversight and for any confusion it may have caused.

The online version of the original article can be found at <https://doi.org/10.1007/s10886-017-0890-4>

✉ Courtney Marneweck
courtney.marneweck@gmail.com

¹ School of Life Sciences, University of KwaZulu-Natal, Durban, South Africa

² Chemical Plant Ecology, Technische Universität Darmstadt, Darmstadt, Germany

³ Present address: Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa

Fig. 5 The importance of volatile organic compounds (VOCs) for distinguishing the age of dung from territorial males, **a**) wet season, **b**) dry season; non-territorial males **c**) wet season, **d**) dry season; estrus

females **e**) wet season, **f**) dry season; and non-estrus females **g**) wet season, **h**) dry season. Importance was based on mean decrease in accuracy (MDA). Only the top ten compounds are presented in the figure

Table A1 Chemical information pertaining to the tentative identification of compounds listed in Fig. 5 (the importance of volatile organic compounds for distinguishing the age of dung)

Compound	Retention time (min.)	Retention index RI (calc.)	Retention Index RI (lit.)	Observed Match and R. Match values	Lit. Source of RI/Lit. Source of MS (NIST 2013)	MS-Spectrum in Sample (most abundant peaks)
Alkanes						
Nonane ^c	6.41	900	900	Match: 813 R.Match: 890		
Alkenes						
(<i>E</i>)-2-Octene ^b	4.32	799	798	Match: 799 R.Match: 886	Xu et al. (2003)	
(<i>Z</i>)-2-Octene ^b	4.38	802	817	Match: 779 R.Match: 853	Garcia-Estaban et al. (2004)	
(<i>E</i>)-3-Decene ^a	7.77	961	–	Match: 801 R.Match: 854	41=999 55=868 69=691 56=630 43=483 70=482 39=320 42=311 29=273 27=263	69= 999 41= 732 70= 651 55= 615 56= 348 43= 226 140= 222 39= 207 57= 199 67= 194
(<i>Z</i>)-3-Dodecene ^a	13.06	1193	–	Match: 828 R.Match: 886	55=999 41=949 56=898 69=881 43=715 70=650 57=582 83=396 42=331 29=287	55= 999 43= 751 41= 731 69 = 725 56= 667 70= 551 83= 410 105= 409 77= 407 84= 363
Aliphatic Aldehydes						
Octanal ^c	8.45	991	1004	Match: 786 R.Match: 903	Moio et al. (1996)	
Nonanal ^c	10.98	1112	1103	Match: 808 R.Match: 850	Flamini et al. (2007)	
Decanal ^c	13.17	1198	1205	Match: 862 R.Match: 887	Baccouri et al. (2007)	
Aliphatic Ketones						
Pentan-2-one ^b	2.43	664	691	Match: 848 R.Match: 859	Macku and Shibamoto (1991)	
Hexan-2-one ^b	4.03	780	790	Match: 856 R.Match: 905	Wu et al. (2007)	
Undecan-2-one ^b	14.61	1269	1291	Match: 652 R.Match: 858	Adams (1999)	
Aliphatic Acids						
Acetic acid ^c	1.82	601	600	Match: 901 R.Match: 920	Whetstone et al. (2005)	
Isobutanoic acid ^b	3.39	740	765	Match: 808 R.Match: 870	Mateo and Zumalacarregui (1996)	
Butanoic acid ^b	4.28	797	790	Match: 813 R.Match: 820	Quijano et al. (2007)	
3-Methylbutanoic acid ^b	4.98	831	831	Match: 664 R.Match: 724	Ventanas et al. (2008)	
2-Methylbutanoic acid ^b	5.40	851	853	Match: 690 R.Match: 745	Kondjoyan et al. (1997)	
Aromatic compounds						
Toluene ^c	3.73	761	762	Match: 923 R.Match: 950	Moon et al. (2006)	
<i>p</i> -Xylene ^c	5.70	866	869	Match: 824 R.Match: 922	Helmig et al. (1996)	
Styrene ^c	6.22	891	894	Match: 773 R.Match: 837	Insausti et al. (2005)	
Benzaldehyde ^c	7.77	960	948	Match: 817 R.Match: 892	Jordan et al. (2001)	
<i>p</i> -Cymene ^b	9.26	1030	1060	Match 643 R.Match: 683	Marongiu et al. (2004)	
Acetophenone ^c	10.14	1073	1062	Match: 914 R.Match: 946	Schwambach and Peterson (2006)	
<i>m</i> -Cresol ^c	10.53	1093	1073	Match: 925 R.Match: 934	Re-Poppi and Santiago-Silva (2002)	
Terpenoids						
Tricyclene ^b	6.98	925	923	Match: 876 R.Match: 910	Pino et al. (2005)	
Camphene ^b	7.62	954	945	Match: 919 R.Match: 962	Zhang et al. (2005)	
3,4-Dihydro- β -ocimene ^b	7.40	944	939	Match: 915 R.Match: 916	Diaz and Kite (2002)	
(<i>E</i>)-1,4-Dihydro- β -ocimene ^b	8.33	985	990	Match: 899 R.Match: 959	Buchin et al. (2002)	
(<i>Z</i>)-1,4-Dihydro- β -ocimene ^b	8.64	999	990	Match: 866 R.Match: 928	Buchin et al. (2002)	

Table A1 (continued)

Compound	Retention time (min.)	Retention index RI (calc.)	Retention Index RI (lit.)	Observed Match and R. Match values	Lit. Source of RI /Lit. Source of MS (NIST 2013)	MS-Spectrum in Sample (most abundant peaks)
Unknown terpenoid A ^a	7.07	929	—	—	—	55= 999 41= 981 67= 847 39= 653 68= 473 95= 406 53= 272 56= 262 69= 260 81= 245
Limonene ^c	9.41	1037	1039	Match: 907 R.Match: 927	Blagojevic et al. (2006)	
β-Caryophyllene ^c	17.72	1442	1440	Match: 904 R.Match: 908	Bylaite and Meyer (2006)	
Cyclosativene ^b	16.73	1393	1370	Match: 886 R.Match: 893	Andrade et al. (2007)	
β-Gurjunene ^b	17.91	1451	1428	Match: 911 R.Match: 936	Saroglou et al. (2007)	
Geranylacetone ^b	17.96	1453	1445	Match: 756 R.Match: 819	Dhanda et al. (2003)	
Unknown terpenoid B ^a	19.49	1533	—	—	—	161= 999 122 = 670 107 = 588 105= 399 91 = 389 81 = 355 204 =338 79 = 305 41 = 299 93 = 273
Unknown terpenoid C ^a	23.17	1779	—	—	—	32 = 999 43 = 844 70 = 736 55 = 669 69 = 621 71 = 567 57 = 564 41 = 562 56 = 312 58 = 289

^a comparison of MS with published data^b comparison of MS and retention time with published data^c comparison of MS and retention time with authentic standard

References

- Adams RP (1999) Systematics of multi-seeded eastern hemisphere *Juniperus* based on leaf essential oils and RAPD DNA fingerprinting. *Biochem Syst Ecol* 27:709–725
- Andrade EHA, Oliveira J, Zoghbi MGB (2007) Volatiles of *Anaxagorea dolichocarpa* Spreng. & Sandw. and *Annona densicoma* Mart. Growing Wild in the State of Para, Brazil. *Flavour Fragr J* 22:158–160
- Baccouri B, Ben Temime S, Campeol E, Cioni PL, Daoud D, Zarrouk M (2007) Application of solid-phase microextraction to the analysis of volatile compounds in virgin olive oils from five new cultivars. *Food Chem* 102:850–856
- Blagojevic P, Radulovic N, Palic R, Stojanovic G (2006) Chemical composition of the essential oils of Serbian wild-growing *Srtemisia absinthium* and *Artemisia vulgaris*. *J Agric Food Chem* 54:4780–4789
- Buchin S, Salmon J-C, Carnat A-P, Berger T, Bugaud C, Bosset JO (2002) Identification de composés monoterpéniques, sesquiterpéniques et benzeniques dans un lait d'alpage très riche en ces substances. *Mitt. Lebensmittelunters. Hyg.* 93:199–216
- Bylaite E, Meyer AS (2006) Characterisation of volatile aroma compounds of orange juices by three dynamic and static headspace gas chromatography techniques. *Eur Food Res Technol* 222:176–184
- Dhanda JS, Pegg RB, Shand PJ (2003) Saskatchewan specialty livestock value-added program. Saskatchewan agri-food innovation fund (AFIF) Project #98000016
- Diaz A, Kite GC (2002) A comparison of the pollination ecology of *Arum maculatum* and *A. italicum* in England, Watsonia. 24:171–181
- Flamini G, Cioni PL, Morelli I, Bader A (2007) Essential oils of the aerial parts of three *Salvia* species from Jordan: *Salvia lanigera*, *S. spinosa* and *S. syriaca*. *Food Chem*, 100:732–735
- Garcia-Estaban M, Ansorena D, Astiasaran I, Ruiz J (2004) Study of the effect of different fiber coatings and extraction conditions on dry cured ham volatile compounds extracted by solid-phase microextraction (SPME). *Talanta* 64:458–466
- Helmig D, Pollock W, Greenberg J, Zimmerman P (1996) Gas chromatography mass spectrometry analysis of volatile organic trace gases at Mauna Loa Observatory, Hawaii. *J Geophys Res* 101:14697–14710
- Insausti K, Goni V, Petri E, Gorraiz C, Beriain MJ (2005) Effect of weight at slaughter on the volatile compounds of cooked beef from Spanish cattle breeds. *Meat Sci* 70:83–90
- Jordan MJ, Shaw PE, Goodner KL (2001) Volatile components in aqueous essence and fresh fruit of *Cucumis melo* cv. Athena (muskmelon) by GC-MS and GC-O. *J Agric Food Chem* 49:5929–5933
- Kondjoyan N, Viallon C, Berdague JL, Daridan D, Simon M-N, Legault C (1997) Analyse comparative de la fraction volatile de jambons secs de porcs Gascon et Large-White x Landrace Français. *J Rech CNRS* 29:405–410
- Macku C, Shibamoto T (1991) Headspace volatile compounds formed from heated corn oil and corn oil with glycine. *J Agric Food Chem* 39:1265–1269

- Marongiu B, Piras A, Porcedda S (2004) Comparative analysis of the oil and supercritical CO₂ extract of *Elettaria cardamomum* (L.) maton. J Agric Food Chem 52:6278–6282
- Mateo J, Zumalacarregui JM (1996) Volatile compounds in chorizo and their changes during ripening. Meat Sci 44:255–273
- Moio L, Rillo L, Ledda A, Addeo F (1996) Odorous constituents of ovine milk in relationship to diet. J Dairy Sci 79:1322–1331
- Moon S-Y, Cliff MA, Li-Chan ECY (2006) Odour-active components of simulated beef flavour analysed by solid phase microextraction and gas chromatography-mass spectrometry and -olfactometry. Food Res Int 39:294–308
- Pino JA, Mesa J, Munoz Y, Marti MP, Marbot R (2005) Volatile components from mango (*Mangifera indica* L.) cultivars. J Agric Food Chem 53:2213–2223
- Quijano CE, Salamanca G, Pino JA (2007) Aroma volatile constituents of Colombian varieties of mango (*Mangifera indica* L.). Flavour Fragr J 22:401–406
- Re-Poppi N, Santiago-Silva MR (2002) Identification of polycyclic aromatic hydrocarbons and methoxylated phenols in wood smoke emitted during production of charcoal. Chromatographia 55:475–481
- Saroglou V, Marin PD, Rancic A, Veljic M, Skaltsa H (2007) Composition and antimicrobial activity of the essential oil of six *Hypericum* species from Serbia. Biochem Syst Ecol 35:146–152
- Schwambach SL, Peterson DG (2006) Reduction of Stale Flavor Development in Low-Heat Skim Milk Powder via Epicatechin Addition. J Agric Food Chem 54:502–508
- Ventanas S, Estevez M, Andres AI, Ruiz J (2008) Analysis of volatile compounds of Iberian dry-cured loins with different intramuscular fat contents using SPME-DED. Meat Sci 79:172–180
- Whetstone MEC, Cadwallader KR, Drake MA (2005) Characterization of aroma compounds responsible for the rosy/floral flavor in cheddar cheese. J Agric Food Chem 53:3126–3132
- Wu S, Zom H, Krings U, Berger RG (2007) Volatiles from submerged and surface-cultured beefsteak fungus, *Fistulina hepatica*. Flavour Fragr J 22:53–60
- Xu X et al. (2003) Comprehensive two-dimensional gas chromatography (GC*GC) measurements of volatile organic compounds in the atmosphere. Atmos Chem Phys, 3:665–682
- Zhang F, Xu Q, Fu S, Ma X, Xiao H, Liang X (2005) Chemical constituents of the essential oil of *Asarum forbesii* Maxim (Aristolochiaceae). Flavour Fragr J 20:318–320