

An ICT-Based Hydrogen Sulfide Sensor with Good Water Solubility for Fluorescence Imaging in Living Cells

Qiuyan Xu¹ · Longwei He¹ · Haipeng Wei¹ · Weiyang Lin^{1,2}

Received: 19 March 2015 / Accepted: 4 May 2015 / Published online: 19 May 2015
© Springer Science+Business Media New York 2015

Introduction

Hydrogen sulfide (H₂S), traditionally considered to be a toxic gas with the typical smell of rotten eggs and an antioxidant or scavenger for reactive oxygen species (ROS) [1], has recently emerged as a member of the endogenous gaseous transmitter family of signaling molecules including nitric oxide (NO) and carbon monoxide (CO) [2–3]. The endogenous H₂S is produced by the enzymes (such as cystathionine β-synthase, cystathionine γ-lyase, and 3-mercaptopyruvate sulphurtransferase) catalysis action of cysteine or cysteine derivatives in mammalian tissues [4–6]. H₂S plays an important physiological role in many biological processes, for instance, ischemia reperfusion injury, vasodilation, apoptosis, insulin signaling, and oxygen sensing [7–11]. However, abnormal levels of H₂S is associated with various diseases, like Alzheimer's disease [12], Down's syndrome [13], diabetes [10], and liver cirrhosis [14]. Therefore, it's very significant for monitoring H₂S using sensitive, selective, and water-soluble fluorescent probe in the native biological environment.

Electronic supplementary material The online version of this article (doi:10.1007/s10895-015-1582-5) contains supplementary material, which is available to authorized users.

✉ Weiyang Lin
weiyanglin2013@163.com

¹ State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha, Hunan 410082, China


² Institute of Fluorescent Probes for Biological Imaging, School of Chemistry and Chemical Engineering, School of Biological Science and Technology, University of Jinan, Jinan, Shandong 250022, People's Republic of China

Fluorescence imaging has been widely used as a powerful tool for monitoring biomolecules within the context of living systems with high spatial and temporal resolution [15–16]. Naphthalimide dye is the favorable building block for constructing fluorescent probes because of its excellent photophysical properties, such as high extinction coefficients, excellent quantum yields, and great photostability. In addition, azido group is well-known for sensitively and selectively responding to H₂S, the electron withdrawing azide is very easy reduced by H₂S affording the electron donating amine [17–26]. Thus, in this work, we employed the naphthalimide chromophore as the signal reporter and azido group as the responding site for H₂S to construct a H₂S probe based on internal charge transfer (ICT) process (Scheme 1). In addition, to enhance the water solubility, we introduced a hydrophilic alcoholic group to the naphthalimide core. Prior to react with H₂S, we supposed that probe **Nap-N₃** has negligible fluorescence. However, interaction of naphthalimide azide with H₂S affords 4-amino-1,8-naphthalimide, which would elicit a significant fluorescence enhancement via ICT process caused by the electron donating amine and the electron withdrawing amide. So, probe **Nap-N₃** might be employed as a turn-on fluorescent probe for detecting H₂S.

Experimental

Materials and Instruments

Unless otherwise stated, all reagents were purchased from commercial suppliers and used without further purification. Solvents used were purified by standard methods prior to use. Twice-distilled water was used throughout all experiments; High resolution mass spectrometric (HRMS) analyses were measured on a Finnigan MAT 95 XP spectrometer;



Scheme 1 A turn-on fluorescent H₂S probe based on the proposed ICT switching mechanism

NMR spectra were recorded on an INOVA-400 spectrometer, using TMS as an internal standard; Electronic absorption spectra were obtained on a LabTech UV Power spectrometer; Photoluminescent spectra were recorded with a HITACHI F4600 fluorescence spectrophotometer with a 1 cm standard quartz cell; The fluorescence imaging of cells was performed with OLYMPUS FV1000 (TY1318) confocal microscopy; The pH measurements were carried out on a Mettler-Toledo Delta 320 pH meter; TLC analysis was performed on silica gel plates and column chromatography was conducted over silica gel (mesh 200–300), both of which were obtained from the Qingdao Ocean Chemicals.

Synthesis

The target compound **Nap-N₃** was readily synthesized in just two steps as shown in Scheme 2.

Synthesis of Compound **Nap-N₃**

The mixture of compounds 4-bromo-1,8-naphthalic anhydride (82.8 mg, 0.3 mmol) and ethanolamine (90 mg, 1.5 mmol) in 5 mL of ethanol was heated at 80 °C under reflux for 4 h. After cooling to room temperature, the reaction solvent was removed under reduced pressure. The resulting residue was a light yellow solid as the crude product of intermediate compound **2**, which was directly carried on the next reaction without further process. The mixture of compound **2** (16 mg, 0.05 mmol) and sodium azide (13 mg, 0.20 mmol) in 3 mL of dry N,N-dimethylformide (DMF) was heated to 90 °C for 3 h in the dark. After cooling to room temperature, the reaction mixture was poured into 100 mL of water and

then extracted three times with dichloromethane. The organic phase was collected, washed with brine, and dried with anhydrous MgSO₄. The solvent was removed under reduced pressure and the solid residue was purified by flash chromatography column using methanol/dichloromethane (*v/v* 1:20) to afford a yellow solid as compound **Nap-N₃** (10 mg, yield 70.9 %). ¹H NMR (400 MHz, CDCl₃) δ 8.64 (d, *J* = 7.2 Hz, 1H), 8.58 (d, *J* = 8.0 Hz, 1H), 8.45 (d, *J* = 8.4 Hz, 1H), 7.75 (t, *J* = 7.8 Hz, 1H), 7.47 (d, *J* = 8.0 Hz, 1H), 4.45 (t, *J* = 5.2 Hz, 2H), 3.96 (t, *J* = 5.2 Hz, 2H). ¹³C NMR (100 MHz, CDCl₃) δ 164.87, 164.50, 143.84, 132.55, 132.06, 129.13, 126.91, 124.32, 122.29, 118.52, 114.72, 61.85, 42.80. MS (EI) *m/z* 282.1 [M]⁺. HRMS (EI) *m/z* calcd for C₁₄H₁₀N₄O₃ [M]⁺: 282.0747. Found 282.0758.


Synthesis of Compound **Nap-NH₂**

Sodium sulfide (20 mg, 0.25 mmol) was added to the solution of compound **Nap-N₃** (14 mg, 0.05 mmol) in 3 mL DMF/H₂O (*v/v* 9:1) mix-solvent. The mixture was stirred for 1.5 h at room temperature in the dark. The reaction mixture was poured into 100 mL of water and then extracted two times with dichloromethane and ethyl acetate (*v/v* 5:1). The organic phase was collected, washed with brine, and dried with anhydrous MgSO₄. The solvent was removed under reduced pressure and the solid residue was purified by flash chromatography column using methanol/dichloromethane (*v/v* 1:10) to afford a yellow solid as compound **Nap-NH₂** (9.5 mg, yield 74.2 %). ¹H NMR (400 MHz, DMSO) δ 8.69 (d, *J* = 8.4 Hz, 1H), 8.42 (d, *J* = 7.2 Hz, 1H), 8.19 (d, *J* = 8.4 Hz, 1H), 7.65 (t, *J* = 8.2 Hz, 1H), 7.56 (s, 2H), 6.89 (d, *J* = 8.4 Hz, 1H), 4.13 (t, *J* = 6.6 Hz, 2H), 3.59 (m, 2H). ¹³C NMR (100 MHz, DMSO) δ 164.39, 163.49, 153.23, 134.35, 131.42, 130.18, 129.92, 124.35, 122.23, 119.79, 108.57, 107.91, 58.42, 41.78. MS (EI) *m/z* 256.1 [M]⁺. HRMS (EI) *m/z* calcd for C₁₄H₁₂N₂O₃ [M]⁺: 256.0842. Found 256.0844.

Determination of the Fluorescence Quantum Yield

Fluorescence quantum yields for compounds **Nap-N₃** and **Nap-NH₂** were determined by using fluorescein ($\Phi_f = 0.95$

Scheme 2 Synthesis of H₂S sensor **Nap-N₃** and the reduced product **Nap-NH₂**






Fig. 1 **a** Fluorescence spectra ($\lambda_{\text{ex}} = 408$ nm) of $10 \mu\text{M}$ Nap- N_3 with 0–3.5 eq. of Na_2S in 25 mM phosphate buffer (pH 7.4, containing 2 % ethanol). The inset shows the visual fluorescence color of Nap- N_3

before (left) and after (right) addition of H_2S (UV lamp, 365 nm). **b** The linear relationship between the fluorescence intensity at 546 nm and the concentration of H_2S

in 0.1 M NaOH aqueous solution) as fluorescence standard [27]. The quantum yields were calculated using the following equation.

$$\Phi_{\text{F}(X)} = \Phi_{\text{F}(S)} (A_S F_X / A_X F_S) (n_X / n_S)^2$$

Where Φ_{F} is the fluorescence quantum yield, A is the absorbance at the excitation wavelength, F is the area in the corrected emission spectrum, and n is the refractive index of the solvent used. Subscripts S and X refer to the standard and to the unknown, respectively.

HeLa Cells Culture

HeLa cells were cultured in DMEM (Dulbecco's modified Eagle's medium) supplemented with 10 % FBS (fetal bovine serum) in an atmosphere of 5 % CO_2 and 95 % air at 37 °C.

Imaging of Exogenous H_2S in Living Cells

HeLa cells were incubated with $5.0 \mu\text{M}$ Nap- N_3 for 20 min in an atmosphere of 5 % CO_2 and 95 % air, and then treated with $20 \mu\text{M}$ Na_2S for 10 min. Subsequently, the cells were imaged using OLYMPUS FV1000 (TY1318) confocal microscope with an excitation filter of 405 nm and emission channels of 520–570 nm (green channel).

Imaging of Endogenous H_2S in Living Cells

PC-3 cells were incubated with $200 \mu\text{M}$ cysteine for 60 min in an atmosphere of 5 % CO_2 and 95 % air, and then treated with $5.0 \mu\text{M}$ Nap- N_3 for 20 min. Subsequently, the cells were imaged using OLYMPUS FV1000 (TY1318) confocal microscope with an excitation filter of 405 nm and emission channels of 520–560 nm (green channel).

Results and Discussion

Fluorescence Responses to H_2S

With compound Nap- N_3 in hand, we first evaluated the capability of Nap- N_3 to detect H_2S in aqueous buffer. The titration of H_2S to the probe Nap- N_3 ($10.0 \mu\text{M}$) was performed in 25 mM PBS buffer (pH 7.4) with just 2 % ethanol. As designed, upon excitation at 438 nm, the free sensor displayed faint fluorescence at around 546 nm (Fig. 1a). However, addition of Na_2S (a standard source for hydrogen sulfide) elicited a significant emission enhancement, suggesting that Nap- N_3 was reduced affording Nap- NH_2 in the presence of H_2S and ICT process occurred. The fluorescence intensities at 546 exhibited 7.9-fold increase, it's the ICT resonance structure of compound Nap- NH_2 that contributes mainly to the enhanced fluorescence. The quantum yields of compounds Nap- N_3 and Nap- NH_2 were determined 0.0056 and 0.1175, respectively, using a reference fluorescein dye (with a quantum yield of 0.95 in 0.1 M NaOH aqueous solution). Importantly, the





Fig. 2 Absorption spectra of probe Nap- N_3 with 0–3.5 eq. of Na_2S in 25 mM phosphate buffer (pH 7.4, containing 2 % ethanol)


Fig. 3 The comparison of the partial ^1H NMR spectra of **a** Nap- N_3 in CD_3Cl and **b** Nap- NH_2 in d_6 -DMSO



chemodosimeter shows an excellent linear relationship between the fluorescent emission intensities at 546 nm and the concentrations of H_2S from 0.2 to 3.0 equivalent (Fig. 1b), suggesting that the chemodosimeter is potentially useful for quantitative determination of H_2S . The ratio changes were

observed in absorption spectra, as shown in Fig. 2, upon addition of H_2S , the absorption of naphthalimide azide at around 376 nm gradually faded and simultaneously a new red-shifted (62 nm) absorption band at around 438 nm (characteristic absorption of amino naphthalimide) was enhanced. There is

Fig. 4 Interfacial plots of the orbitals for Nap- N_3 (left) and Nap- NH_2 (right) in the excited states



a clear isosbestic point (at 407 nm) was appeared, suggesting that the conversion from **Nap-N₃** to **Nap-NH₂** is a concerted process.

Mechanism Studies

To shed light on the H₂S-triggered fluorescence turn-on response, we decided to characterize the reduced product. Incubation of **Nap-N₃** with Na₂S afforded the product, which was isolated and characterized by standard NMR and mass spectrometry (Figs. S1–3). The ¹HNMR spectra comparison of probe **Nap-N₃** and the product **Nap-NH₂**, the peak of NH₂ in **Nap-NH₂** was obviously observed, which suggested the resulting product should be reduced compound **Nap-NH₂** (Fig. 3). We further employed time-dependent density function (TD-DFT) to calculate the molecular orbital plots of **Nap-N₃** and **Nap-NH₂** by a suite of Gaussian 03 programs (6-31G(d) basis sets). As shown in Fig. 4, for the probe **Nap-N₃**, both the highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) in the excited states are distributed on the naphthalimide backbone and azido group, however, the LUMO +1 is mainly located on the electron-withdrawing azido group. The theoretical data suggested the π electrons do not transfer from the azido unit to the naphthalimide moiety and ICT process could not occur in **Nap-N₃**. By contrast, in the case of **Nap-NH₂**, the LUMO +1 is primarily resided on the naphthalimide backbone, but not on the electron-donating amino group, which indicated the π electrons transfer to the naphthalimide moiety and ICT process occurs. These experimental facts and theoretical data supported our proposed ICT switching mechanism.




Fig. 5 Fluorescence intensity (at 546 nm) of sensor **Nap-N₃** (10 μ M) in the presence of various analytes in aqueous solution (pH 7.4 PBS, containing 2 % ethanol): 1, blank; 2, F⁻; 3, Cl⁻; 4, I⁻; 5, N₃⁻; 6, CO₃²⁻; 7, SO₄²⁻; 8, HPO₄²⁻; 9, NO₂⁻; 10, NO₃⁻; 11, Ac⁻; 12, SCN⁻; 13, citrate; 14, S₂SO₃²⁻; 15, ClO⁻; 16, H₂O₂; 17, SO₃²⁻; 18, NO; 19, ascorbic acid; 20, Cys; 21, GSH; 22, Na₂S




Fig. 6 Time-dependent fluorescence intensity (at 546 nm) responses of sensor **Nap-N₃** (10 μ M) to 3 eq. Na₂S in PBS buffer

Selectivity Studies

The detection limit was calculated to be 1.09×10^{-6} M (S/N = 3), which locates in the range of physiological concentration of H₂S in vivo, indicating that the probe is sensitive to H₂S and might be suitable for detecting endogenous H₂S in biological samples. To examine the selectivity, the probe **Nap-N₃** (10 μ M) was treated with various biologically relevant species in the aqueous buffer, such as the representative anions, reactive oxygen species, reducing agents, small-molecule thiols, and Na₂S. As shown in Fig. 5, addition of the representative interfering species including F⁻, Cl⁻, I⁻, N₃⁻, CO₃²⁻, SO₄²⁻, HPO₄²⁻, NO₂⁻, NO₃⁻, AcO⁻, SCN⁻, citrate at 1 mM, and S₂O₃²⁻, SO₃²⁻, ClO⁻, H₂O₂, NO, ascorbic acid at 200 μ M induced negligible changes. Notably, small-molecule biothiols such as glutathione (GSH) and cysteine at 1 mM triggered only a small emission intensity increase.




Fig. 7 The pH influence on the fluorescence intensity (at 546 nm) of **Nap-N₃** (10 μ M) in the absence (black square) or presence (red circle) of Na₂S (30 μ M)




Fig. 8 Confocal fluorescence images of HeLa cells incubated with 5 μM Nap- N_3 for 20 min (**a, b**) and then 20 μM Na_2S for 20 min (**c, d**). Images (**a, c**) were acquired using 405 nm excitation and green emission channels of 520–570 nm; (**b, d**) bright field images. Scale bar = 40 μm

However, just upon addition of 30 μM Na_2S could elicit obvious increase of emission intensity.

Reaction-Time

In addition, the fluorescence intensity reached its maximum at about 15 min (Fig. 6). The results suggested the probe Nap- N_3 has a high selectivity for H_2S over other biological species.

pH Effect

What's more, probe Nap- N_3 could respond well to H_2S at round physiological pH (Fig. 7). The results indicated Nap- N_3 may be suitable for studies of H_2S in the living systems.

Fluorescence Imaging in Living Cells

Firstly, to examine the cell membrane permeability of naphthalimide dye, we employed compound Nap- NH_2 for imaging living cells. The cells incubated with Nap- NH_2 exhibited strong fluorescence, while control cells showed none fluorescence (Fig. S4), suggesting the naphthalimide dye is cell membrane permeable. Then to examine the utility of the sensor, we intended to image H_2S in living cells. For proof-of-concept, Nap- N_3 (5 μM) was initially incubated with HeLa cells for 20 min, after rinsed three times by PBS buffer, then treated with 20 μM Na_2S for another 20 min. As shown in

Fig. 8, the cells incubated with only Nap- N_3 exhibited faint fluorescence in the green channel. However, after co-incubation with Na_2S , strong green fluorescence was observed in living cells, suggesting Nap- N_3 was reduced by H_2S affording fluorescent compound Nap- NH_2 . Thus, these results revealed that Nap- N_3 is cell membrane permeable and capable of monitoring H_2S in living cells.

In addition to detection of extraneous H_2S , we further detected the biosynthesis H_2S inside the cells. It's well-known that cystathionine β -synthase (CBS) and cystathionine γ -lyase (CSE) could catalyze cysteine for producing H_2S in living cells [28–29]. HeLa cells were firstly incubated with 200 μM cysteine for 1 h, after rinsed three times by PBS buffer, followed addition of 5 μM Nap- N_3 and incubated another 20 min. As exhibited in Fig. 9, in comparison with the cells loaded with only the probe, the cells co-incubated with cysteine and Nap- N_3 elicited a marked increase of fluorescence intensity in the green channel. These results further indicated that Nap- N_3 is capable of detecting not only external H_2S in living cells, but also endogenous H_2S biologically produced by the cells.

Conclusions

In summary, we have introduced a high water-soluble H_2S fluorescent probe based on naphthalimide chromophore. The




Fig. 9 Confocal fluorescence images of HeLa cells incubated with 5 μM Nap- N_3 only for 20 min (**a, b**) and 200 μM cysteine for 1 h followed by 5 μM Nap- N_3 for 20 min (**c, d**). Images (**a, c**) were acquired using 405 nm

excitation and green emission channels of 520–570 nm; (**b, d**) bright field images. Scale bar = 40 μm

probe **Nap-N₃** exhibits a turn-on fluorescence signal for responding H₂S via ICT switching mechanism. It has a selectivity and sensitivity for in vitro H₂S over other various biologically relevant species, and could detect both the exogenous and endogenous H₂S in living cells.

Acknowledgments This work was financially supported by the National Science Foundation of China (Nos. 21172063, 21472067) and the startup fund of University of Jinan.

References

- Calvert JW, Jha S, Gundewar S, Elrod JW, Ramachandran A, Pattillo CB, Kevil CG, Lefler DJ (2009) Hydrogen sulfide mediates cardioprotection through Nrf2 signaling. *Circ Res* 105:365–374
- Szabó C (2007) Hydrogen sulphide and its therapeutic potential. *Nat Rev Drug Discov* 6:917–935
- Culotta E, Koshland Jr DE (1992) NO news is good news. *Science* 258:1862–1865
- Boehning D, Snyder SH (2003) Novel neural modulators. *Annu Rev Neurosci* 26:105–131
- Tanizawa K, Biochem J (2011) Production of H₂S by 3-mercaptopyruvate sulphurtransferase. *J Biochem* 149:357–359
- Hughes MN, Centelles MN, Moore KP (2009) Making and working with hydrogen sulfide: the chemistry and generation of hydrogen sulfide in vitro and its measurement in vivo: a review. *Free Radic Biol Med* 47:1346–1353
- Elrod JW, Calvert JW, Morrison J, Doeller JE, Kraus DW, Tao L, Jiao X, Scalia R, Kiss L, Szabo C, Kimura H, Chow CW, Lefler DJ (2007) Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc Natl Acad Sci U S A* 104:15560–15565
- Yang G, Wu L, Jiang B, Yang W, Qi J, Cao K, et al. (2008) H₂S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 322:587–590
- Yang G, Wu L, Wang R (2006) Pro-apoptotic effect of endogenous H₂S on human aorta smooth muscle cells. *FASEB J* 20:553–555
- Yang W, Yang G, Jia X, Wu L, Wang R (2005) Activation of K_{ATP} channels by H₂S in rat insulin-secreting cells and the underlying mechanisms. *J Physiol Lond* 569:519–531
- Peng YJ, Nanduri J, Raghuraman G, Souvannakitti D, Gadalla MM, Kumar GK, Snyder SH, Prabhakar NR (2010) H₂S mediates O₂ sensing in the carotid body. *Proc Natl Acad Sci U S A* 107:10719–10724
- Eto K, Asada T, Arima K, et al. (2002) Brain hydrogen sulfide is severely decreased in Alzheimer's disease. *Biochem Biophys Res Commun* 293:1485–1488
- Kamoun P, Belardinelli MC, Chabli A, et al. (2003) Endogenous hydrogen sulfide overproduction in Down's syndrome. *Am J Med Genet* 116A:310–311
- Fiorucci S, Antonelli E, Mencarelli A, Orlandi S, Renga B, Rizzo G, Distrutti E, Shah V, Morelli A (2005) The third gas: H₂S regulates perfusion pressure in both the isolated and perfused normal rat liver and in cirrhosis. *Hepatology* 42:539–548
- Ueno T, Nagano T (2011) Fluorescent probes for sensing and imaging. *Nat Methods* 8:642–645
- Kobayashi H, Ogawa M, Alford R, Choyke PL, Urano Y (2010) New strategies for fluorescent probe design in medical diagnostic imaging. *Chem Rev* 110:2620–2640
- Li W, Sun W, Yu X, Du L, Li M (2014) Coumarin in-based fluorescent probes for H₂S detection. *J Fluoresc* 23:181–186
- Cai Y, Li L, Wang Z, Sun JZ, Qin A, Tang BZ (2014) Sensitivity tuneable tetraphenylethene-based fluorescent probe for directly indicating the concentration of hydrogen sulfide. *Chem Commun* 50:8892–8895
- Qiao Q, Zhao M, Lang H, Mao D, Cui J, Xu Z (2014) A turn-on fluorescent probe for imaging lysosomal hydrogen sulfide in living cells. *RSC Adv* 4:25790–25794
- Ozdemir T, Sozmen F, Mamur S, et al. (2014) Fast responding and selective near-IR Bodipy dye for hydrogen sulfide sensing. *Chem Commun* 50:5455–5457
- Adarsh N, Krishnan MS, Ramaiah D (2014) Sensitive naked eye detection of hydrogen sulfide and nitric oxide by aza-BODIPY dyes in aqueous medium. *Anal Chem* 86:9335–9342
- Sun W, Fan J, Hu C, Cao J, Zhang H, Xiong X, Wang J, Cui S, Sun S, Peng X (2013) A two-photon fluorescent dye with near-infrared emission for hydrogen sulfide imaging in biosystems. *Chem Commun* 49:3890–3892
- Wan Q, Song Y, Li Z, Gao X, Ma H (2013) In vivo monitoring of hydrogen sulfide using a cresyl violet-based ratiometric fluorescence probe. *Chem Commun* 49:502–504
- Wang K, Peng H, Ni N, Dai C, Wang B (2014) 2,6-dansyl azide as a fluorescent probe for hydrogen sulfide. *J Fluoresc* 24:1–5
- Liu XL, Du XJ, Dai CG, et al. (2014) Ratiometric two-photon fluorescent probes for mitochondrial hydrogen sulfide in living cells. *J Org Chem* 79:9481–9489
- Yu C, Li X, Zeng F, Zheng F, Wu S (2013) Carbon-dot-based ratiometric fluorescent sensor for detecting hydrogen sulfide in aqueous media and inside live cells. *Chem Commun* 49:403–405
- Lakowicz JR (1999) Principles of fluorescence spectroscopy, 2nd edn. Kluwer Academic/Plenum Publishers, New York
- Qian Y, Karpus J, Kabil O, Zhang SY, Zhu HL, Banerjee R, Zhao J, He C (2011) Selective fluorescent probes for live-cell monitoring of sulphide. *Nat Commun* 2:495–501
- Yang S, Qi Y, Liu C, Wang Y, Zhao Y, Wang L, Li J, Tan W, Yang R (2014) Design of a simultaneous target and location-activatable fluorescent probe for visualizing hydrogen sulfide in lysosomes. *Anal Chem* 86:7508–7515