## **ERRATUM**

## Erratum to: Nanomagnet-Bound Imidazole as a Heterogeneous Axial Ligand for Mn<sup>III</sup>(salophen)Cl: An Efficient, Recoverable and Recyclable Catalyst for Epoxidation of Alkenes with Sodium Periodate

Maryam Torki · Shahram Tangestaninejad · Valiollah Mirkhani · Majid Moghadam · Iraj Mohammadpoor-Baltork · Ahmad Reza Khosropour

Published online: 23 July 2013

© Springer Science+Business Media New York 2013

Erratum to: J Inorg Organomet Polym (2013) 23:923–929 DOI 10.1007/s10904-013-9865-2

The original version of this article unfortunately contained a mistake. The structures in Table 3 were missing. The corrected Table 3 is given below.

The online version of the original article can be found under doi:10.1007/s10904-013-9865-2.

M. Torki  $\cdot$  S. Tangestaninejad ( $\boxtimes$ )  $\cdot$  V. Mirkhani ( $\boxtimes$ )  $\cdot$ 

M. Moghadam (⋈) · I. Mohammadpoor-Baltork ·

A. R. Khosropour

Department of Chemistry, Catalysis Division, University of Isfahan, 81746-73441 Isfahan, Iran

e-mail: stanges@sci.ui.ac.ir

V. Mirkhani

e-mail: mirkhani@sci.ui.ac.ir

M. Moghadam

e-mail: moghadamm@sci.ui.ac.ir



Table 3 Epoxidation of alkenes with sodium periodate catalyzed by 1 in acetonitrile/water at room temperature

Entry	Alkene	Conversion (%) <sup>a,b</sup>	Epoxide selectivity (%)	Time (h)
1		97 (92)	100	1
2		100° (94)	57	3
3		90 <sup>d</sup> (86)	91	3
4		91 <sup>e</sup> (88)	87	3
5	MeO	92 (87)	94	3
6		77 (71)	100	3
7		89 (82)	100	3
8		80 (74)	100	3
9	<b>/////</b>	58 (52)	100	7

 $Reaction\ conditions:\ alkene\ (0.5\ mmol),\ NaIO_4\ (1\ mmol),\ catalyst\ (0.150\ g,\ 0.008\ mmol\ Mn)\ and\ CH_3CN/H_2O\ (7.5\ ml)$ 

<sup>&</sup>lt;sup>a</sup> GC yield based on starting alkene

<sup>&</sup>lt;sup>b</sup> The yields in the parenthesis refer to isolated yield

<sup>&</sup>lt;sup>c</sup> The by-product is allylic ketone

<sup>&</sup>lt;sup>d</sup> The by-product is benzaldehyde

<sup>&</sup>lt;sup>e</sup> The by-product is acetophenone