CORRECTION

Correction to: A 50-year sediment record of algal assemblage changes in Hanabanilla Reservoir, Cuba

Yeny Labaut · Pablo A. Macchi · Augusto A. Comas · Carmen R. Betancourt · Misael Díaz-Asencio

Published online: 31 March 2020
© Springer Nature B.V. 2020

Correction to: J Paleolimnol (2020) 63:235–250
<https://doi.org/10.1007/s10933-020-00113-5>

Unfortunately, several plots in Fig. 3 labels are missed in the original version of this article. The author has provided the figure with the labels that were missed during editing (Fig. 3).

The authors apologize for this error.

The original article can be found online at <https://doi.org/10.1007/s10933-020-00113-5>.

Y. Labaut (✉) · A. A. Comas · M. Díaz-Asencio
Centro de Estudios Ambientales de Cienfuegos, Calle 17
esq. Ave. 46 s/n, 55100 Reina, Cienfuegos, Cuba
e-mail: yenylabautcu@gmail.com;
ylabautbetancourt@unrn.edu.ar

M. Díaz-Asencio
Centro de Investigación Científica y de Educación
Superior de Ensenada, Carretera Tijuana-Ensenada 3918,
22860 Ensenada, Baja California, Mexico

Y. Labaut · P. A. Macchi · C. R. Betancourt
Universidad Nacional de Río Negro, Instituto de
Investigación en Paleobiología y Geología,
8332 General Roca, Río Negro, Argentina

Y. Labaut
Consejo Nacional de Investigaciones Científicas y
Técnicas, Instituto de Investigación en Paleobiología y
Geología, 8332 General Roca, Río Negro, Argentina

C. R. Betancourt
Universidad de Cienfuegos “Carlos Rafael Rodríguez”,
Carretera a Rodas km 3, Cienfuegos, Cuba

◀**Fig. 3** Changes in Hanabanilla Reservoir, Cuba, between 1960 and 2012. [Maximum water volume: Vmax, total phosphorus: TP, organic carbon: OC, relative abundances of C–S–R (Competitive, Stress tolerant and Ruderal strategists) functional groups, and abundances of selected algal groups in the sedimentcore]

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.