Controlling the Critical Temperature in $\text{Mg}_{1-x}\text{Al}_x\text{B}_2$

A. Bianconi · Y. Busby · M. Fratini · V. Palmisano ·
L. Simonelli · M. Filippi · S. Sanna · F. Congiu ·
A. Saccone · M. Giovannini · S. De Negri

Published online: 16 October 2007
© Springer Science+Business Media, LLC 2007

Erratum to: J Supercond Nov Magn DOI 10.1007/s10948-007-0279-7

The incorrect version of Fig. 1 was inadvertently printed.
The correct version is as follows:

The online version of the original article can be found under
doi:10.1007/s10948-007-0279-7.

A. Bianconi · Y. Busby (✉) · M. Fratini · V. Palmisano ·
L. Simonelli
Dipartimento di Fisica, Università “La Sapienza”,
P. le Aldo Moro 2, 00185 Rome, Italy
e-mail: yan.busby@roma1.infn.it

A. Bianconi
e-mail: antonio.bianconi@roma1.infn.it

M. Filippi
Laboratoire CRISMAT, UMR CNRS ISMRA 6508,
6 bd Maréchal Juin, 14050 Caen Cedex, France

S. Sanna
CNISM Dipartimento di Fisica, Università di Parma,
via G.P. Usberti 7, 43100 Parma, Italy

F. Congiu
Dipartimento di Fisica, Università di Cagliari,
S.P. Monserrato-Sestu, 09042 Monserrato (Cagliari), Italy

A. Saccone · M. Giovannini · S. De Negri
Dipartimento di Chimica e Chimica industriale, Università
di Genova, via Dodecaneso 31, 16146 Genoa, Italy

Fig. 1 SEM micrographs of $\text{Mg}_{1-x}\text{Al}_x\text{B}_2$ sample with $x = 0.28$ with two different magnification (scan dimensions are indicated by the markers on the figure)