



Correction to: Impact of environmental conditions on sorption of $^{210}\text{Pb}(\text{II})$ to NKF-5 zeolite

Donglin Zhao^{1,2} · Caicai Zhang^{2,3} · Junzheng Xu³ · Zhiwei Niu³

Published online: 4 January 2020
© Akadémiai Kiadó, Budapest, Hungary 2020

Correction to: J Radioanal Nucl Chem (2011) 289:671–678
<https://doi.org/10.1007/s10967-011-1011-4>

The original publication of the article contains error in Figs. 1 and 2. The corrected Figs. 1 and 2 are provided in this correction.





Fig. 1 FT-IR spectrum of NKF-5 zeolite

The original article can be found online at <https://doi.org/10.1007/s10967-011-1011-4>.

✉ Donglin Zhao
zhaodlin@126.com

- ¹ School of Materials and Chemical Engineering, Anhui University of Architecture, Hefei 230601, Anhui, People's Republic of China
- ² Key Laboratory of Novel Thin Film Solar Cells, Institute of Plasma Physics, Chinese Academy of Sciences, P.O. Box 1126, Hefei 230031, People's Republic of China
- ³ Radiochemistry Laboratory, Lanzhou University, Lanzhou 730000, People's Republic of China

Fig. 2 Power *XRD* pattern of NKF-5 zeolite. A: the *XRD* pattern of NKF-5 zeolite sample used in the experiment, and B: the standard *XRD* pattern in reference



Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.