RESEARCH ARTICLE

Characterising the relationship between suitable habitat and gene flow for *Myotis bechsteinii* and *Eptesicus serotinus* in Britain

P. G. R. Wright · C. Bellamy · P. B. Hamilton · H. Schofield ·
D. Finch · F. Mathews

Received: 14 October 2020 / Accepted: 3 August 2021 / Published online: 19 August 2021
© The Author(s) 2021

Abstract

Context Habitat suitability models (HSM) have been used to understand the impacts of landscape-scale habitat connectivity and gene flow mostly by assuming a regular decrease in the cost of movement as habitat improves. Yet, habitat selection and gene flow are governed by different behavioural processes which may limit the reliability of this approach as individuals are likely to disperse through unsuitable habitat for breeding.

Objectives The aim of this study was to identify the optimal relationship between gene flow and HSMs for two bat species (*Myotis bechsteinii* and *Eptesicus serotinus*) in Britain by testing a range of nonlinear negative exponential functions for the transformation of HSMs into resistance surfaces.

Methods We modelled habitat suitability using a hierarchical, multi-level approach that integrates models across three nested levels. Then, we measured the relationship between published genetics data of both species and six negative exponential transformations of the predicted outputs.

Results The two most extreme transformations provided the best fit to genetic data for both *M. bechsteinii* ($c = 32$; $R^2 = 0.87$) and *E. serotinus* ($c = 16$; $R^2 = 0.42$). The negative linear transformations had the poorest fit.

Conclusions These results suggest that bats are able to disperse through areas of poor habitat for breeding, but will avoid the most unsuitable areas. We recommend comparing multiple transformations of HSMs at different resolutions to gain a more accurate representation of gene flow across heterogeneous landscapes and to inform cost-effective, targeted management.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10980-021-01317-4>.

P. G. R. Wright · H. Schofield
Vincent Wildlife Trust, Ledbury HR8 1EP, UK

P. G. R. Wright · D. Finch · F. Mathews (✉)
University of Sussex, Brighton BN1 9RH, UK
e-mail: f.mathews@sussex.ac.uk

C. Bellamy
Centre for Ecosystems, Forest Research, Roslin, UK

P. B. Hamilton
University of Exeter, Exeter EX4 4PS, UK

Keywords Bats · Chiroptera · Landscape genetics · Habitat suitability models · Dispersal · Movement · Connectivity

Introduction

Land use change and intensification are major contributors towards the rapid and ongoing decline of

biodiversity (Ceballos et al. 2017). Their negative effects can be amplified when coupled with climatic and other environmental change leading to population declines (De Chazal and Rounsevell 2009), the loss of genetic diversity, and ultimately species loss (Bailey 2007). Understanding how landscape composition and structure influence the genetic connectivity of a species is essential for wildlife conservation. This is particularly true for rare or elusive species for which we have limited knowledge, and species more at risk of genetic drift and local extinctions due to traits such as narrow niche requirements or low dispersal abilities (e.g. Razgour et al. 2011). Habitat Suitability Modelling (HSM) is increasingly used to fill these knowledge gaps (Elith and Leathwick 2009; Bellamy et al. 2013). This approach predicts the distribution of a species and provides information on the underlying drivers using occurrence records and environmental data layers (Guisan et al. 2017; Guisan and Zimmermann 2000). As well as understanding and predicting habitat suitability, effective conservation action requires information on how landscapes influence species movements and gene flow (Razgour et al. 2016). Although major barriers, such as mountain ranges or seas, are easy to identify and detect for some species (e.g. Garroway et al. 2011; McLeod et al. 2015; Moussy et al. 2015), the heterogeneous structure and composition of many landscapes will affect population processes in a more complex way. Therefore, detailed analytical methods are needed to help understand and predict these relationships. Landscape genetic approaches have been used to assess the amount of connectivity between habitats in a fragmented landscape (e.g. Clauzel et al. 2015; Ramirez-Reyes et al. 2016); characterise features affecting genetic variation; identify isolated populations; map and design potential wildlife corridors to facilitate dispersal and gene flow (Balkenhol et al. 2016; Razgour et al. 2016) and predict the impact of landscape changes on wildlife (Razgour et al. 2017).

Predicted habitat suitability often correlates with gene flow (e.g. Razgour et al. 2017). Most studies have used negative linear transformations (resistance decreases at a constant rate as suitability increases) of habitat suitability models into resistance surfaces (the degree to which movement of an organism is affected in a grid cell; Spear et al. 2010) to assess the relationship between gene flow and habitat suitability (Pullinger and Johnson 2010; Richards-Zawacki

2009). However, Beier et al. (2009) challenged this assumption as habitat selection and dispersal/gene flow are governed by different behavioural processes. The first is mainly driven by adult home ranges, whereas gene flow is primarily influenced by the dispersal or movement of individuals and will vary depending on the species' dispersal abilities (Balkenhol et al. 2016). It has been argued that dispersing individuals will be more tolerant to 'sub-optimal' habitat than individuals foraging within their home ranges. Therefore, the creation of resistance surfaces, where the species' ability to move decreases slightly as habitat deteriorates with the exception of lower suitability values where movement is almost impossible, is likely to better reflect dispersal and gene flow. Several studies applying negative exponential functions to transform HSMs into resistance surfaces (Fig. 1) to test this hypothesis have proven successful for different species (Trainor et al. 2013; Keeley et al. 2016; Zeller et al. 2018). Yet, these are still not commonly applied when deriving HSMs into resistance surfaces.

Bats, being small and volant, are highly mobile and able to travel long distances within a short period of time in comparison to other terrestrial mammals (Ahlén et al. 2009; Bryja et al. 2009). Behaviours, such as autumnal swarming, where individual bats travel significant distances to aggregate at

Fig. 1 Six curves used to transform habitat suitability values into resistance values. The curves are based on the transformation function where higher transformation values result into more habitat being classified with low resistance values

underground sites are also known to be linked with mating and facilitate gene flow (Kerth et al. 2003; Rivers et al. 2005). Such behaviours can be observed in highly specialist and sedentary species, such as *Myotis bechsteinii* (Dekeukeleire et al. 2016). Therefore, the transformation of HSMs into resistance surfaces by applying a negative exponential function, where only very poor habitat will act as a significant barrier for movement, as opposed to the more commonly negative linear transformation is likely to better predict landscape-scale genetic connectivity for most bat species.

The aim of this study is to identify the optimal relationship between gene flow and HSMs for two bat species exhibiting different reproductive and foraging strategies in Britain by testing a range of nonlinear negative exponential functions. We tested the functions on *Myotis bechsteinii*, a woodland specialist that performs autumnal swarming and *Eptesicus serotinus*, a more generalist species that can forage and roost in built-up areas but is not known to perform autumnal swarming. We hypothesised that higher transformations of HSMs would better reflect the relationship between suitable habitat and gene flow.

Materials and methods

Habitat suitability models

We used the hierarchical, multi-level HSM framework developed by Bellamy et al. (2020) to model habitat suitability in Britain for *M. bechsteinii*, a woodland specialist that performs autumnal swarming, and *E. serotinus*, a more generalist species that is not known to occur at swarming sites. This method nests models across three levels (population range, home range, local habitat —See below) to limit predictor collinearity and enable context dependency. MaxEnt (Phillips et al. 2006), a presence-only HSM approach, was used to predict suitability at each level. All analyses were carried out using R (v. 3.4.2; R Core Team 2018) in R Studio (v.1.1.383; RStudio Team).

We used landscape variables reflecting environmental features and conditions likely to influence the distribution of temperate bats (See supplementary 1 for predictor variables; Boughey et al. 2011; Bellamy et al. 2013; Fuller et al. 2018; Rebelo et al. 2010). We performed an initial population range model focusing

on climate at a 5 km resolution. This was followed by a summer home range (1 km resolution), and a local habitat (100 m resolution) model. The local habitat model was restricted geographically to southern Britain (i.e. were within both species' known current range), and employed land cover, landscape structure and land management variables.

We collected presence records between 2000 and 2016 during the months of April to September at a ≤ 10 km resolution which were extracted from the National Biodiversity Network (NBN; <https://nbnatlas.org/>) gateway, national and local bat monitoring schemes, and local record centres. For *M. bechsteinii*, we only used confirmed roosts and bats identified in hand, because the species' echolocation calls overlaps with that of other *Myotis* species making the use of acoustic data unreliable. The lack of information in the reporting of *E. serotinus* presence data throughout Britain meant that we also included confirmed acoustic records, as the species can be differentiated from others, along with roosts and bats identified in the hand. We filtered records to match the resolution of each model (i.e. 5 km, 1 km, and 100 m respectively) and to retain a single record per grid square. To reduce the amount of sampling bias towards heavily sampled areas (Graham et al. 2015), we used 10,000 pseudoabsence records of all bat species within the study extent from the same sources recorded over the same time period (Phillips et al. 2009). It should be noted that this approach does not entirely control sampling bias for *M. bechsteinii* as most bat records originate from acoustic data and *M. bechsteinii* can only be reliably identified in hand.

The population range model was run first, and the output Habitat Suitability Index (HSI) values from each of the three levels were disaggregated to the resolution of the next lower level and incorporated as an explanatory variable. For each level, we removed highly correlated variables using the 'vifstep' stepwise function of the 'usdm' package (Naimi et al. 2014) and a conservative VIF threshold of three (Zuur et al. 2010). The package 'ENMeval' (Muscarella et al. 2014) was used to identify the optimal MaxEnt model settings. We tested combinations of feature types (L, linear; H, hinge; Q, quadratic; P, product) and disabled threshold features to reduce overfitting. We varied the regularisation multiplier in steps of 0.5, from 0.5 to 4. A final model was then created using the optimal settings to produce model predictions.

Landscape genetics

Myotis bechsteinii and *Eptesicus serotinus* genotypes were taken from previously published papers (Moussy et al. 2015; Wright et al. 2018), but excluded samples collected outside Britain. The *M. bechsteinii* dataset included genotypes from 260 individuals using 14 microsatellite loci collected at 8 sites. The *E. serotinus* dataset comprised 593 samples using 10 loci from 28 sites. We used the genetic landscape-shape analysis from the Alleles in Space software (Miller 2005) to identify patterns of genetic structure throughout the study areas (Moussy et al. 2015; Wright et al. 2018). The connectivity networks produced from this software, were based on Delaunay triangulations (Watson 1992; Brouns et al. 2003) and were used to build a three-dimensional map, where the Z-axis corresponded to genetic distance between populations.

To transform HSI values into resistance surfaces, we used a negative linear transformation and six transformation values on a negative exponential function (Fig. 1) as reported by Trainor et al. (2013), Mateo-Sánchez et al. (2015) and Keeley et al. (2016):

$$R = 100 - 99(1 - \exp(-cH))/(1 - \exp(-c))$$

H = habitat suitability value, $c = 1, 2, 4, 8, 16$ and 32 .

Finally, because woodland connectivity is frequently used as an indicator of habitat connectivity for bats, we also created a woodland cover resistance surface where the highest resistance values (100) were associated with the lowest amount of woodland.

We used Circuitscape v4.0.5 (McRae and Nurnberger 2006) to calculate resistance distance matrices between study populations (focal nodes) using the “pairwise” modelling mode. To test the relationship between the transformed HSM variables and gene flow for *M. bechsteinii* and *E. serotinus* in Britain, we performed multiple regressions on distance matrices (MRDMs) using 10,000 permutations (Legendre et al. 1994; Goslee and Urban 2007). We performed MRDMs on the output matrices from Circuitscape, along with genetic and geographic distance matrices, as described by Santos (2017). Distance matrices were initially standardized from 0 to 1 using the ‘scales’ package (Wickham 2014). The effect of geographic distance was then removed by extracting the residuals of linear models of geographic distance against each resistance surface (Dyer et al. 2010). Univariate MRDMs of genetic distance were finally calculated

for each resistance surface, in order to assess the correlation of different HSM transformations with gene flow. MRDMs were performed for all transformations for which the sea was assigned a resistance value of 70 and 120 to identify the optimal effect of the sea separating the Isle of Wight (an island on the coast of southern England) from mainland Britain on gene flow.

Results

Habitat suitability models

A total of 283 *M. bechsteinii* and 3,460 *E. serotinus* records across Britain were used for the final local model (Table 1). The multi-scale frameworks performed well on both *M. bechsteinii* and *E. serotinus* with good AUC scores (*M. bechsteinii*: $AUC \geq 0.79$; *E. serotinus*: $AUC \geq 0.74$). Overall, the framework performed better for *M. bechsteinii*; AUC values increased to 0.89 (local habitat). The highest AUC for *E. serotinus* was 0.81 (local habitat).

For both species, suitable conditions predicted from the population range model were concentrated in southern Britain and were mainly influenced by temperature, precipitation and decreased woodland cover (Supplementary Information 2–8 for permutation importance and 3–9 for response curves). At the home range and local levels, disaggregated HSI values from preceding levels were the strongest explanatory variables with permutation importance’s ranging from 76 to 86%. In the home range level model, high cover of both protected broadleaf and ancient woodland were associated with the presence of *M. bechsteinii*; whilst no clear patterns were identified with *E. serotinus* outside the importance of the preceding population range level (Supplementary Information 4–10 for permutation importance and 5–11 for response curves). Woodlands were also associated with *M. bechsteinii* at the local level as the species was primarily influenced by woodland patch shape and broadleaved woodland cover. Finally, *E. serotinus* habitat showed strong associations to areas close to buildings and hedgerows within a 100 m scale (Fig. 2a and b; Supplementary Information 6–12 for permutation importance and 7–13 for response curves).

Table 1 Model performance of the final HSM at a population range (5 km), home range (1 km) and local range (100 m) for both *Myotis bechsteinii* (*M. b.*) and *Eptesicus serotinus* (*E. s.*) in Britain

Species	Model type	n	Feature types	Regularisation multiplier	AIC _c	Non-independent AUC	Mean AUC $\pm$ var
<i>M. b.</i>	Population range	109	LQ	0.5	1700.3	0.79	0.78 $\pm$ 0.03
	Home range	183	LQ	0.5	3751.9	0.88	0.82 $\pm$ 0.002
	Local habitat	283	LQ	1	8080.7	0.89	0.88 $\pm$ 0.0003
<i>E. s.</i>	Population range	1337	LQHPT	2.5	22,027.7	0.74	0.66 $\pm$ 0.007
	Home range	2494	LQHPT	2.5	54,666.9	0.81	0.74 $\pm$ 0.002
	Local habitat	3460	LQHPT	2	105,076.8	0.81	0.75 $\pm$ 0.002

Fig. 2 Predicted logistic habitat suitability indices (HSI) of the final local level model (100 m resolution) for (a.) *M. bechsteinii* and (b.) *E. serotinus*; and the resistance surface with the highest correlation with gene flow for (c.) *M. bechsteinii* and (d.) *E. serotinus*

Landscape genetics

The genetic landscape-shape analysis revealed contrasting population structure between *M. bechsteinii* and *E. serotinus*. While only two *M. bechsteinii* populations showed signs of isolation, many *E.*

serotinus populations were estimated to be isolated (Fig. 3).

For *M. bechsteinii*, the most extreme transformation of the local HSM output ($c = 32$) showed the highest correlation coefficient ($R^2 = 0.87$, $p < 0.001$), while the second highest transformation had the

Fig. 3 Predicted genetic distances of location of sampling sites of a. *Myotis bechsteinii* and b. *Eptesicus serotinus*. Peaks (red colour) represent areas of high pairwise genetic distance,

indicating isolated populations, and valleys (blue colour) are indicative of areas of low pairwise genetic distance

highest correlation coefficient ($c = 16$) for *E. serotinus* ($R^2 = 0.42$, $p < 0.001$; Table 2). The correlation coefficients did not all increase with increased transformation levels. In most models, coefficients were highest at $c = 16$ and, in the case of *M. bechsteinii*, these notably declined at $c = 32$ in the population and home range models (Table 2). The models also revealed that woodland cover did not correlate with gene flow for both species ($R^2 < 0.05$ and $p > 0.5$ for both models).

Discussion

Maintaining gene flow between populations is crucial for securing a species' viability. However, understanding how small, mobile species, such as bats, move through the landscape in order to maintain gene

flow is hard to measure directly. In this study, we assessed the strength of association between a range of nonlinear negative exponential functions derived from HSMs to describe the relationship between predicted suitable habitat and gene flow of two bat species (*Myotis bechsteinii* and *Eptesicus serotinus*) in Britain. We found that the two highest transformations ($c = 32$ and $c = 16$) of HSMs into resistance surfaces had the strongest relationship with gene flow for both species.

Negative linear transformations of HSMs into resistance surfaces are frequently used for assessing the degree of permeability of different habitats to species, yet their sole use would have underestimated the ability for both species to maintain genetic connectivity in a fragmented landscape. Our results suggest that the design of corridors and other management efforts to maintain gene flow between

Table 2 Summary of MRDM R^2 for *M. bechsteinii* and *E. serotinus* (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$)

HSM transformation ^a	Negative Linear		1		2		4		8		16		32	
	70	120	70	120	70	120	70	120	70	120	70	120	70	120
<i>M. bechsteinii</i>														
Population Range	0.58**	0.59**	0.62**	0.65**	0.66**	0.68**	0.71**	0.71**	0.77**	0.77**	0.45*	0.43*	0	0
Home Range	0.33	0.39*	0.4*	0.47*	0.48*	0.55*	0.63**	0.68**	0.82**	0.85***	0.85***	0.74***	0.18	0.09
Local Habitat	0.02	0.05	0.05	0.11	0.1	0.62**	0.25	0.35	0.53*	0.62**	0.76***	0.80***	0.87***	0.71***
<i>E. serotinus</i>														
Population Range	0.14**	0.20**	0.17***	0.24***	0.20***	0.26***	0.25***	0.28***	0.30***	0.31***	0.22	0.22***	0.20***	0.20***
Home Range	0.09	0.17**	0.11*	0.21***	0.13*	0.23***	0.20*	0.29***	0.35***	0.40***	0.40***	0.40***	0.38***	0.38***
Local Habitat	0.06	0.14**	0.07	0.17***	0.09	0.19**	0.14**	0.23**	0.28***	0.36***	0.42***	0.42***	0.41***	0.41***

^aHSM transformation values represent the different curves in Fig. 1 and the highest correlations appear in bold

populations of both species require a high negative exponential transformation of habitat suitability models. These findings also mirrored the few studies having previously assessed the use of such transformations on other species (Trainor et al. 2013; Mateo-Sánchez et al. 2015; Keeley et al. 2016; Zeller et al. 2018). These results also have important implications for bat conservation as they could provide new insight into the ability of populations to shift their distribution under climate change (Razgour et al. 2017).

The hierarchical framework has been found to improve the accuracy of HSMs by integrating predictors at their scales of effect (Bellamy et al. 2020). For landscape genetics, the framework also offers insight into the models that better represent gene flow, albeit we can only make assumptions on the reasons why certain models show higher correlations (e.g. higher resolution, higher AUC values, different environmental variables). Interestingly, the two highest transformations of the local level HSM model (highest resolution) were found to best explain gene flow. However, for *M. bechsteinii*, the correlations for the highest transformations for the population and home range models dropped significantly once the highest correlation was reached at transformation 16. This suggests that a wide range of transformations of different resolutions must be tested as correlations do not always increase along with HSM transformations.

The highest correlation level for *M. bechsteinii* was approximately twice as high as for *E. serotinus*. This could be explained by the higher performance of HSMs, which may be linked to the fact that the ecological niche of habitat specialists, such as *M. bechsteinii*, tends to be better predicted (Evangelista et al. 2008). *Myotis bechsteinii* was also associated with the highest transformation as opposed to the second highest for *E. serotinus* meaning that resistance values were higher throughout the landscape for *E. serotinus*. Some bat species, such as *M. bechsteinii*, are known to perform autumnal swarming which is linked with mating. While *M. bechsteinii* tend to forage over short distances in old growth woodlands (~ 3 km, Schofield and Morris 2000), they will readily travel distances of up to 20 km to reach swarming sites (Dekeukeleire et al. 2016). In a fragmented landscape, as found in Britain, such movements would inevitably entail crossing some unsuitable habitat while the most unsuitable habitat will act as a barrier. The effective dispersal

mechanism of *M. bechsteinii* has likely reduced the impact of habitat fragmentation. However, the isolation of certain populations raises questions on a possible dispersal threshold for the species. In other words, it is important to identify how much unsuitable habitat individuals are able to traverse before populations become genetically isolated.

As for *E. serotinus*, very little is known about their mating system and dispersal abilities. The species ranges over significantly larger areas (Catto et al. 1996) and will forage and roost in a broader range of habitats, including built-up areas, making it possibly less sensitive to human development than *M. bechsteinii*. However, it does not exhibit swarming behaviour like *Myotis* bats. Limited dispersal for mating could result in the greater genetic differentiation between populations observed in Britain as the species may be less likely to travel long distances and move through unsuitable habitat for mating. This could explain the stronger association with the second highest transformation ($c = 16$) as opposed to the highest for *M. bechsteinii* ($c = 32$). Differences between the dispersal abilities of both species can be observed on the Isle of Wight (UK) located 1–5 km from the south coast of England. Here, genetic structure between the Isle of Wight and Britain for *E. serotinus* has been identified (Moussy et al. 2015), while no apparent genetic structure was observed for *M. bechsteinii* (Wright et al. 2018). This reinforces the proposal that gene flow in *M. bechsteinii* is less restricted by the presence of unsuitable habitat than in the case of *E. serotinus*.

Whilst still being rarely performed the use of negative exponential transformations of HSMs tend to provide better insight into the dispersal mechanisms of wildlife (Trainor et al. 2013; Keeley et al. 2016; Zeller et al. 2018) and must continue to be encouraged. These may be particularly relevant for the design of wildlife corridors for species willing to travel through large areas of unsuitable habitat to establish new territories and breed. For example, terrestrial carnivores, such as *Canis lupus* or *Puma concolor* (Zeller et al. 2018), may disperse over hundreds of kilometres and will inevitably use unsuitable habitat. The added information from these models will help identify the most likely paths animals will use and where small improvements in habitat can be made to encourage the successful dispersal of a species as opposed to its establishment.

Conclusions

Our aim was to identify the optimal relationship between gene flow and HSMs for two bat species, *M. bechsteinii* and *E. serotinus*. For this we tested multiple negative exponential transformations of HSMs into resistance surfaces as opposed to the more commonly used single negative linear transformation. Our study highlights the importance of testing a multitude of transformations as the highest correlations for both species was identified from the two highest transformations ($c = 16$ and 32). These findings have important implications for bat conservation as the sole use of negative linear transformations would have underestimated the ability of both species to maintain gene flow between populations. Bats exhibit complex mating behaviour and may travel significant distances outside of high quality habitat for mating. It is therefore important that such transformations are applied on HSMs for landscape genetic studies as they may provide a more accurate representation of dispersal in a heterogeneous landscape.

Acknowledgements This work was supported by the Woodland Trust, the People's Trust for Endangered Species and the Vincent Wildlife Trust. Fiona Mathews is supported by a NERC KE Fellowship NE/S006486/1 and the University of Sussex. We would also like to thank Caroline Moussy for making her dataset publicly available.

Author contributions The study was designed by PW, FM, HS, PH and CB. All data were analysed by PW with the help of CB with the preparation of GIS data. Manuscript preparation was carried out by PW and DF. FM, HS, PH, CB & DF provided comments and reviewed the manuscript.

Data availability Data used for this study was obtained from previously published resources (Moussy et al. 2015; Wright et al. 2018). Code and map outputs have been made available on figshare (<https://doi.org/10.6084/m9.figshare.15036192>).

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not

included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ahlén I, Baagøe HJ, Bach L (2009) Behavior of Scandinavian bats during migration and foraging at sea. *J Mammal* 90(6):1318–1323
- Bailey S (2007) Increasing connectivity in fragmented landscapes: an investigation of evidence for biodiversity gain in woodlands. *For Ecol Manage* 238(1):7–23
- Balkenhol N, Cushman SA, Storfer A, Waits LP (2016) Introduction to landscape genetics-concepts, methods, applications. Wiley-Blackwell, Chichester
- Beier P, Majka DR, Newell SL (2009) Uncertainty analysis of least-cost modeling for designing wildlife linkages. *Ecol Appl* 19(8):2067–2077
- Bellamy C, Scott C, Altringham J (2013) Multiscale, presence-only habitat suitability models: fine-resolution maps for eight bat species. *J Appl Ecol* 50(4):892–901
- Bellamy C, Boughhey K, Hawkins C, Reveley S, Spake R, Williams C, Altringham J (2020) A sequential multi-level framework to improve habitat suitability modelling. *Landsc Ecol* 35:1–20
- Boughhey KL, Lake IR, Haysom KA, Dolman PM (2011) Effects of landscape-scale broadleaved woodland configuration and extent on roost location for six bat species across the UK. *Biol Conserv* 144(9):2300–2310
- Brouns G, De Wulf A, Constaes D (2003) Delaunay triangulation algorithms useful for multibeam echosounding. *J Surv Eng* 129(2):79–84
- Bryja J, Kaňuch P, Fornůsková A, Bartonička T, Řehák Z (2009) Low population genetic structuring of two cryptic bat species suggests their migratory behaviour in continental Europe. *Biol J Linn Soc* 96(1):103–114
- Catto CMC, Hutson AM, Racey PA, Stephenson PJ (1996) Foraging behaviour and habitat use of the serotine bat (*Eptesicus serotinus*) in southern England. *J Zool* 238:623–633
- Ceballos G, Ehrlich PR, Dirzo R (2017) Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proc Natl Acad Sci USA* 114:E6089–E6096
- Clauzel C, Bannwarth C, Foltete J-C (2015) Integrating regional-scale connectivity in habitat restoration: An application for amphibian conservation in eastern France. *J Nat Conserv* 23:98–107
- De Chazal J, Rounsevell MD (2009) Land-use and climate change within assessments of biodiversity change: a review. *Glob Environ Chang* 19(2):306–315
- Dekeukeleire D, Janssen R, Haarsma A-J, Bosch T, Schaik JV (2016) Swarming behaviour, catchment area and seasonal movement patterns of the Bechstein's bats: implications for conservation. *Acta Chiropterologica* 18(2):349–358
- Dyer RJ, Nason JD, Garrick RC (2010) Landscape modelling of gene flow: improved power using conditional genetic distance derived from the topology of population networks. *Mol Ecol* 19(17):3746–3759
- Elith J, Leathwick JR (2009) Species distribution models: ecological explanation and prediction across space and time. *Annu Rev Ecol Evol Syst* 40:677–697
- Evangelista PH, Kumar S, Stohlgren TJ, Jarnevich CS, Crall AW, Norman JB III, Barnett DT (2008) Modelling invasion for a habitat generalist and a specialist plant species. *Divers Distrib* 14(5):808–817
- Fuller L, Shewring M, Caryl FM (2018) A novel method for targeting survey effort to identify new bat roosts using habitat suitability modelling. *Eur J Wildl Res* 64(3):31
- Garraway CJ, Bowman J, Wilson PJ (2011) Using a genetic network to parameterize a landscape resistance surface for fishers *Martes Pennanti*. *Mol Ecol* 20(19):3978–3988
- Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological data. *J Stat Softw* 22(7):1–19
- Graham LJ, Haines-Young RH, Field R (2015) Using citizen science data for conservation planning: methods for quality control and downscaling for use in stochastic patch occupancy modelling. *Biol Conserv* 192:65–73
- Guisan A, Zimmermann NE (2000) Predictive habitat distribution models in ecology. *Ecol Model* 135(2):147–186
- Guisan A, Thuiller W, Zimmermann NE (2017) Habitat suitability and distribution models: with applications in R. Cambridge University Press, Cambridge
- Keeley AT, Beier P, Gagnon JW (2016) Estimating landscape resistance from habitat suitability: effects of data source and nonlinearities. *Landsc Ecol* 31(9):2151–2162
- Kerth G, Kiefer A, Trappmann C, Weishaar M (2003) High gene diversity at swarming sites suggest hot spots for gene flow in the endangered Bechstein's bat. *Conserv Genet* 4(4):491–499. <https://doi.org/10.1023/a:1024771713152>
- Legendre P, Lapointe FJ, Casgrain P (1994) Modeling brain evolution from behavior: a permutational regression approach. *Evolution* 48(5):1487–1499
- Mateo-Sánchez MC, Balkenhol N, Cushman S, Pérez T, Domínguez A, Saura S (2015) A comparative framework to infer landscape effects on population genetic structure: are habitat suitability models effective in explaining gene flow? *Landsc Ecol* 30(8):1405–1420
- McLeod B, Burns L, Frasier T, Broders H (2015) Effect of oceanic straits on gene flow in the recently endangered little brown bat (*Myotis lucifugus*) in maritime Canada: implications for the spread of white-nose syndrome. *Can J Zool* 93(6):427–437
- McRae BH, Nünberger B (2006) Isolation by resistance. *Evolution* 60(8):1551–1561
- Miller M (2005) Alleles In Space (AIS): computer software for the joint analysis of interindividual spatial and genetic information. *J Hered* 96(6):722–724
- Moussy C, Atterby H, Griffiths A, Allnutt T, Mathews F, Smith G, Hosken D (2015) Population genetic structure of serotine bats (*Eptesicus serotinus*) across Europe and implications for the potential spread of bat rabies (European bat lyssavirus EBLV-1). *Heredity* 115(1):83–92
- Muscarella R, Galante PJ, Soley-Guardia M, Borja RA, Kass JM, Uriarte M, Anderson RP (2014) ENM eval: An R

- package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models. *Methods Ecol Evol* 5(11):1198–1205
- Naimi B, Hamm NA, Groen TA, Skidmore AK, Toxopeus AG (2014) Where is positional uncertainty a problem for species distribution modelling? *Ecography* 37(2):191–203
- Phillips S, Anderson RP, Schapire R (2006) Maximum entropy modeling of species geographic distributions. *Ecol Model* 190:231–259
- Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecol Appl* 19(1):181–197
- Pullinger MG, Johnson CJ (2010) Maintaining or restoring connectivity of modified landscapes: evaluating the least-cost path model with multiple sources of ecological information. *Landsc Ecol* 25(10):1547–1560
- R Core Team (2018) R: a language and environment for statistical computing. The R foundation for Statistical Computing, Vienna
- Ramirez-Reyes C, Bateman BL, Radeloff VC (2016) Effects of habitat suitability and minimum patch size thresholds on the assessment of landscape connectivity for jaguars in the Sierra Gorda, Mexico. *Biol Conserv* 204:296–305
- Razgour O, Hanmer J, Jones G (2011) Using multi-scale modelling to predict habitat suitability for species of conservation concern: the grey long-eared bat as a case study. *Biol Conserv* 144(12):2922–2930
- Razgour O, Rebelo H, Di Febbraro M, Russo D (2016) Painting maps with bats: species distribution modelling in bat research and conservation. *Hystrix* 27:1–8
- Razgour O, Taggart JB, Manel S, Juste J, Ibáñez C, Rebelo H, Park K (2017) An integrated framework to identify wildlife populations under threat from climate change. *Mol Ecol Resour* 18(1):18–31
- Rebelo H, Tarroso P, Jones G (2010) Predicted impact of climate change on European bats in relation to their biogeographic patterns. *Glob Change Biol* 16(2):561–576
- Richards-Zawacki CL (2009) Effects of slope and riparian habitat connectivity on gene flow in an endangered Panamanian frog *Atelopus varius*. *Divers Distrib* 15(5):796–806
- Rivers NM, Butlin RK, Altringham JD (2005) Genetic population structure of Natterer's bats explained by mating at swarming sites and philopatry. *Mol Ecol* 14(14):4299–4312
- RStudio Team (2018) RStudio: integrated development environment for R. Boston, RStudio
- Santos H (2017) Combining molecular analyses with predictive modelling to study the population history of Iberian cryptic bats. (Doctoral thesis). Universidade do Porto, Porto
- Schofield H, Morris C (2000) Ranging behaviour and habitat preferences of female Bechstein's bat, *Myotis bechsteinii* (Kuhl, 1818), in summer: Report held by The Vincent Wildlife Trust
- Spear SF, Balkenhol N, Fortin MJ, McRae BH, Scribner K (2010) Use of resistance surfaces for landscape genetic studies: considerations for parameterization and analysis. *Mol Ecol* 19(17):3576–3591
- Trainor AM, Walters JR, Morris WF, Sexton J, Moody A (2013) Empirical estimation of dispersal resistance surfaces: a case study with red-cockaded woodpeckers. *Landsc Ecol* 28(4):755–767
- Watson DF (1992) Contouring: a guide to the analysis and display of spatial data. Pergamon Press, New York
- Wickham H (2014) Package 'scales. Scale functions for graphics. <http://cran.rproject.org/web/packages/scales/scales.pdf>
- Wright P, Hamilton P, Schofield H, Damant C, Davidson-Watts I, Glover AM, Mathews F (2018) Genetic structure and diversity of a rare woodland bat, *Myotis bechsteinii*: comparison of continental Europe and Britain. *Conserv Genet* 19(4):777–787
- Zeller KA, Jennings MK, Vickers TW, Ernest HB, Cushman SA, Boyce WM (2018) Are all data types and connectivity models created equal? Validating common connectivity approaches with dispersal data. *Divers Distrib* 24(7):868–879
- Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical problems. *Methods Ecol Evol* 1(1):3–14

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.