



Meaning-Preserving Translations of Non-classical Logics into Classical Logic: Between Pluralism and Monism

Gerhard Schurz¹ 

Received: 3 October 2020 / Accepted: 31 May 2021 / Published online: 6 September 2021

© The Author(s) 2021

Abstract

In order to prove the validity of logical rules, one has to assume these rules in the metalogic. However, rule-circular ‘justifications’ are demonstrably without epistemic value (sec. 1). Is a non-circular justification of a logical system possible? This question attains particular importance in view of lasting controversies about classical versus non-classical logics. In this paper the question is answered positively, based on meaning-preserving translations between logical systems. It is demonstrated that major systems of non-classical logic, including multi-valued, paraconsistent, intuitionistic and quantum logics, can be translated into classical logic by introducing additional intensional operators into the language (sec. 2–5). Based on this result it is argued that classical logic is representationally optimal. In sec. 6 it is investigated whether non-classical logics can be likewise representationally optimal. The answer is predominantly negative but partially positive. Nevertheless the situation is not symmetric, because classical logic has important *ceteris paribus* advantages as a unifying metalogic.

Keywords Translations of non-classical into classical logics · Meaning-preserving translations · Representational optimality, logical pluralism

1 Introduction: The Significance of Non-circular Justifications for Contemporary Philosophy of Logic

The background of this paper is a foundation-theoretic epistemology whose class of ‘basic’ beliefs is *minimalistic*, consisting only of immediately evident *analytical* and *introspective* beliefs (cf. [59, 60]). In this framework the major epistemic load has to be carried by deductive, inductive or abductive reasoning.

✉ Gerhard Schurz
schurz@hhu.de; <http://www.philosophie.hhu.de/schurz>

¹ Düsseldorf Center for Logic and Philosophy of Science (DCLPS), Heinrich Heine University Düsseldorf, Universitätsstraße 1, Geb. 24.52, 40225 Düsseldorf, Germany

Therefore the ‘higher order’ justification of the reasonability or reliability of the rules governing these inferences acquires central importance. Many contemporary epistemologists consider the problem of higher order justification as unsolvable, because it leads into a circle or infinite regress. In epistemological movements such as coherentism and (partly) externalism it is argued that *rule-circular* justifications, in which the rule to be justified is employed in the justificatory argument, are a viable solution to this problem.¹ However, it can be demonstrated that rule-circular justifications are epistemically worthless, because with such a justification both a rule and the opposite rule can be proved. For example, both the rule of induction and the rule of counter-induction (that infers the opposite of what induction infers) can be rule-circularly ‘justified’ as follows ([56], 46):

<p>(1) <i>Rule-circular justification of induction:</i> <i>Premise:</i> Past inductions have been successful. <u><i>Therefore, by rule of induction:</i></u> Future inductions will be successful.</p>	<p><i>Rule-circular justification of counter-induction:</i> <i>Premise:</i> Past counter-inductions have not been successful. <u><i>Therefore, by rule of counter-induction:</i></u> Future counter-inductions will be successful.</p>
---	---

A similar situation arises when the semantic validity of logical rules is proved by employing these rules in the *metalogic*. To substantiate this claim let us introduce a bit of notation: In what follows, the indexed letter L_i varies over systems of propositional object logics and \mathcal{L}_i designates their language (identified with the set of their well-formed formulas). ML_i denotes a metalogic in which we can express truth assertions about L_i -sentences. It is sufficient for our purpose to represent ML_i as a modal extension of L_i , obtained by adding an intensional truth operator T (a generalized modal operator) to L_i for which the correspondence (or ‘redundancy’) axiom (C): $TA \leftrightarrow A$ holds.² L_2 denotes the classical (bivalent) propositional logic, with \mathcal{L}_2 containing \neg, \wedge, \vee as Boolean connectives and $\rightarrow, \leftrightarrow$ for the material implication and equivalence. $\Gamma \models_{L_2} A$ stands for the corresponding inference relation (defined via truth preservation). Moreover, we use $p_1, p_2, \dots, q, r, \dots$ as (propositional) variables, A, B, \dots as schematic letters for arbitrary formulas (sentences); Γ, Δ, \dots designate arbitrary sets of formulas.

Already Haack [28] pointed out in her example of “Modus morons” that with help of rule-circular arguments rather weird rules could be ‘proved’ as valid. We illustrate this point by the following example of two circular metalogical ‘proofs’ differing only in their final lines: the left one ‘proving’ the validity of Modus Ponens, MP, and the right one ‘proving’ the validity of the opposite rule, called *Modus Nonsense*, MN: “ $p, p \rightarrow q / \neg q$ ” (truth tables are assumed to be the classical ones):

¹ Among others, the following philosophers supported rule-circular justifications. *Coherentists*: Sellars [62], Lehrer [41] and the early Bonjour [7]. *Externalists*: Goldman [25], van Cleve [68] and Greco [27]. *Philosophers of science*: Papineau [48], Lipton [43] and Psillos [53].

² For truth as a sentence operator the axiom (C) cannot produce antinomies. The modal logic for (C) is semantically characterized by the class of worlds seeing exactly themselves.

(2) Rule-circular ‘proofs’ of the validity of Modus ponens and Modus nonsense:

	1. $T(p \rightarrow q)$	Premise
	2. Tp	Premise
	3. $Tp \rightarrow Tq$	From 1 by \rightarrow 's truth table
4. Tq	From 2, 3 and MP	4*. $\neg Tq$
		From 2, 3 and MN
		5*. $T\neg q$
		From 4* by \neg 's truth table

In conclusion, rule-circular ‘justifications’ may be semantically illuminating, but they do not provide real justification. Rather, the validity or invalidity of these rules is assumed as *basic* ‘knowledge’ that is simply accepted and has no further positive justification.³ Technically this basic knowledge can be explicated either semantically, in propositional logic by means of truth tables, or syntactically by means of an axiomatic system. In our example, we ‘know’ that only step 4 but not step 4* is correct, because only the rule MP but not the rule MN is validated by the classical truth tables, that are assumed as basic.

If systems of logic cannot be justified by means of logical demonstration, what other possibilities of justifying them are there? Prima facie, the purpose of logics is to explicate valid inference (cf. Beall and Restall’s Generalized Tarski Thesis; [4], 29), but the rules of valid inference depend on the basic logical operators. To a certain extent, the basic logical operators can be justified in a Kantian ‘transcendental’ sense, as a presupposition of the *possibility of cognition at all*. For example, the possibility of a true description of a *manifold* presupposes the operation of *conjunction*; expressing that a certain description is false presupposes the operation of *negation*; and similar arguments can be given for disjunction and implication. However, these a priori reasons do *not* determine the precise logical meaning of these operators; they constitute necessary conditions, but are too weak for a justification of a system of logic. The so-called *classical* logic is characterized by an additional semantic principle, the principle of *bivalence*: every statement p that is expressed in a semantically complete (non-indexical) way is *either true or false*. Thus, p cannot be ‘undetermined’, i.e. neither true nor false (the principle of excluded middle), nor can p be ‘dialethic’, i.e. both true and false (the principle of non-contradiction). Note that the principle of bivalence is not meant in an epistemic but in an *ontological* way. Ultimately this principle expresses the *determinateness* of reality: if “ p ” is a semantically complete (non-indexical) sentence, then either p or not p must obtain.

Most non-classical logics reject the principle of two-valuedness, but in different ways and for different reasons. For example, Lukasiewicz’ *three-valued logic* [44] originated from the thesis that there are sentences whose truth-value is objectively undetermined, with the consequence that the classical law of the excluded middle ($p \vee \neg p$) becomes invalid. Brouwer’s *intuitionistic* logic [9] originated in the domain of mathematics and consisted in the replacement of the Platonist notion of mathematical truth by the concept of constructive mathematical verification, with the consequence

³ In the sense of an inference from further reasons. Weaker forms of justifications for logical principles are possible, for example epistemic entitlement in the sense of Wright [74]. My argument from representational optimality developed in this paper is itself a weaker form of (internalist) justification.

that for intuitionistic logic the classical law of double negation ($\neg\neg p \rightarrow p$) becomes invalid. *Paraconsistent* logic was motivated by the attempt to avoid the irrelevant inference of ‘explosion’ ($p \wedge \neg p / q$) by assuming the possibility of *dialethic* statements that are both true and false [51]. Still different is the motivation of *quantum logic*, attempting to represent the possibility of superposed quantum states by a non-classical disjunction that violates the expansion direction of distribution, $p \wedge (q \vee r) \rightarrow (p \wedge q) \vee (p \wedge r)$. For all these non-classical logics there have been lasting philosophical controversies about whether their objectives can be equally or even better realized by measures within classical logic. Let us avoid taking sides in these controversies, but consider the epistemological consequences of this situation. For a foundation-theoretic epistemology aiming at objective justification, these consequences are highly problematic. The key problem is this: Assessing the adequacy and the relative merits of a logic requires external standards, but apparently there are no such external standards, since every rational argument and even every description of evidence presupposes some kind of logic.

In contemporary philosophy of logic the view of logical *anti-exceptionalism* has been developed, as an opposition to the traditional logical apriorism. Defenders of this view (e.g., [10, 32]) argue that logics are not ‘exceptional’ compared to the empirical sciences. More precisely, they argue that (a) it is false that the laws of logic have an exceptional *apriori* status, as it is traditionally assumed (for contemporary defenses of the apriori view cf. [17, 20]). Rather (b) the laws of logic can be supported and revised according to empirical facts, similarly as has been the case for physical geometries [54]. In our view, thesis (a) is right, but thesis (b) is not tenable. The situation in logic is very different from the case of geometry. The justification of different systems of *physical* geometry (say Euclidean versus non-Euclidean) is based on independent systems of logico-mathematical description that do *not* presuppose a particular geometry. But every description of evidence for or against a given logic L presupposes a metalogic for describing this evidence and drawing conclusions from it about the status of L (cf. [63, 73]).⁴

There are also good reasons to accept a metalogic that is at least as strong as the object-logics one accepts. First, as Woods [73] has pointed out, if ML is weaker than L, then a purported conflict between a piece of evidence e and a rule R of L could simply be removed by removing the inference from R to $\neg e$ in ML. Second, as Bremer [8] has shown, the common core logic of competing logics $L_1, L_2 (\dots)$ is too weak as a framework for comparing those logics (cf. [63], 9). These considerations fit the common (though not ubiquitous) practice to use the strongest logic, namely classical logic, for the philosophical evaluation of non-classical logics. Typically the evidence for or against a non-classical logic is also described within classical metalogic, which is self-undermining in a sense.

On the other hand, as we have seen, for every ‘logical’ rule, no matter how bizarre, it is possible to adopt a suitably strong metalogic that permits a rule-circular ‘proof’ of the rule’s validity. In conclusion, a non-circular strategy of justifying a system of logic is not in sight. All this seems to speak in favor of the *exceptional* status of logic as irreducibly basic. In combination with the plurality of competing logical systems, this

⁴ Besides experiential facts, a further sort of evidence used by anti-exceptionalists are intuitions about validity. Intuition-based evidence has been criticized as subjective and unreliable (cf. [33, 61]).

exceptional feature of logic seems to undermine the possibility of foundation-oriented epistemology. There are two threats emerging from this situation. *First*, there is the threat of *relativism*: It appears to be largely a *subjective* matter which logic one chooses, based on intuition or ‘cognitive taste’, but not upon objective reason. The threat of relativism would not be so bad if the choice of a logic were largely a matter of convention, like the choice of a conventional terminology. But the opposite appears to be the case: logic seems to constitute the innermost core of our belief system, determining all other parts. This brings us to the second threat - that of *incommensurability*: Proponents of different logics appear to be in a fundamental disagreement (about the logic one should use) that cannot be resolved by rational argumentation, nor even be formulated in a neutral way, because every formulation presupposes the adoption of a logic.⁵

2 Optimality Justifications Based on Translations between Logics: Escape from Incommensurability and Relativism?

Theoretically the problem of incommensurability appears to be even worse than that of relativism, as it seems to prevent a rational discourse between proponents of alternative logics. This theoretical diagnosis stands in remarkable contrast to the practice of logicians: proponents of different logics have no problem understanding each other, but exchange rational arguments all the time. How is this possible? The answer lies in hidden translation possibilities between non-classical and classical logics.

The study of translations methods between different logics has become an area of increasing research in contemporary logic [12]. The possibility of a particular kind of translations, namely *meaning-preserving* translations, constitutes one of the two major ideas underlying the present paper. Meaning-preserving translations between different systems of logic remove the threat of incommensurability, but not yet the threat of relativism, because they don’t answer the question of how one can justify the choice of a logic in a non-circular way. The second idea of this paper is to answer this question by the method of *optimality justifications*.

The method of optimality justifications has been developed as an approach to the problem of induction, or Hume’s problem [58, 60]. An optimality justification does not attempt to demonstrate that a given epistemic method is strictly or probabilistically reliable, in the sense of leading to the truth in all or most cases. It pursues a more modest goal, namely to demonstrate that a given method is *optimal* in regard to a given epistemic goal, among all competing methods that are *cognitively accessible* to the given epistemic agent. It can be proved that in all possible worlds a certain method of *meta-induction* is predictively optimal in the long run, among all prediction methods that are accessible to the forecaster.⁶

In this paper the method of optimality justifications is applied to the domain of logic, focusing on propositional logic and its modal extensions. Optimality

⁵ The problem of ‘incommensurability’ was introduced by Kuhn [40] for the rational comparison of competing paradigms in science.

⁶ More information in Schurz [60], Henderson [30], Sterkenburg [66], Feldbacher [19].

justifications are relative to a given epistemic *goal*. In the case of induction, this goal is predictive success. What could the epistemic goal of logic be? In contemporary philosophical logic one often discusses goals such as intuitive naturalness, usefulness in certain domains, or agreement with assumed metaphysical positions. The problem with these goals is that they are too subjective and context-dependent to enable a robust objective optimality justification. For example, Priest [51], Williamson [71] as well as Bueno [10] and Hjortland [32] are anti-exceptionalists, but they draw opposite inferences about the preferred logic: Williamson prefers classical logic, Priest argues for paraconsistent logic, while Hjortland and Bueno opt for a version of logical pluralism.⁷ For a robust epistemic justification we need a more general epistemic goal, indeed a *most* general one, since logic is the most general level of description. The goal we propose for this purpose is power of linguistic representation, in short: *representational power*. Searching for optimality in regard to this goal is possible by applying meaning-preserving translations between different systems of logic.

In the next sections (3–5) it will be demonstrated that the most well-known systems of non-classical logics can be translated into classical logic. Our technique of achieving a meaning-preserving translation will consist in extending the language of classical logics by intensional operators who reflect the non-classical structure inside of their scope but behave classically outside of their scope. Based on this result it is argued that classical logic is representationally optimal: by using classical logic one can only win but not lose, because if a non-classical logic turns out to have advantages for certain purposes, it can be translated into classical logic. However, optimality doesn't imply dominance, but is compatible with a plurality of equally optimal logics. In sec. 6 it will be investigated whether non-classical logics can be likewise representationally optimal. The answer is predominantly negative but partially positive. Nevertheless the situation is not symmetric, because arguably classical logic has important *ceteris paribus* advantages in its role as a unifying metalogic.

3 Translation of Many-Valued Logics

3.1 Translation of Lukasiewicz' Three-Valued Logic into Classical Logic

In this section we develop the translation approach in application to Lukasiewicz' three-valued (propositional) logic L_3 with the truth-values true (t), false (f) and undetermined (u). Let us assume that the sentence "there is exactly one electron" is indeed neither true nor false but undetermined. Then, despite the assumed indeterminacy of this sentence, the statement that the sentence is undetermined is itself two-valued, either true or false, and certainly not undetermined. More generally, even in the trivalent logical framework, all

⁷ Different notions of logical pluralism have been developed, e.g. *local* pluralism versus *global* pluralism ([31], 356 f.) and *contextual* pluralism [11]. Bueno's pluralism is contextualist and localist (different logics can be adequate for different domains and purposes).

statements asserting the truth, indeterminateness or falsity of a certain statement are bivalent - since assuming a trivalent semantic assignment just means that these statements are bivalent. If we manage to translate all statements of the three-valued framework into combinations of bivalent statements of this sort, we have found a translation function.⁸

Let us carry out this translation in a precise way. The language L_3 has four basic truth-functional connectives \neg, \vee, \wedge and \rightarrow , where the three-valued conditional \rightarrow is not definable in terms of the other three connectives. As usual one assumes a linear ordering among the truth-values of a (finite) multi-valued logic; in the case of L_3 the ordering is $f < u < t$; or represented as ranks: $-1, 0, +1$. Based on this ordering, Lukasiewicz' three-valued truth-tables for the four connectives are described as follows: the truth value of $\neg p$ is the inverse of p 's truth value, that of $p \wedge q$ is the minimum and that of $p \vee q$ the maximum of the truth values of p and q . Finally, $p \rightarrow q$'s truth value equals $t/u/f$ if the rank difference between q 's truth value and p 's truth value is not smaller than $0 / -1 / -2$, respectively.

p	$\neg p$	p	q	$p \wedge q$	$p \vee q$	$p \rightarrow q$
t	f	t	t	t	t	t
u	u	t	u	u	t	u
f	t	t	f	f	t	f
		u	t	u	t	t
		u	u	u	u	t
		u	f	f	u	u
		f	t	f	t	t
		f	u	f	u	t
		f	f	f	f	t

The notion of logical truth and validity in L_3 is defined analogously as in bivalent logics. Let \wp be the denumerable set of propositional variables, and $\text{val}_3: \wp \rightarrow \{t, u, f\}$ range over trivalent truth-valuations of the variables, recursively extended to arbitrary complex L_3 -formulas via the above truth tables. Then an L_3 -formula A is *logically true* in L_3 , in short $\models_3 A$ iff $\text{val}_3(A) = t$ for all (possible) trivalent valuations, and A *follows* from a formula set Γ in L_3 , in short $\Gamma \models_3 A$, iff all trivalent valuations making all formulas in Γ true make A true. Here are some examples:

$$\text{Some theorems of } L_3 : p \rightarrow (q \rightarrow p), (\neg q \rightarrow \neg p) \leftrightarrow (p \rightarrow q), (p \vee q) \leftrightarrow ((p \rightarrow q) \rightarrow q).$$

⁸ A possible objection comes from the account of high-order vagueness, or indeterminacy. Proponents of this position argue that the truth-value statements Tp, Up and Fp – short for “proposition p is true, undetermined or false, respectively” – may themselves be true, undetermined or false. A possible defeat of this objection goes as follows. There are two possibilities ([64], sec. 1 + 4): Either (i) one stops the vagueness-iteration at some finite level k , or (ii) the iteration goes on forever. In case (i), the truth-values of degree k ($X(p)$ for $X \in \{T, U, F\}^k$) are bivalent. In case (ii) one usually assumes an assignment of real numbers; then the statements of the form “the truth-degree of p is r ” ($T(p) = r$) are bivalent. Thus, there seems to be always an ultimate level of description at which descriptions are bivalent.

Some non-theorems of L_3 : $p \vee \neg p$, $\neg(p \wedge \neg p)$, $(p \vee q) \leftrightarrow (\neg p \rightarrow q)$.

Conditional proof $\Gamma, A \models B / \Gamma \models A \rightarrow B$ fails for L_3 , because in L_3 the logical truth of $A \rightarrow B$ is *stronger* than the validity of A/B : while the latter implies merely that whenever A is true, B is true, the former implies in addition that when A is undetermined, B is undetermined or true. For example, $p \wedge \neg p \models_3 q$ but $\not\models_3 (p \wedge \neg p) \rightarrow q$.

Our translation is based on the expansion of \mathcal{L}_2 by three operators T , U and F that express the truth values of being true, undetermined, and false in three-valued logic. If p is a sentence of the three-valued logic, the sentences $T(p)$, $U(p)$ and $F(p)$ are bivalent and obey the following truth table:

p	$T(p)$	$U(p)$	$F(p)$
t	t	f	f
u	f	t	f
f	f	f	t

We don't need to introduce these operators in \mathcal{L}_3 , because they are definable in L_3 as follows:

$$\begin{aligned} U(A) &=_{\text{def}} (A \vee \neg A) \rightarrow (A \wedge \neg A), \\ T(A) &=_{\text{def}} A \wedge \neg U(A), \text{ and} \\ F(A) &=_{\text{def}} \neg A \wedge \neg U(A). \end{aligned}$$

It is easy to check that the truth-functions of these formulas in L_3 coincide with the above truth tables for T , U and F .

By adding the Lukasiewicz-operators T , U , F to the classical language \mathcal{L}_2 we obtain the extended classical language $L_{2,\text{Luk}}$ whose formulas are still evaluated bivalently and whose basic logical laws are still the classical laws of L_2 . Within L_2 the operators T , U and F figure as *intensional* (non-bivalently-truth-functional) operators, similar to the operators of modal logic. Based on the truth tables of these three operators, every semantic rule of three-valued logic can be translated into a set of corresponding axioms formulated in the extended language of classical logic as follows:

$$\begin{aligned} \text{--For the negation: } & T(\neg A) \leftrightarrow F(A) \quad U(\neg A) \leftrightarrow U(A) \quad F(\neg A) \leftrightarrow T(A) \\ \text{--For the conjunction: } & T(A \wedge B) \leftrightarrow T(A) \wedge T(B) \quad F(A \wedge B) \leftrightarrow F(A) \vee F(B) \\ & U(A \wedge B) \leftrightarrow (U(A) \wedge \neg F(B)) \vee (U(B) \wedge \neg F(A)) \\ \text{--For the disjunction: } & T(A \vee B) \leftrightarrow T(A) \vee T(B) \quad F(A \vee B) \leftrightarrow F(A) \wedge F(B) \\ & U(A \vee B) \leftrightarrow (U(A) \wedge \neg T(B)) \vee (U(B) \wedge \neg T(A)) \end{aligned}$$

$$\begin{aligned}
 \text{--For the implication : } & T(A \rightarrow B) \leftrightarrow F(A) \vee (U(A) \wedge \neg F(B)) \vee (T(A) \wedge T(B)) \\
 & U(A \rightarrow B) \leftrightarrow (T(A) \wedge U(B)) \vee ((U(A) \wedge F(B))) \\
 & F(A \rightarrow B) \leftrightarrow T(A) \wedge F(B)
 \end{aligned}$$

- Finally we add the trivalent truth-value axiom: $T(A) \vee U(A) \vee F(A)$ (“ \vee ” for exclusive disjunction, defined as usual).

The set of these axiom schemata forms the axiom system Ax_{Luk} of Lukasiewicz’ logic in the extended language of classical logic $\mathcal{L}_{2.Luk}$.

We now turn to our proposed method of translating \mathcal{L}_3 -assertions into \mathcal{L}_2 -assertions. Our method is based on the *truth view of assertion*: asserting a sentence A means to assert that A is true (more on this in sec. 3.2.1). Thus our translation function “ $\tau_{3 \rightarrow 2}$ ” (from \mathcal{L}_3 into \mathcal{L}_2) is this:

$$(3) \text{ For all } A \in \mathcal{L}_3: \tau_{3 \rightarrow 2}(A) = T(A).$$

Note that the translation based on the assertion view is not recursive but *holistic*: it translates every complex \mathcal{L}_3 -assertion at once into a statement of $\mathcal{L}_{2.Luk}$. By applying the axioms in Ax_{Luk} we can then transform every sentence of $\mathcal{L}_{2.Luk}$ into a truthfunctional combination of modalized variables (Tp_i , Up_i or Fp_i) and thus find out what the translation means for the truth-value of the modalized variables of the sentence.

Some examples of translations and their equivalent transformations (“=” for identity, “ \leftrightarrow ” for material L_2 -equivalence given Ax_{Luk} , “ τ ” short for “ $\tau_{3 \rightarrow 2}$ ”):

$$\begin{aligned}
 \tau(p) &= T(p), \tau(\neg p) = T(\neg p) \leftrightarrow F(p), \tau(p \vee \neg p) = T(p \vee \neg p) \leftrightarrow Tp \vee Fp, \tau(p \wedge \neg p) \\
 &= T(p \wedge \neg p) \leftrightarrow Tp \wedge Fp, \tau(p \rightarrow q) = T(p \rightarrow q) \leftrightarrow Fp \vee (Up \wedge (Uq \vee Tq)) \vee (Tp \wedge Tq).
 \end{aligned}$$

Note that we use the same logical symbols for the two-valued and the three-valued propositional connectives (e.g., both “ \rightarrow ” for two-valued and three-valued implication). This is not a problem, because whenever we translate an \mathcal{L}_3 -formula into \mathcal{L}_2 , the three-valued connectives are *hedged* in the scope of the intensional truth value operators T , U and F . When we bring them out of their scope by applying the equivalences of Ax_{Luk} , the three-valued connectives are transformed into the two-valued connectives of L_2 .

We now show that the translation $\tau_{3 \rightarrow 2}$ preserves meaning of modalized formulas and L_3 -logical truth (or validity) in a precise sense. For this purpose, we introduce some further terminology. In what follows, O_i ranges over the three trivalent truth-value operators, T , U and F . Let $P(A) = \{p_1, \dots, p_n\}$ be the set of variables occurring in sentence A . We speak of the “ p_i ” as “unmodalized variables” and of the statements “ $O_i p_j$ ” as the “modalized” variables. For \wp a set of unmodalized variables, $O_\wp = \bigcup_{p \in \wp} \{Tp, Up, Fp\}$ denotes the corresponding set of modalized variables. If \wp is the (denumerable) set of unmodalized variables common to \mathcal{L}_3 and $\mathcal{L}_{2.Luk}$, then

following from the intensional nature of the operators O_i , bivalent truth-valuations over $\mathcal{L}_{2,\text{Luk}}$ are defined over the set of elementary formulas $\wp \cup O\wp$. Let $\text{Val}_3(\wp)$ be the set of all trivalent valuations over \wp and Val_3 be the set of (recursively extended) trivalent valuations over sentences of \mathcal{L}_3 . Moreover, let $\text{Val}_{2,\text{Luk}}(O\wp)$ be the set of all bivalent valuations over $O\wp$ satisfying the axiom $T(A) \dot{\vee} U(A) \dot{\vee} F(A)$. Moreover, let $\mathcal{L}_{2,\text{Luk},\tau}$ be the set of $\mathcal{L}_{2,\text{Luk}}$ -formulas that are Ax_{Luk} -equivalent with translations of \mathcal{L}_3 -formulas, and let $\text{Val}_{2,\text{Luk},\tau}$ be the set of all recursively extended bivalent truth-valuations over formulas of $\mathcal{L}_{2,\text{Luk},\tau}$ that satisfy the equivalence axioms of Ax_{Luk} . Then we can prove:

(4) **Theorem** (L_3 into L_2):

4.1 For every three-valued valuation function $\text{val}_3 \in \text{Val}_3$ there exists exactly one two-valued valuation function $\text{val}_2 =_{\text{def}} f(\text{val}_3) \in \text{Val}_{2,\text{Luk},\tau}$ (f an injective function) such that for every \mathcal{L}_3 -formula A

$$\text{val}_3(A) = o \text{ iff } \text{val}_2(O(A)) = t, \text{ where } O = T/U/F \text{ iff } o = t/u/f.$$

4.2 An \mathcal{L}_3 -statement A is logically true in L_3 iff $T(A)$ follows logically from Ax_{Luk} in L_2 , and analogously for logical consequence. Thus

$$|=_{L_3} A \text{ iff } \text{Ax}_{\text{Luk}} |=_{L_2} \tau(A), \text{ and}$$

$$\Gamma |=_{L_3} A \text{ iff } \text{Ax}_{\text{Luk}} \cup \tau(\Gamma) |=_{L_2} \tau(A), \text{ where } \tau(\Gamma) =_{\text{def}} \{\tau(B) : B \in \Gamma\}.$$

According to theorem 4.2 the translation function is conservative in the sense of Carnielli et al. [12]. The proof of theorem (4) is based on two lemmata:

Lemma 1 There exists an injective function $f: \text{Val}_3(L) \rightarrow \text{Val}_{2,\text{Luk},\tau}$ whose restriction to propositional atoms satisfies

$$(*) \text{ for all } p \in \wp : \text{val}_3(p) = o \text{ iff } \text{val}_2(Op) = t, \text{ where } O = T/U/F \text{ iff } o = t/u/f.$$

Proof For every $\text{val}_3 \in \text{Val}_3(\wp)$, we define $f(\text{val}_3) = \text{val}_2 \in \text{Val}_{2,\text{Luk}}(O\wp)$ by the condition (*). The definition satisfies the axiom $T(A) \dot{\vee} U(A) \dot{\vee} F(A)$. The bijection f extends uniquely to all \mathcal{L}_3 -formulas by the requirement that val_2 satisfies the equivalence axioms in Ax_{Luk} ; this requirement determines the truth values of the translations of all \mathcal{L}_3 -formulas and their Ax_{Luk} -equivalents in $\mathcal{L}_{2,\text{Luk},\tau}$. Q.E.D.

Lemma 2 For every $A \in L_3$ with $\wp(A) = \{p_1, \dots, p_n\}$, Ax_{Luk} L_2 -entails that $O(A)$ is equivalent with a distinguished disjunctive and negationless normal form $DN(OA)$, each elementary disjunct being a conjunction of modalized variables $O_1 p_1 \wedge \dots \wedge O_n p_n$ (the ‘trivalent constituents’).

Proof By successive application of the equivalence transformations corresponding to the axioms of Ax_{Luk} , one can drive the truth-value operators successively inside the formula until they stand immediately before the unmodalized variables; negations are thereby eliminated.⁹ By applying $\wedge - \vee$ distribution laws and expanding conjuncts in which certain modalized variables Op_i in $O_{\wp}(A)$ are missing (by replacing such a conjunct X by $(X \wedge Tp_i) \vee (X \wedge Up_i) \vee (X \wedge Fp_i)$), one can produce the required disjunction of trivalent constituents, abbreviated as $DN(OA)$. Q.E.D.

Proof of Theorem 4.1 Let $A \in \mathcal{L}_3$. The proof rests on the fact that every line (' val_3 ') of A 's trivalent truth table with $val_3(A) = o$ corresponds exactly to one of $O(A)$'s trivalent constituents in L_2 . Here is the formal proof: $val_3(A) = o$ iff $val_3 \models O(A)$ iff $val_3 \models DN(OA)$ (since $val_3 \models A \leftrightarrow DN(OA)$) iff $val_3 \models C$ for some conjunct C of $DN(OA)$ iff $val_2 = f(val_3)$ verifies C (by lemma 1) iff $val_2 \models DN(OA)$ iff $val_2 \models O(A)$ (by lemma 2). Q.E.D.

Proof of Theorem 4.2 $\models_{L_3} A$ iff $\forall val_3 \in Val_3: val_3 \models A$ iff $\forall val_2 \in Val_{2,Luk,\tau}: val_2 \models TA$ (by theorem 4.1) iff $\models_{L_2} \tau(A)$. - Analogously for inferences. Q.E.D.

3.2 Discussion of the Proposed Translation Method

In the next four subsections we discuss subtleties and achievements of the proposed translation method.

3.2.1 Meaning Preservation

Our translation preserves meaning as well as possible. It is not literally possible to translate a three-valued atomic proposition p into a two-valued logic, because in L_2 it has only two but in L_3 it has three truth values. What can be literally translated, however, are the more *fine-grained* propositions Tp , Up and Fp , whose meaning is strictly the same in L_3 and in L_2 . This makes it possible to translate all assertions expressible in L_3 , given the truth view of assertions. Moreover, by applying the Ax_{Luk} -equivalences (that are valid in $L_{2,Luk}$ as well as in L_3), the semantic composition of A in L_3 is fully reflected in L_2 . We conclude that *every proposition that can be expressed in L_3 can be also expressed in L_2* .

Kooi and Tamminga [37] also use a translation function for truth-value assertions in their translation of L_3 into the modal logic S4. Their translation differs from ours in two respects: First, they translate the truth-value assertions Tp , Up and Fp into the modal

⁹ Here is the formal inductive proof: We must show that (*) for every $O_i \in \{T,U,F\}$, $Ax_{Luk} \models O_i(A) \leftrightarrow A'$ holds, where A' satisfies the condition (**) that all O_i stand in front of propositional atoms. Induction hypothesis: Assume (*) is proved for all $O_i(A)$ formulas of degree n , where "degree(A)" is the maximal number of propositional operators in the scope of a truth operator in A . Induction step: Assume $O_i(A)$ has degree $n + 1$. Then A has one of the forms $\neg B$, $B \wedge C$, $B \vee C$ or $B \rightarrow D$. By applying the respective Ax_{Luk} equivalence axioms, $O_i(A)$ is Ax_{Luk} -equivalent with the corresponding formula schemata A'' at the right side, whose degree is n . By induction hypothesis we can replace every subformula E of A'' of degree n by a subformula E' satisfying condition (**). This replacement leads to a formula A' equivalent with $O_i(A)$ and satisfying (**). Q.E.D.

S4-sentences $\Box p$, $\Diamond p \wedge \neg \Box p$ and $\neg \Box p$, respectively (ibid., 1064), while we use the intensional operators T, U and F. *Second*, they build the equivalences that we expressed by means of the axioms $Ax_{L_{luk}}$ directly into their translation function for complex L_3 -sentences, by translating a complex L_3 -sentence directly into the disjunction of its modal constituents, while we proved in lemma 2 that a sentence is $Ax_{L_{luk}}$ -equivalent with this disjunction. We prefer our translation method for the epistemological reason that it makes the embedding of L_3 into an axiom set in $L_{2,L_{luk}}$ explicit.

3.2.2 Holistic Nature of the Translation

The deeper reason behind the ‘holistic’ nature of our translation function is the fact that in L_3 the correspondence schema of truth is no longer valid: $TA \leftrightarrow A$ has the value f if A has the value u, since $f \leftrightarrow u = f$. In combination with the truth view of assertion this implies that a three-valued proposition and the assertion of this proposition are not logically equivalent. Therefore the translation of A by T(A) cannot be applied in a recursive fashion. For example, the assertion “p” is translated as “Tp” (because this is what “p” asserts), but the assertion “ $\neg p$ ” cannot be recursively translated as “ $\neg Tp$ ”, because what it asserts is “T($\neg p$)” ($\text{val}(p) = f$) and this is stronger than $\neg Tp$ ($\text{val}(p) \in \{u, f\}$). This epistemological subtlety has been reflected in early writings about three-valued logic, e.g. in Woodruff’s distinction between a strong and a weak mode of assertion ([72], 122).

3.2.3 Comparison with the Literature

The expansion of classical truth-functional logic by the intensional operators T, U, F makes it possible to preserve the meaning of L_3 -assertions. This meaning-preservation feature distinguishes my account from most translation functions between logics studied in the literature (for an overview cf. [12]). In these studies, translations are usually not accompanied by *expansions* of the (classical) language, at the cost that the translation functions do not preserve the meaning and semantic composition of the translated statements; they only preserve the consequence operation. One example is the abstract translation functions studied by Jerábek [35]. These translation functions map the formulas of the language \mathcal{L} of a propositional logic L into formulas of a language \mathcal{L}' of a logic L', such that if $A \vdash_L B$, then $f(A) \vdash_{L'} f(B)$, roughly speaking as follows: given an enumeration of all \mathcal{L} -formulas and the nth formula A_n of \mathcal{L} , $f(A_n)$ is defined as $X \vee (q_n \wedge Y)$, where X is the disjunction of the translations of all premises with indices smaller than n that entail A_n , q_n is a new variable and Y is the conjunction of all translations of implications $C \rightarrow D$ with indices smaller than n such that $\{C, A_n\}$ entails D (ibid., 669). Jerábek [35] proves that L_2 is translation-universal in the sense that every finitary deductive system in countably many formulas can be conservatively translated into L_2 . Many other but not all non-classical propositional logics are translation-universal in this sense. These results are technically impressive. However, the defender of a non-classical logic can object that the translation is not meaning-preserving; it does not reflect what the non-classical assertions express and thus is not suitable for the epistemic purpose of an optimality justification in regard to representation power.

An example of a non-meaning-preserving translation at the semantic level is the reduction of many-valued logics to bivalent logics proposed by Suszko [67]. Given a many-valued logic with a subset $Des \subset Val$ of designated truth values, Suszko proposed

to translate the disjunction (or set) of the designated truth-values into the bivalent value “true” and the disjunction (or set) of the non-designated truth-values into “false”. Suszko’s translation is useful for many purposes (cf. [5]). However, Suszko’s translation does not preserve the meaning of the propositional connectives; they become intensional under Suszko’s bivalent semantics ([45], 79; [70]). For example, both p and $\neg p$ may have the truth-value false; thus the law of excluded middle, $p \vee \neg p$, is no longer valid in Suszko’s bivalent semantics. Therefore Suszko’s bivalent semantics is not classical and does not yield a translation of L_3 into a classical logic.

3.2.4 Bridge Axioms between L_3 and L_2

For every $S \in L_3$, the truth value of $T(S)$ depends only on the truth values of its modalized but not of its unmodalized variables. So far, the bivalent truth-values of the modalized variables ($O_i p$) have not been related to the bivalent truth-values of the unmodalized variables. For the semantic coherence between L_3 and L_2 we require the following

(5) Bridge axioms (from $\tau(L_3)$ to L_2): $Tp \rightarrow p$ and $Fp \rightarrow \neg p$.

In words, a trivalently true (or false) atomic sentence is also bivalently true (or false, respectively), while for undetermined sentences their bivalent truth value is left *open*. One can prove that (5) entails the validity of the bridge axioms $TA \rightarrow A$ and $FA \rightarrow \neg A$ for all \mathcal{L}_3 -formulas built up from \neg, \wedge and \vee . Interestingly, the unrestricted bridge axioms may be violated for Lukasiewicz’ implicational formulas¹⁰; but they are valid for Kleene’s three-valued implications, which differ from Lukasiewicz’ implications in having $u \rightarrow u = u$ instead of $u \rightarrow u = t$.

The *converse implications* ($p \rightarrow Tp$ and $\neg p \rightarrow Fp$) *must not* hold, since otherwise the translation would not be conservative and the translated three-valued logic would collapse into two valued logic: If we would accept the inverse bridge axioms, then $\models_{L_2} p \vee \neg p$ would imply $\models_{L_3} T(p \vee \neg p)$ and thus $\models_{L_3} p \vee \neg p$.

3.3 Generalization to Many-Valued Logics

Lukasiewicz’ L_3 was later mathematically generalized to *many-valued* logics with arbitrarily many ‘truth-values’, abstracting from a particular philosophical interpretation of these truth-values [26, 45]. A multi-valued logic L in the generalized sense is semantically characterized by A set of truth-values Val and a subset $Des \subset Val$ of designated truth-values; a formula A is defined as logically true if all possible valuations of the propositional variables assign to A a designated value. The triple $\langle Val, Des, \{t_c: c \in C\} \rangle$ is called a Val -valued *logical matrix*, with $\{t_c: c \in C\}$ being the set of truth-tables for a set of connectives C .

The translation strategy of sec. 3.1 applies to all many-valued logics that are representable by means of *finitely* many truth values. Thus, if an n -valued logic L_n is based on a matrix $\langle Val_n, Des_k, \{t_c: c \in C\} \rangle$ with $|Val_n| = n$, $|Des_k| = k < n$ and $C = \{\neg, \wedge, \vee, \rightarrow\}$, then we extend L_2 by n intensional truth value operators O_1, \dots, O_n

¹⁰ A counterexample: if $val_3(p_1) = val_3(p_2) = u$, then $val_3(\pm p_1 \rightarrow \pm p_2) = t$ and thus $val_2(T(\pm p_1 \rightarrow \pm p_2)) = t$ for all $\pm p_i \in \{p_i, \neg p_i\}$; but $val_2(\pm p_1 \rightarrow \pm p_2) = t$ can not hold for all $\pm p_i \in \{p_i, \neg p_i\}$.

together with the equivalence axioms describing L_n 's truth tables for \neg , \wedge , \vee and \rightarrow , and add the n -valent truth-value axiom $O_1(A) \dot{\vee} \dots \dot{\vee} O_n(A)$. For all many-valued logics it is natural to assume the *designated-value view* of assertions: asserting p in a many-valued logic means asserting that p has a designated value. Based on this view, we can then prove the translation theorem for the assertions of the many-valued logic L_n in the same way as above, by demonstrating that $\models_{L_n} A$ iff $\models_{L_2} \text{Des}(A)$, with $\text{Des}(A)$ abbreviating $\dot{\vee} \{O_i(A) : v_i \in \text{Des}_k\}$ (and likewise for inferences). Of course, if the number of truth values is high, the disjunctive normal forms of the L_2 -translations of L_n -formulas become rather complex (for m propositional variables there are n^m possible constituents). However, we conjecture that in many cases fewer intensional operators than truth values are needed to express the translation function; an investigation of this question is work for the future.

4 Translating Paraconsistent and Intuitionistic Logics

4.1 Translating the Paraconsistent Logic LP

As explained in sec. 1, the major motivation of paraconsistent logics is to admit the possibility of statements that are both true and false, but to prohibit the classically valid inference of ‘explosion’ or ‘ex falso quod libet’ (EFQ): $p \wedge \neg p / q$. Not all but many paraconsistent logics can be characterized by means of finite truth value matrices ([52], sec. 3.6). The simplest paraconsistent logic is the *logic of paradox*, LP, developed by Priest [50]. It is a species of a three-valued logic, having instead of the third value “undetermined” the value “both true and false”, abbreviated as b (or “ p ” for “paradoxical”, as in [50]). The values “ t ” and “ f ” now mean “true only” and “false only”, respectively. If “ u ” is replaced by “ b ”, then the three-valued matrices for the propositional connectives of LP are those of Kleene’s [36] three-valued logic, which are ‘almost’ identical with Lukasiewicz’ three-valued logic L_3 , with the exception of one line in the truth table for the implication, namely $b \rightarrow b = b$ (instead of $u \rightarrow u = t$, as in [44]). The important difference of LP compared to L_3 is that LP considers both values t and b as *designated*, which changes the resulting LP-valid theorems and inferences radically. One the one hand, LP’s logical theorems are now identical with the theorems of classical logic, i.e., $LP = L_2$. For example, $p \vee \neg p$ is LP-true, since under all valuations $\text{val}: P \rightarrow \text{Val}$, $\text{val}(p \vee \neg p)$ comes out either as t or as b . On the other hand, many classically valid inferences become invalid. For example, (EFQ) $p \wedge \neg p \models q$ is invalid, since there are valuations assigning b to p and thus to $p \wedge \neg p$, but f to q . While the elimination of (EFQ) is the central intended feature of LP, four further highly plausible inferences become LP-invalid, too, namely (DS): $p \vee q, \neg p / q$, (MP): $p, p \rightarrow q / q$, (MT): $p \rightarrow q, \neg q / \neg p$ and (Trans \rightarrow): $p \rightarrow q, q \rightarrow r / p \rightarrow r$ ([50], 228).

In spite of these differences, the strategy of translating the logic LP into classical logic L_2 is the same as for L_3 as outlined in section 3.1: we extend \mathcal{L}_2 by the three intensional operators T (true only), B (both true and false) and F (false only) and represent their truth-tables as axioms of the modal theory AX_{LP} in the extended language $\mathcal{L}_{2,\text{para}}$. As in all other many-valued logics, we translate LP’s assertions into L_2 based on the designated-value view of assertions. Let “ $D(A)$ ” stand for “ A has a designated truth value”, being an abbreviation for “ $T(A) \vee B(A)$ ”. Then we can prove a

translation theorem analogously to theorem (4), but with $T(A)$ and $T(\Gamma)$ replaced by $D(A)$ and $D(\Gamma)$, respectively (where $D(\Gamma) := \{D(B) : B \in \Gamma\}$):

$$(6) \textbf{Theorem (LP into } L_2\text{): } \Gamma \models_{LP} A \text{ iff } Ax_{LP} \cup D(\Gamma) \models_{LP} D(A).$$

Based on this translation we can show, for example, that $D(p \wedge \neg p) \not\models D(q)$, and likewise that $D(p \vee q), D(\neg p) \not\models D(q)$ and $D(p \rightarrow q), D(p) \not\models D(q)$ (etc.), i.e., the translations of LP-invalid inferences come out as invalid, too.

So far the situation seems to be similar as that for Lukasiewicz' L_3 , but there is a deep hidden difference: While the trivalent truth operators T, U and F , whose values are always bivalent, are explicitly definable in L_3 , the corresponding truth operators T, B and F are undefinable in LP - ultimately because of the tiny difference that $b \rightarrow b = b$ (instead of t). This makes it impossible to define "Bp" via " $(p \vee \neg p) \rightarrow (p \wedge \neg p)$ " and implies that "Tp" and "Fp" are also undefinable in LP. Therefore, the translation of LP into L_2 yields a proper expressive extension of LP: in $L_2 \cup Ax_{LP}$ we can express and prove inferences that are inexpressible in LP. For example, we can prove

$$(7) T(p \wedge \neg p) \models_2 Tq, \quad T(p \vee q), T(\neg p) \models_2 Tq, \quad T(p \rightarrow q), Tp \models_2 Tq,$$

which are the T-modalized versions of EFQ, DS and MP, that are valid in LP, but inexpressible in LP. Proponents of LP may see this as a disadvantage, but we see it as an advantage, because 'true-only'-assertions are part of LP's semantics, so why should these assertions not also be expressible in the object language? But within LP it is impossible to find formulas that are necessarily bivalent, i.e. either true or false but not both. A consequence of this expressive weakness of LP explained in section 6 is that it makes an inverse translation of L_2 into LP impossible.

4.2 Translating Intuitionistic Logic into S4

As explained in sec. 1, in intuitionistic logic L_i the classical notions of truth and falsity are replaced by the concepts of mathematical verification and falsification, where a verification of p consists in a constructive proof with conclusion p , and a falsification in a constructive proof from the assumption p to a contradiction $q \wedge \neg q$. Since by the results of Gödel and Church there exist mathematical propositions whose truth value is undecidable, this has the consequence that certain laws of classical logic are no longer valid, foremost the laws of double negation ($\neg\neg p \rightarrow p$) and of excluded middle ($p \vee \neg p$). Intuitionistic logic cannot be algebraically represented by finite matrices [23]; thus the translation strategy of sec. 3 does not apply. A well-known 'translation' of L_i into L_2 is based on Glivenko's double-negation theorem: $\Gamma \models_2 A$ iff $\neg\neg(\Gamma) \models_1 \neg\neg A$ (with $\neg\neg(\Gamma) = \{\neg\neg B : B \in \Gamma\}$). Since this translation does not preserve meaning (as p and $\neg\neg p$ mean different propositions in L_i), it is not suitable for our purpose. There is, however, a well-known translation of L_i into a modal extension of L_2 , namely into the modal logic S4, that fits perfectly into our account of meaning-preserving translations of the non-classical *assertions*. The philosophical characteristics of intuitionistic logic, as opposed to classical or many-valued logics, is that it replaces the truth view of assertion by the *provability view of assertion*: asserting that p in L_i means to assert that

p has a constructive proof. Thus it is natural to translate the assertion that p in intuitionistic logic by $\Box p$, where the modal operator “ \Box ” is understood (not as meta-physical necessity, but) as constructive provability. As in the translation of many-valued logics, this translation gives us bivalence outside the provability operator: although $\Box p \vee \Box \neg p$ is not L-true, $\Box p \vee \neg \Box p$ is again L-true.

Constructive provability satisfies the laws of the modal logic S4, semantically characterized by the class of reflexive and transitive Kripke frames (W,R) (i.e., the accessibility relation $R \subseteq W^2$ between worlds is reflexive and transitive). The S4-axioms are (T): $\Box A \rightarrow A$ (what is provable is true, corresponding to reflexivity), (4): $\Box A \rightarrow \Box \Box A$ (provability of propositions is itself provable, corresponding to transitivity), and the axioms and rules satisfied by all Kripke frame logics, namely (K): $\Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$ and (N) $\models A / \models \Box A$.

Before we turn to the syntactic translation of L_i into S4 we briefly recapitulate how the Kripke semantics for L_i implements the translation idea in a semantic fashion ([47], sec. 5.1). The worlds of these frames are interpreted as epistemic states and the accessibility relation R expresses a natural (e.g. temporal) *successor* relation between these states. The valuation function over these frames is merely *partial*: a propositional atom may either be intuitionistically true or false or undefined at a world α . This makes sense, since intuitionistic truth means provable truth, but not every p is either provable or disprovable, i.e., $p \vee \neg p$ is not L_i -true. Since intuitionistic truth is provable truth, the valuation function val has to satisfy the following constraint: if p is intuitionistically true at a state α ($\alpha \models p$), then p is true at all states β accessible from α ($R\alpha\beta$ implies $\beta \models p$). Intuitionistic truth of conjunction and disjunction is recursively defined as classical truth, i.e. $\alpha \models A \wedge B / A \vee B$ iff $\alpha \models A$ and/or $\alpha \models B$. Intuitionistic truth of negation and implication, however, is more subtle: $\alpha \models \neg A$ iff for no β accessible from α , $\beta \models A$ holds (i.e., every β accessible from α can access a world in which A is not intuitionistically true), and $\alpha \models A \rightarrow B$ iff for every world β accessible from α , $\beta \models A$ implies $\beta \models B$. Kripke [39] proved that L_i is complete for this semantics, using a classical metalogic.¹¹

By transforming the Kripke semantics for L_i into corresponding syntactical formulas, one arrives at the famous translation function from L_i -assertions into S4-assertions ([55], 265):

(8) Translation (τ) from L_i to S4:

$$\begin{aligned} \tau(p) &= \Box p \text{ (for all propositional atoms } p \in \rho) \\ \tau(\neg A) &= \Box \neg \tau(A) \\ \tau(A \wedge B) &= \tau(A) \wedge \tau(B) \\ \tau(A \vee B) &= \tau(A) \vee \tau(B) \\ \tau(A \rightarrow B) &= \Box(\tau(A) \rightarrow \tau(B)) \end{aligned}$$

The translation function (8) can be compactified as follows: $\tau(A)$ results from boxing all subformulas of A that are either atoms, negations or implications. For example, $\tau(\neg p) = \Box \neg \Box p = \Box \Diamond \neg p$, $\tau(p \vee q) = \Box p \vee \Box q$, $\tau(p \rightarrow q) = \Box(\Box p \rightarrow \Box q)$, and $\tau((p \rightarrow \neg q) \wedge r) =$

¹¹ Kreisel [38] showed that there is no purely intuitionistic completeness proof for intuitionistic predicate logic (cf. [47] s. 5.1).

$\Box(\Box p \rightarrow \Box\neg\Box q)\wedge\Box r$). With this translation function the following translation theorem has been proved [46]:

$$(9) \textbf{Theorem}(L_i \text{ into } S4): \Gamma \models_i A \text{ iff } \tau(\Gamma) \models_{S4} \tau(A).$$

For example, $\models_{S4} \tau(p\vee\neg p)$ holds, since $\tau(p\vee\neg p) = p \vee \neg p$ and $\models_{S4} p \vee \neg p$.

The major philosophical conclusion of this translation of L_i into S4 is that the logic of mathematical verification may also be represented *within* an extended classical logic. This seems to refute Bueno’s thesis that “if we want to accommodate the constructive features of mathematical reasoning, classical logic will be clearly inadequate” ([10], 108).

5 Translating Quantum Logics

5.1 Quantum-Logical Disjunction and Negation

Quantum logic intends to explicate the structure of ‘experimental’ propositions about quantum-physical systems, having the form “the magnitude m of system x has value v ”, in short “ $m(x) = v$ ” (cf. [69]). For example, $m(x)$ can be the (real-valued) position or the (discrete-valued) spin of an electron x . The system x is described by its physical state or wave-function $\psi(x)$ (the set of all possible states is called ‘Hilbert space’). Each magnitude m corresponds to an operator M describing M ’s measurement whose application to $\psi(x)$ determines the value of $m(x)$ either (i) *sharply*, in which case $\psi(x)$ is a *pure* state w.r.t. the magnitude m and m is an ‘eigenvalue’ of M (i.e., $M\psi(x) = m\cdot\psi(x)$ holds), or (ii) merely *probabilistically*, in which case $\psi(x)$ is in a so-called superposition (or ‘entangled’) state w.r.t. m , meaning that ψ is a linear combination of two or more pure states w.r.t. m . In this case a *measurement* of m changes the systems’s state from a superposition state to a pure state for one of the possible values of m according to a probability distribution derivable from computing $M\psi$; this measurement-induced change is also called the ‘collapse’ of the wave function.

Importantly, certain pairs of quantum-physical magnitudes cannot simultaneously have a sharp value. For example, according to Heisenberg’s uncertainty relation it is impossible that both the position s and the momentum p can both be sharply realized: thus if $\psi(x)$ is a pure state w.r.t. p , it is a superposition state w.r.t. s . It is the crucial feature of the standard version of quantum logic (QL) going back to Birkhoff and von Neumann [6] that the superposition state of a system is expressed as a non-classical disjunction (\vee_q) of pure states. Thus, if the electron in a two-slit experiment is in a superposition of the two possible positions states s_r (going though the right slit) and s_l (going to the left slit), then QL describes this state as $s_r \vee_q s_l$. Note that this view of the matter is by no means mandatory: already early critics of QL objected that a superposition state is genuinely different from a (classical) disjunction of pure states ([49], 685; [22], 523). The defenders of QL replied that quantum disjunction is different from classical disjunction (\vee) and that the description of superposition states by non-classical disjunctions has advantages for realism [54] as well as for empiricism [69].

Instead of entering this controversy we simply observe that the resulting quantum-logical disjunction is radically different from classical disjunction. Assume that the electron in the two-slit experiment is in the superposition state $s_r \vee_q s_l$ and in the pure state p w.r.t. momentum, then QL describes this situation as $p \wedge (s_r \vee_q s_l)$, but by the incommensurability of momentum and position, both conjunctions $p \wedge s_r$ and $p \wedge s_l$ are false. Thus the expansion direction of the classical *law of distribution*, (D_c) : $p \wedge (q \vee r) \rightarrow (p \wedge q) \vee (p \wedge r)$, is invalid according to QL.

Putnam [54] argued that the failure of (D_c) would allow for a *fully* realistic interpretation of quantum physics, because we could now say that the electron is always in a definite position and momentum state; so the only mistake is accepting the logical law of distribution (D_c) . But as several authors (e.g., [16]) pointed out, this argument is flawed, because although the quantum-logical disjunction $s_r \vee_q s_l$ is true in the superposed state, *both* disjuncts of the disjunction are false ([69], 584; [13], 133). Thus quantum-logical disjunction \vee_q is so radically different from classical disjunction that it may be better not to call it a disjunction at all [34]. The reason why many QL-laws - all except of (D_{exp}) - resemble the laws of L_2 is that QL introduces a corresponding quantum-logic *negation* \neg_q that is likewise substantially different from classical negation \neg , but fits with \vee_q in similar ways as \neg fits with \vee . Let Σ be the Hilbert space (of a given quantum-physical system x) and the semantical extension $\|A\| \subseteq \Sigma$ of a quantum proposition A be defined as the set of states in Σ verifying A . Then the extension of the quantum negation $\neg_q A$ is defined as the set of all states that are *orthogonal* to all states verifying A (where two state vectors α, β are orthogonal, in short: $\alpha \perp \beta$, if their scalar product is zero). Thus $\|\neg_q A\|$ is a proper subset of the extension $\Sigma - \|A\|$ of classical negation, and $\|(A \vee_q B)\|$ is a proper superset of the extension $\|A\| \cup \|B\|$ of classical disjunction. Only the QL-extension of the conjunction \wedge is the classical intersection, $\|A \wedge B\| = \|A\| \cap \|B\|$. As a consequence, the extensions of QL-propositions are always closed under linear combinations.

Because of these differences, it is broadly agreed that QL cannot replace classical logic: classical logic is needed to express the full structure and consequences of quantum physics ([65, 29, 3]). On the other hand, quantum logic can express physical operations (e.g., linear combinations) that go beyond what classical logic can express. Moreover, there is no ordinary quantum implication ([13], 146–152). What is needed is a unified system containing the resources of both classical and quantum logic. Such a unification is possible by translating QL into an extension of classical logic, which is the topic of the next section.

5.2 Two Possibilities of Translating Quantum Logics

There are two well-established methods of translating QL into classical logic. The first method translates *orthologic* OL, which is a weakening of QL, into the modal logic B. This modal logic is semantically characterized by reflexive and symmetric Kripke frames and satisfies the axiom (B): $\Box A \rightarrow \Diamond A$ (corresponding to the symmetry of R) as well as (T), (K) and the rule (N) (see sec. 4.2). OL results from classical logic by removing the expansion version of distribution (D_c) ; algebraically OL is characterized by the class of ortholattices (Boolean lattices without distributivity, cf. [13], 137). QL strengthens OL by a weakened distribution law satisfied by the lattice of quantum propositions, namely ‘orthomodularity’ (O): $p \wedge (\neg_q p \vee_q (p \wedge q)) \equiv_{QL} q$ (ibid., 142).

Dishkant [15] proved that OL is complete for the class of generalized KB-frames $\langle W, R, \Pi \rangle$ whose worlds are physical states, whose accessibility relation is interpreted

as the negation of orthogonality ($R\alpha\beta$ iff not $\alpha\perp\beta$), and whose algebra of admissible propositions $\Pi \subseteq \text{Pow}(W)$ contains \emptyset and W , is closed under the operation of orthocomplement, $X^{oc} =_{\text{def}} \{\alpha \in W : \alpha \perp X\}$ (the semantic counterpart of quantum negation), and satisfies the following condition:

$$(*) \forall \beta \in R_\alpha \exists \gamma \in R_\beta (\gamma \in X) \rightarrow \alpha \in X, \text{ for all } X \in \Pi$$

where R_δ is the set of worlds R -accessible by world δ ([13], 139). Condition (*) guarantees closure of worlds under linear combinations and implies $X = (X^{oc})^{oc}$. Generalized Kripke frames of this kind (with $\text{val}(\neg_q A) = \|A\|^{oc}$ and $\text{val}(A \wedge B) = \text{val}(A) \cap \text{val}(B)$) are called *orthoframes*. Following from (*) one can prove that a statement A is true at a world α in an orthoframe F if and only if $\Box A$ is true at α in F (ibid. 140, theorem 11), where the *only-if* direction is the non-trivial part (the if-direction is an instance of axiom B). This fact leads directly to the method of translating OL-statements into the logic KB, characterized by reflexive-symmetric Kripke frames with the classical (i.e. unrestricted) algebra of propositions ($\Pi = \text{Pow}(W)$), namely by translating OL-statements into their ‘necessarily-possibly’-variants as follows:

(10) Translation (τ) from OL into B:

$$\tau(p) = \Box \Diamond p \text{ (for all propositional atoms } p \in \wp)$$

$$\tau(\neg_q A) = \Box \neg \tau(A)$$

$$\tau(A \wedge B) = \tau(A) \wedge \tau(B)$$

$$\tau(A \vee_q B) =_{\text{def}} \tau(\neg_q(\neg_q A \wedge \neg_q B)) = \Box \neg (\Box \neg \tau(A) \wedge \Box \neg \tau(B)) \leftrightarrow \Box (\Diamond \tau(A) \vee \Diamond \tau(B)).$$

Based on this translation one can prove ([24, 13], 156, theorem 53):

(11) **Theorem** (OL into B): $\Gamma \models_{OL} A$ iff $\tau(\Gamma) \models_B \tau(A)$.

The second translation method does not only apply to OL but to full QL and is the most powerful one. It was developed by theoretical physicists [14, 18] and logicians [2, 3] and translates quantum logic into *dynamic* modal logic. This modal logic (abbreviated DML) extends the language of ordinary modal logic by a set of terms a_1, a_2, \dots denoting actions. Each action a_i is semantically characterized by an accessibility relation R_{a_i} between worlds, expressing that performing action a_i leads to a *transition* from the actual state α to an R_{a_i} -accessible state β . For each action term a , $[a]$ is the corresponding action-necessity operator satisfying the standard axioms and rules K and N of Kripke modal logics; $\alpha \models [a]A$ expresses “ A holds in all states produced by action a applied to state α ” (and dually for $\langle a \rangle A$). DML introduces three sorts of complex action terms: “ $a;b$ ” for the sequential combination of a and b , definable as $[a;b]A \leftrightarrow [a][b]A$, “ $a \cup b$ ” for indeterministic choice, definable as $[a \cup b]A \leftrightarrow [a]A \wedge [b]A$, and the classical test operation $?A$ (“is A true?”), characterized by reflexive or empty actions (depending on whether A is true or false) and definable as $[?A]B \leftrightarrow (A \rightarrow B)$. Fine and Schurz [] proved that DML (without the loop-operator) is a definitional extension of a combined multimodal logic.

Dynamical logic is also suited to describe quantum measurements, which may *change* the underlying state if it is a superposed state. Dynamic quantum logic, in short DQL, arises

from DML by leaving out the definition of the classical test operation and assuming instead two types of quantum actions: (i) quantum test operations $A_1?, A_2?, \dots$ for experimental propositions A_i and (ii) unitary actions a_1, a_2, \dots , describing undisturbed evolutions of nonstationary quantum systems (the latter actions are not needed for translating QL into DQL). Experimental quantum propositions A are now translated as follows. “ A holds in a state α ” means that all A -tests applied to state α lead to a state in which A is true, which (as explained) can only hold if α is a pure A -state and $A?$ applied to α leads back to α ([3], 297). “ $\neg_q A$ holds in a state α ” means that no A -test applied to α leads to a positive result, i.e. every A -test fails. This is expressed in DQL as $[A?]_{\perp}$, i.e., the test operation $A?$ leads to nowhere (ibid. 298). Note that $[A?]_{\perp}$ is stronger than the classical negation “ $\neg A$ ” expressing that not all tests of A fail; moreover it satisfies the quantum law of double negation: $\alpha \models \neg_q \neg_q A$ iff all $([A?]_{\perp})$ -tests applied to α fail iff $\alpha \models A$. Conjunction is interpreted classically: “ $A \wedge B$ ” holds in α iff both A and B hold. With the mentioned definition of the indeterministic choice, the test condition for a conjunction, $(A \wedge B)?$, can be expressed as $[\neg_q A? \cup \neg_q B?]_{\perp}$, since $[\neg_q A?]_{\perp} \wedge [\neg_q B?]_{\perp} \equiv \neg_q \neg_q A \wedge \neg_q \neg_q B \equiv A \wedge B$. The QL-disjunction is defined as $A \vee_q B =_{\text{def}} \neg_q (\neg_q A \wedge \neg_q B)$, being equivalent with $[(\neg_q A?)_{\perp} \wedge (\neg_q B?)_{\perp}]?_{\perp}$. The assertions of standard quantum logic can now be translated into DQL as follows:

(12) Translation (τ) from QL into DQL:

$$\begin{aligned} \tau(p) &= p \\ \tau(\neg_q A) &= [\tau(A)?]_{\perp} \\ \tau(A \wedge B) &= \tau(A) \wedge \tau(B) \\ \tau(A \vee_q B) &= \tau(\neg_q (\neg_q A \wedge \neg_q B)), \text{DQL-equivalent with } [([\tau(A)?]_{\perp} \wedge [\tau(B)?]_{\perp})?]_{\perp}. \end{aligned}$$

With this translation one can show ([3], 301):

(13) **Theorem** (from QL into DQL): $\Gamma \models_{\text{QL}} A$ iff $\tau(\Gamma) \models_{\text{DQL}} \tau(A)$.

Thus QL’s non-classical operators \neg_q and \vee_q can be fully transformed into classical assertions about the results of dynamic quantum test operations. Since DQL contains the full resources of classical logic, it fits precisely our goal of offering a unified framework for representing the classical connectives as well as quantum operations. For example, in DQL one can express that a state α is superposed w.r.t. proposition A via $\alpha \models \neg[?A]A \wedge \neg[?A]_{\perp}$. Finally we mention that an equivalent translation of QL is also possible into predicate logic instead of DQL, since it is well known from modal correspondence theory that every modal formula can be equivalently translated into a formula of first or higher order predicate logic (cf. [57], sec. 2.3).

6 Epistemological Conclusions and Discussion of Possible Objections

We have shown that major kinds of non-classical logics are translatable into extended classical logic. We even conjecture that *all* non-classical logics can be translated into a suitably extended classical logic. Our reason for this conjecture is that all non-classical

logics known to us can be semantically described in a classical ML. Further tests of this ‘bold’ conjecture are left to future work.

If our arguments are correct, they give us an *optimality justification* of classical logic: By using classical logic our conceptual representation system can only *gain*, but not *lose*, because if another logic has advantages for certain purposes, we can embed it into classical logic. What is primarily achieved by this result is the removal of the first threat mentioned in section 2: the threat of epistemic *incommensurability*. We rather have a situation of commensurability and ‘pluralism in harmony’. In this harmonic logical pluralism, classical logic plays the role of a *global* unifying framework. One may counter that if classical logic were the only logic that could play this role, this would not speak for logical *pluralism*, but for logical *monism*. However, even if this were the case (see below), it would give us only a representational monism, but not a metaphysical monism, since it would still leave us the options of describing a particular domain by means of determinate or indeterminate, consistent or dialethic propositions (etc.), just that these options can now be realized within the classical framework. This result has significant consequences for the controversy on logical pluralism. For example, it makes it possible to overcome the restrictions of contextualism, insofar as non-classical and classical assertions, that according to contextualism can only be asserted in different contexts (cf. [11]), can now be simultaneously expressed in a unified ‘context’. For example, in $L_{2,L_{uk}}$ we can express that a proposition p about the position-value of an electron in a superposition state is undetermined from a three-valued perspective but false from a two-valued perspective: $U^3p \wedge F^2p$. Assertions of this sort are of obvious importance for rational deliberations about which ‘context’ one should assume.

We now turn to consider possible limitations of our results. Let us first ask: Can the role of a unifying framework only be played by classical logic? This question leads us to the first of two possible objections considered in this final section.

6.1 First Objection: The Possibility of Inverse Translation - Optimality without Dominance?

The first objection points out that there may also be an inverse translation of classical logic into a non-classical logic. Before we investigate this question, let us clarify the possible reach of this objection: if it were true, it would not refute the optimality thesis, but merely the *dominance* thesis for classical logic.

The two notions come from *game theory* (cf. [60], sec. 5.7). A method (here a logic) is optimal in a given class of methods M iff it has maximal value in M in regard to the given goal, here linguistic representation power. Universal optimality coincides with optimality in the class of all methods. The optimality of a logic L does not exclude the possibility of other logics that have the same maximal value; thus there may be *many* logics which are equally optimal. This situation is only excluded if a logic L is dominant, which means that L has a higher value than all other logics.

Although optimality is not refuted by the intertranslatability objection, the objection is not fully defeated either. Our opponent may continue: if some non-classical logics are also optimal, what has been achieved by the optimality justification of classical logic? The answer was given above: the first threat, epistemic incommensurability, is defeated by optimality, but not the second threat, subjective *relativism*. If there are *many* equally representation-optimal logics L_1, \dots, L_n , and there are no further *ceteris paribus*

preferences between them, then the rational choice among them is still undetermined and relative to subjective purpose or taste. As argued in section 2, given translatability, this sort of relativism would not be so disastrous, as we can always intertranslate - yet, given what we said in the preceding section, we would expect that some dominance arguments in favor of classical logic could also be given. Before we turn to this point (in section 6.3) we consider the possibilities of inverse translations in more detail.

Consider our paradigm example in section 3.1, the translation of Lukasiewicz L_3 into classical L_2 . An inverse translation $\tau_{2 \rightarrow 3}$ is possible in complete analogy to $\tau_{3 \rightarrow 2}$ by extending \mathcal{L}_3 by two operators T^2 and F^2 of two-valued logics with the obvious truth tables in L_2 (we write t^2, f^2 to distinguish two-valued from three-valued values):

p	T ² (p)	F ² (p)
t ²	t ²	f ²
f ²	f ²	t ²

In L_2 the two operators are explicitly definable as $T_2A \leftrightarrow A$ and $F_2A \leftrightarrow \neg A$. In L_3 , T^2 and F^2 figure as intensional operators axiomatized by the following set of axioms Ax_{Class} in the extended language $\mathcal{L}_{3,Class}$:

$$\begin{array}{ll}
 T^2(\neg A) \leftrightarrow F^2(A) & F^2(\neg A) \leftrightarrow T^2(A) \\
 T^2(A \wedge B) \leftrightarrow T^2(A) \wedge T^2(B) & F^2(A \wedge B) \leftrightarrow F^2(A) \vee F^2(B) \\
 T^2(A \vee B) \leftrightarrow T^2(A) \vee T^2(B) & F^2(A \vee B) \leftrightarrow F^2(A) \wedge F^2(B) \\
 T^2(A \rightarrow B) \leftrightarrow F^2(A) \vee T^2(B) & F^2(A \rightarrow B) \leftrightarrow T^2(A) \wedge F^2(B) \\
 T^2(A) \dot{\vee} F^2(A) &
 \end{array}$$

Note that from $T^2(A) \vee F^2(A)$ and $F^2(A) \rightarrow \neg T^2(A)$ (following from $T^2(A) \dot{\vee} F^2(A)$) we can derive $T^2(A) \vee \neg T^2(A)$ within $L_3 \cup Ax_{Class}$; thus the T^2 - and F^2 -modalized formulas of $L_{3,Class}$ are bivalent and satisfy the law of excluded middle.

The translation of \mathcal{L}_2 - into \mathcal{L}_3 -formulas is again based on the *truth view* of assertion (which is adequate for all many-valued logics; recall sec. 3.3). Thus we translate:

$$(14) \quad \tau_{2 \rightarrow 3}(A) = T^2(A), \text{ for every } A \in \mathcal{L}_2.$$

Some examples: $\tau(p) = T^2p, \tau(\neg p) = F^2p, \tau(p \vee \neg p) = T^2(p \vee \neg p) \leftrightarrow T^2p \vee F^2p, \tau(p \wedge \neg p) = T^2(p \wedge \neg p) \leftrightarrow T^2p \wedge F^2p, \tau(p \rightarrow q) = T^2(p \rightarrow q) \leftrightarrow F^2p \vee T^2q$, etc. The counterexample to conditional proof in $L_3 - p \wedge \neg p \models q$ but $\not\models (p \wedge \neg p) \rightarrow q$ - fails for the L_3 -translation of L_2 because $Ax_{Class} \models T^2(p \wedge \neg p) \rightarrow T^2(q)$.

More generally, we can prove the following translation theorem in complete analogy to theorem (4) of section 3.1:

(15) **Theorem** (L_2 into L_3):

(15.1) Every bivalent valuation function val_2 over \wp corresponds exactly to a valuation function val_3 over $L_{3,Class,\tau}$ that satisfies the axioms Ax_{Class} such that for every \mathcal{L}_2 -formula A :

$$\text{val}_2(A) = o^2 \text{ iff } \text{val}_3(O^2(A)) = t, \text{ where } o^2 = t^2/f^2 \text{ iff } O^2 = T^2/F^2.$$

$$(15.2) \Gamma \models_{L_2} A \text{ iff } A_{X_{\text{Class}}} \cup T^2(\Gamma) \models_{L_3} T^2(A).$$

For the sake of the semantic coherence between L_2 and L_3 we require again two bridge axioms, but now we must be careful. Intuitively the bridge axioms should say that if an atomic sentence p is trivalently true (resp. false), then it is also bivalently true (resp. false). However, we cannot express the bridge axioms as (i) $p \rightarrow T^2(p)$ and (ii) $\neg p \rightarrow F^2(p)$ (like we did this in (5) of section 3.2, with $T = T^3$ instead of T^2 , and likewise for F), because in L_3 , p and $T^3(p)$, are not equivalent. Implications (i) and (ii) would lead to failure when we contrapose (ii), obtaining $\neg F^2(p) \rightarrow p$, which together with $T^2(p) \rightarrow \neg F^2(p)$ would imply (iii) $T^2(p) \rightarrow p$; but (iii) and (i) would entail $p \leftrightarrow T^2(p)$, i.e., the ‘collapse’ of L_3 into L_2 . Therefore we have to explicate the two bridge axioms as (i*) $T^3(p) \rightarrow T^2(p)$ and (ii*) $F^3(p) \rightarrow F^2(p)$, where T^3 and F^3 are defined in L_3 as explained in sec. 3.1 (we now write T^3 and F^3 for T and F of sec. 3.1, to desambiguate them from T^2 and F^2).

A similar translation of L_2 is possible for all n -valued logics that have “truth” as a designated value and within which the operations of n -valued truth and falsity are definable. For many other non-classical logics, however, it is impossible to embed classical logic via translation. There are two possible reasons for the impossibility of such a translation. *First*, the non-classical logic may be too weak; thus the translation is technically impossible. *Second*, the translation would be technically possible, but is philosophically inadequate. While the first case applies to paraconsistent and to quantum logic, the second case applies to intuitionistic logic.

For the system LP of paraconsistent logic the translation method fails, because LP’s designated values are T^3 and B^3 , and as explained in section 4.1, it is impossible to define truth-only (T^3) in LP. Therefore the axiom $T^2(A) \vee F^2(A)$ (together with $T^2(\neg A) \leftrightarrow F^2(A)$ and $F^2(\neg A) \leftrightarrow F^2(A)$) cannot enforce in LP that $T^2(A)$ and $F^2(A)$ are bivalent (either true-only or false-only); the designatedness of $T^2(A) \vee F^2(A)$ is compatible with $T^2(A)$ and $F^2(A)$ having the value b . Even stronger, there is no formula $X(p)$ of LP whose designatedness could entail that p is true-only. Thus the representational power of LP is strictly weaker than L_2 or L_3 .¹²

In intuitionistic logic, the translation of L_2 into L_i could be technically carried out in the same way as for L_3 , by extending the intuitionistic language \mathcal{L}_i by the bivalent operators T^2 , F^2 and adding the axiom set $A_{X_{\text{Class}}}$. The bridge axioms can be expressed as $A \rightarrow T^2(A)$ and $\neg A \rightarrow F^2(A)$ without entailing collapse (because $\neg\neg A$ does not entail A). Thus the translation is technically possible; however, it is philosophically inadequate. The translation would allow one to derive the bivalence axiom $T^2(A) \vee \neg T^2(A)$ for L_2 -translations, but L_i ’s assertions are based on the proof view of assertion. According to this view the validity of $T^2(A) \vee \neg T^2(A)$ in L_i means that one can prove this disjunction and thus can prove either $T^2(A)$ or $\neg T^2(A)$, for every formula A . But this is against the spirit of intuitionistic logic which admits propositions whose truth value is not decidable by a constructive proof.

A translation of classical logic into quantum logic QL appears also to be impossible. Indeed, since the implicit semantics of QL is bivalent, the introduction of an intensional truth operation T^2 would be pointless, because it would satisfy the correspondence

¹² Interestingly a similar result for LP was achieved by Jerábek, ([35], 677), though for very different translation functions (recall sec. 3.2.3).

axiom $T^2(A) \leftrightarrow A$. What would be needed is a translation of classical negation and disjunction, $\tau(\neg A)$ and $\tau(A \vee B)$, into QL-formulas that behave like the classical connectives, but such formulas do not exist.

6.2 Second Objection - the Metalogic of Translation Theorems

The proof of the translation theorems in sections 3–4 is performed by using a classical metalogic. As mentioned above, not all non-classical logicians allow classical logic in their meta-language. Some non-classical logicians insist on level-coherence and use their non-classical logic also at the meta-level. Such a ‘through-and-through’ proponent of a non-classical logic could object that the translation theorems have not been proved in the ‘right’ logic. Does this problem constitute an ultimate circularity within our account? A comprehensive discussion of this question is not possible here. Instead, we confine ourselves to two replies to this challenge: the first reply is a weakness argument and the second is a dialectic refutation strategy.

6.2.1 The Weakness Argument

The metalogical proof of the translation theorem in sec. 3.1 uses only *very weak* logical rules. Since the axioms Ax_{Luk} are valid in both L_2 and L_3 , what is needed in the proof of lemma 2 is the rule of *replacement* of logical equivalents $\Gamma \models A \leftrightarrow B / \Gamma \models C \leftrightarrow C[B/A]$ (that holds in L_2 and L_3). Beyond that, the proof uses only finitary set theory and the classical max-rule and min-rule for the designated values of conjunction and disjunction, which hold in all multi-valued logics. Thus the metalogical proof of the translation theorem of section 3.1 should be acceptable in all multi-valued logics. Similar arguments may be applied to other translation theorems.

6.2.2 The Dialectic Refutation Strategy

It is not clear whether the weakness argument works for all other translation theorems. For example, certain hyperintensional logics [42] do not accept the rule of replacing equivalents. So let us assume that the proponent of a non-classical logic L^* rejects the metalogical proof of the translation theorem of L^* into classical logic L_2 . Even if this is the case, we can still apply the following dialectic refutation strategy:

There are two possibilities:

- (i) Either L_2 is *re-translatable* into L^* in some way that is acceptable for proponent of L^* . In this case the metalogical proof of the translation theorem of L^* into L_2 can be translated from the L_2 -metalogic into the L^* -metalogic and the L^* -proponent has to accept it.
- (ii) Or L_2 is *not re-translatable* into L^* in some way that is acceptable for the proponent of L^* . In this case, L^* is not representationally optimal from L^* 's own point of view, while L_2 is representationally optimal from L_2 's point of view. We argue that such a situation constitutes a reason to prefer L_2 over L^* , because *if* one uses the resources of L_2 , then provably everything that can be expressed and derived in L^* can also be expressed and derived in L_2 , while this is not so if one uses the resources of L^* .

6.3 From Optimality to Dominance

6.3.1 Global *ceteris paribus* advantages of classical logics

In sec. 6.2 we have seen that L_3 and some other many-valued logics are intertranslatable with L_2 . This fact shows that L_2 is not the only representationally optimal logic; some other many-valued logics are representationally optimal, too. In this section we argue that the situation is nevertheless not symmetric. Classical logic has two global *ceteris paribus* advantages as a unifying metalogic:

- (i) *Determinateness in the metalogic*: We are interested in descriptions that are as informative and thus as determinate as possible. Therefore we are interested in constructing logical frameworks whose syntactic and semantic properties are determined. We do not want a notion of proof for which the question of whether a conclusion follows from a premise set is irreducibly indeterminate, nor are we interested in a logical system whose consistency is indeterminate (etc.). This speaks in favor of using not a trivalent, but a classical bivalent logic as metalogic. Should there really arise a situation where we encounter an irreducible indeterminateness, we can express this by using intensional operators.
- (ii) *Naturalness and induction from practice*: It is common practice to discuss the pros and cons of competing logics within a classical metalogic. Likewise it is common practice in mathematical logic to use classical metalogic for the semantics of a non-classical logic. These facts of our practice don't 'prove' anything, and there are exceptions, e.g., constructivist mathematicians using an intuitionistic metalogic. However, these facts seem to confirm the thesis that a classical metalogic is the most natural choice of a framework for the comparison and evaluation of competing systems of logic.

6.3.2 Local Advantages of Classical and Non-Classical Logics

A further important *ceteris paribus* criterion for choice of logics is *simplicity* (cf. [71]). The simplicity of a logic has many dimensions, concerning the simplicity of its expressions, its proof system, its semantics, as well as the simplicity of the translations by which other logics can be embedded into it. What follows from the simplicity criterion for the choice of a logic? Assume we are in a domain whose description requires the logical operations of a non-classical logic L_n . Then the question is: what is simpler - using L_n directly or using the translation of L_n into $L_{2,n}$? This depends on the context of application. As a first example, consider quantum logic. The logical representation of linear combinations and orthocomplements of quantum-physical states is simpler in QL than in QL's translation into classical logic ([1], 185). On the other hand, if we want to express that a quantum system is in a pure versus a superposed state, we need in addition the resources of classical logic (recall sec. 5.2), which speaks in favour of using QL's translation into classical logic. As a second example, consider Lukasiewicz' three-valued logic. Assume we have good reasons for thinking that certain propositions describing our domain are undetermined. If there are only a few propositions of this sort and most propositions are adequately described in L_2 , then using the extended logic $L_{2,Luk}$ is clearly preferable to a complicated 'context-switching' between

L_3 and L_2 . On the other hand, in a domain where more-or-less all propositions are three-valued (if such a domain exists), the employment of L_3 throughout would be simpler than using L_3 's translation in $L_{2,Luk}$.

6.3.3 Summary of conclusions

In this paper we have demonstrated that four major systems of non-classical logic - multi-valued, intuitionistic, paraconsistent and quantum logics - can be translated into classical logic in a meaning-preserving way, by introducing additional intensional operators into the language. Two major philosophical conclusions were drawn from this result: First, non-classical logics are not incommensurable with classical logic, but can be semantically embedded into it. Second, classical logic is representationally optimal, since if a non-classical logic turns out to have advantages for particular purposes, it can be translated into classical logic. In the last section we investigated the possibility of inverse translations. We have seen that not many, but at least some non-classical logics, e.g. Lukasiewicz' three-valued logic, can be likewise representationally optimal. Despite this fact we have argued that the situation is not symmetric. Classical logic has global *ceteris paribus* advantages as a unifying metalogic, and (in many but not in all domains) local advantages as a most simple object logic. One may ask: Why didn't we bring in these advantages of classical logic from the start? The reason is that without the intertranslatability argument, the incommensurability problem would prevent that these advantages become justificatorily crucial. A proponent of a non-classical logic L^* could insist that L^* is the correct logic, and correctness is more important than these *ceteris paribus* advantages. In combination with translatability, however, these advantages may become decisive, because the proponent of L^* can reformulate L^* in the classical framework L_2 , where (s)he can implement the metaphysics of L^* while enjoying the *ceteris paribus* advantages of L_2 .

Acknowledgements For important discussion and help I am indebted to Heinrich Wansing, Diderik Batens, Graham Priest, Manuel Bremer, Hannes Leitgeb, Alexandre Costa-Leite, Jean-Yves Beziau, Corina Strößner and two unknown referees.

Funding Open Access funding enabled and organized by Projekt DEAL. This work was supported by the DFG (Deutsche Forschungsgemeinschaft), research unit FOR 2495.

Declarations

Conflict of Interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ashcroft, M. (2010). Does science influence the logic we ought to use: A reflection on the quantum logic controversy. *Studia Logica*, 95, 183–206.
2. Baltag, A., & Smets, S. (2005). Complete axiomatizations for quantum actions. *International Journal of Theoretical Physics*, 44(12), 2267–2282.
3. Baltag, A., & Smets, S. (2011). Quantum logic as a dynamic logic. *Synthese*, 179(2), 285–306.
4. Beall, J. C., & Restall, G. (2006). *Logical pluralism*. Oxford Univ. Press.
5. Beziau, J.-Y. (1999). A sequent calculus for Lukasiewicz's three-valued logic based on Susko's bivalent semantics. *Bulletin of the Section of Logic*, 28(2), 89–97.
6. Birkhoff, G., & von Neumann, J. (1936). The logic of quantum mechanics. *Annals of Mathematics*, 37, 823–843.
7. Bonjour, L. (1985). *The structure of empirical knowledge*. Harvard Univ. Press.
8. Bremer, M. (2014). Restall and Beall on logical pluralism: A critique. *Erkenntnis*, 79, 293–299.
9. Brouwer, L. E. J. (1912). Intuitionism and formalism. Reprinted in Benacerraf & Putnam (eds.), *Philosophy of mathematics: selected readings* (pp. 77–89). Cambridge University Press, 2nd ed., 1983.
10. Bueno, O. (2010). Is logic a priori? *The Harvard Review of Philosophy*, 17, 105–117.
11. Caret, C. R. (2017). The collapse of logical pluralism has been greatly exaggerated. *Erkenntnis*, 82, 739–760.
12. Camielli, W. A., Coniglio, M. E., & D'Ottavio, I. M. E. (2009). New dimensions on translations between logics. *Logica Universalis*, 3, 1–18.
13. Dalla Chiara, M. L., & Giuntini, R. (2002). Quantum logics. In D. Gabbay & F. Guenther (Eds.), *Handbook of philosophical logic* (Vol. 6, pp. 129–228). Kluwer.
14. Daniel, W. (1989). Axiomatic description of irreversible and reversible evolution of a physical system. *Helvetica Physica Acta*, 62, 941–968.
15. Dishkant, H. (1972). Semantics of the minimal logic of quantum mechanics. *Studia Logica*, 30, 17–29.
16. Dummett, M. (1976). Is logic empirical?. Reprinted in M. Dummett, *Truth and Other Enigmas* (pp. 269–289). Harvard University Press, 1978.
17. Dummett, M. (1991). *The logical basis of metaphysics*. Harvard University Press.
18. Faure, C., Moore, D. J., & Piron, C. (1995). Deterministic evolution and Schroedinger flows. *Helvetica Physica Acta*, 68, 150–157.
19. Feldbacher, C. (forthcoming). *Epistemic engineering*. Rowman & Littlefield.
20. Field, H. (1996). The apriority of logic. *Proceedings of the Aristotelian Society*, 96, 59–379.
21. Fine, K., & Schurz, G. (1996). Transfer theorems for multimodal logics. In Copeland, J. (ed.), *Logic and reality. Essays on the legacy of Arthur Prior* (pp. 169–213). Cambridge Univ. Press.
22. Gardner, M. (1972). Is quantum logic really logic? *Philosophy of Science*, 38, 508–529.
23. Gödel, K. (1934). Zum intuitionistischen Aussagenkalkül. *Anzeiger der Österreichischen Akademie der Wissenschaften*, 69, 65–66.
24. Goldblatt, R. (1974). Semantical analysis of orthologic. *Journal of Philosophical Logic*, 3, 19–35.
25. Goldman, A. (1999). *Knowledge in a social world*. Oxford Univ. Press.
26. Gottwald, S. (1989). *Mehrwertige Logik*. Berlin; Akademie-Verlag.
27. Greco, J. (2005). Justification is not internal. In M. Steup & E. Sosa (Eds.), *Contemporary debates in epistemology* (pp. 257–270). Blackwell Publishing.
28. Haack, S. (1976). The justification of deduction. *Mind*, 85(337), 112–119.
29. Hellman, G. (1980). "Quantum Logic and Meaning", *PSA 1980* (Proceedings of the Biennial Meeting of the Philosophy of Science Association), 2, 493–511.
30. Henderson, L. (2018). The problem of induction. In *Stanford Encyclopedia of Philosophy* (Summer 2018 Edition), <https://plato.stanford.edu/archives/sum2018/entries/induction-problem/>.
31. Hjortland, O. T. (2013). Logical pluralism, meaning-variance, and verbal disputes. *Australasian Journal of Philosophy*, 91(2), 355–373.
32. Hjortland, O. T. (2017). Anti-exceptionalism about logic. *Philosophical Studies*, 174, 631–658.
33. Horvath, J., & Koch, S. (2021). Experimental philosophy and the method of cases. *Philosophy Compass*, 16, e12716. <https://doi.org/10.1111/phc3.12716>.
34. Jauch, J. M. (1968). *Foundations of quantum mechanics*. Addison-Wesley.
35. Jerábek, E. (2012). The ubiquity of conservative translations. *Review of Symbolic Logic*, 5(4), 666–678.
36. Kleene, S. C. (1952). *Metamathematics*. North Holland.

37. Kooi, B., & Tamminga, A. (2013). Three-valued logics in modal logic. *Studia Logica*, 101, 1061–1072.
38. Kreisel, G. (1958). Elementary completeness properties of intuitionistic logic with a note on negations of prenex formulas. *Journal of Symbolic Logic*, 23, 317–330.
39. Kripke, S. A. (1965). Semantical analysis of intuitionistic logic. In J. Crossley & M. A. E. Dummett (Eds.), *Formal systems and recursive functions* (pp. 92–130). North-Holland Publishing.
40. Kuhn, T.S. (1962). *The structure of scientific revolutions*. Univ. of Chicago press (3rd edition 1996).
41. Lehrer, K. (1990). *Theory of knowledge*. Routledge.
42. Leitgeb, H. (2019). HYPE: A system of hyperintensional logic. *Journal of Philosophical Logic*, 48, 305–405.
43. Lipton, P. (1991). *Inference to the best explanation*. Routledge.
44. Łukasiewicz, J. (1920). O logice trojwartosciowej. *Ruch Filozoficzny*, 5, 170–171. Translated in J. Łukasiewicz (1970), *Selected works*. North-Holland and Warsaw.
45. Malinowski, G. (1993). *Many-valued logics*. Clarendon Press.
46. McKinsey, J., & Tarski, A. (1948). Some theorems about the sentential calculi of Lewis and Heyting. *Journal of Symbolic Logic*, 13, 1–15.
47. Moschovakis, J. (2018). Intuitionistic logic. *Stanford Encyclopedia of Philosophy*, plato.stanford.edu/archives/win2018/entries/logic-intuitionistic.
48. Papineau, D. (1993). *Philosophical naturalism*. B. Blackwell.
49. Popper, K. (1968). Birkhoff and von Neumann's interpretation of quantum Mmechanics. *Nature*, 219, 682–685.
50. Priest, G. (1979). Logic of paradox. *Journal of Philosophical Logic*, 8, 219–241.
51. Priest, G. (2006). *In Contradiction*. Oxford University press (2nd edition).
52. Priest, G. (2013). Paraconsistent logic. In *Stanford Encyclopedia of Philosophy* <http://plato.stanford.edu>.
53. Psillos, S. (1999). *Scientific realism. How science tracks truth*. Routledge.
54. Putnam, H. (1968). Is logic empirical? Reprinted as 'The logic of quantum physics' in H. Putnam, *Philosophical Papers. Vol. 1* (pp. 174–197). Cambridge University Press 1979, 2nd ed.
55. Rautenberg, W. (1979). *Klassische und nichtklassische Aussagenlogik*. Vieweg.
56. Salmon, W. C. (1957). Should we attempt to justify induction? *Philosophical Studies*, 8(3), 45–47.
57. Schurz, G. (2006). Alethic modal logic and semantics. In D. Jacquette (ed.), *A companion to philosophical logic* (pp. 442–477). B. Blackwell, 2nd ed.
58. Schurz, G. (2008). The Meta-inductivist's winning strategy in the prediction game. *Philosophy of Science*, 75, 278–305.
59. Schurz, G. (2018). Optimality justifications: New foundations for foundation-oriented epistemology. *Synthese*, 195, 3877–3897.
60. Schurz, G. (2019). *Hume's problem solved: The optimality of meta-induction*. MIT Press.
61. Schurz, G., & Hertwig, R. (2019). Cognitive success. *Topics in Cognitive Science*, 11(1), 7–36.
62. Sellars, W. (1975). The structure of knowledge. In H.-N. Castaneda (Ed.), *Action, knowledge, and reality* (pp. 295–347). Bobbs-Merrill.
63. Sereni, A., & Sforza Fogliani, M. P. (2020). How to water a thousand flowers. On the logic of logical pluralism. *Inquiry*, 63(3–4).
64. Sorensen, R. (2018). Vaguenes. In *The Stanford Encyclopedia of Philosophy* (Summer 2018 edition), plato.stanford.edu/archives/sum2018/entries/vagueness/.
65. Stachel, J. (1986). Do quanta need a new logic? In R. G. Colodny & A. Coffa (Eds.), *From quarks to quasars: Philosophical problems of modern physics* (pp. 229–347). University of Pittsburgh Press.
66. Sterkenburg, T. (2019). The meta-inductive justification of induction. *Philosophy of Science*, 86, 981–992.
67. Suszko, R. (1977). The Fregean axiom and polish mathematical logic in the 1920's. *Studia Logica*, 36, 373–380.
68. Van Cleve, J. (1984). Reliability, justification, and induction. In P. A. French, T. E. Uehling, & H. K. Wettstein (Eds.), *Causation and causal theories. Midwest studies in philosophy. Vol. 9* (pp. 555–567). Univ. of Minnesota Press.
69. Van Fraassen, B. (1975). The labyrinth of quantum logics. In C. A. Hooker (Ed.), *The logico-algebraic approach to quantum mechanics* (pp. 577–607). Reidel.
70. Wansing, H., & Shramko, Y. (2008). Suszko's thesis, inferential many-valuedness, and the notion of a logical system. *Studia Logica*, 88, 405–429.
71. Williamson, T. (2017). Semantic paradoxes and abductive methodology. In B. Armour-Garb (Ed.), *Reflections on the liar* (pp. 325–346). Oxford University Press.
72. Woodruff, P. M. (1970). Logic and truth value gaps. In K. Lambert (Ed.), *Philosophical problems in logic* (pp. 121–142). Reidel.

73. Woods, J. (2019). Against reflective equilibrium for logical theorizing. *Australasian Journal of Logic*, 16(7), 319–341.
74. Wright, C. (2004). Warrant for nothing (and foundations for free)? *Aristotelean Society Supplementary*, 78(1), 167–212.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.