




Correction: Breastfeeding Experiences in Women from Ten States Reporting Opioid Use Before or During Pregnancy: PRAMS, Phase 8

Madeleine Jade Bremer¹ · Kerri Lynn Knippen² 

Published online: 13 October 2022

© Springer Science+Business Media, LLC, part of Springer Nature 2022

Correction to: Maternal and Child Health Journal
<https://doi.org/10.1007/s10995-022-03497-0>

The original version of this article unfortunately contained a mistake in Table 2. The results and footnotes are missing in Table 2. The correct Table 2 is given below. The original article has been updated.

The original article can be found online at <https://doi.org/10.1007/s10995-022-03497-0>.

✉ Kerri Lynn Knippen
kknippe@bgsu.edu

Madeleine Jade Bremer
maddie.bremer1@gmail.com

¹ Food Service Director and Community Nutrition Educator,
Camp Beechpoint 3212 125th Ave, Allegan, MI, USA

² Department of Public & Allied Health, Bowling Green
State University, 122 Health & Human Services Building,
Bowling Green, OH, USA

Table 2 Breastfeeding attempt and duration among opioid and non-opioid reporting women from 10 states who participated in PRAMS, Phase 8 (2016)

Variable	Non-opioid % ^a	Opioid % ^a	OR ^b	95% CI ^c	P Value	Adj ^d OR	95%CI	P Value
<i>BF attempt</i>	83.8	83.0	0.95	0.71, 1.3	0.71			
<i>BF duration</i>								
≥ 6 weeks	82.7	71.6	0.53	0.39, 0.71	< 0.001	0.61	0.44, 0.84	0.003
≥ 10 weeks	71.0	55.6	0.51	0.39, 0.66	< 0.001	0.62	0.47, 0.82	< 0.001
≥ 20 weeks	14.2	8.7	0.53	0.38, 0.87	< 0.001	0.60	0.38, 0.95	0.03

^aWeighted estimate

^bOR = odds ratio

^cCI = confidence interval

^dConfounders adjusted for included: state, maternal race-ethnicity, income, education, WIC participation during pregnancy, gestational age, delivery, plurality, prior live births, and infant transfer

^eBF = breastfeeding

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.