

A regularity result for the bound states of *N*-body Schrödinger operators: blow-ups and Lie manifolds

Bernd Ammann¹ · Jérémy Mougel² · Victor Nistor^{3,4}

Received: 10 March 2022 / Revised: 27 December 2022 / Accepted: 12 January 2023 / Published online: 21 February 2023 © The Author(s) 2023

Abstract

We prove regularity estimates in weighted Sobolev spaces for the L^2 -eigenfunctions of Schrödinger-type operators whose potentials have inverse square singularities and uniform radial limits at infinity. In particular, the usual *N*-body Hamiltonians with Coulomb-type singular potentials are covered by our result: in that case, the weight is $\delta_{\mathcal{F}}(x) := \min\{d(x, \bigcup \mathcal{F}), 1\}$, where $d(x, \bigcup \mathcal{F})$ is the usual Euclidean distance to the union $\bigcup \mathcal{F}$ of the set of collision planes \mathcal{F} . The proof is based on blow-ups of manifolds with corners and Lie manifolds. More precisely, we start with the radial compactification \overline{X} of the underlying space X and we first blow up the spheres $\mathbb{S}_Y \subset \mathbb{S}_X$ at infinity of the collision planes $Y \in \mathcal{F}$ to obtain the Georgescu–Vasy compactification. Then, we blow up the collision planes \mathcal{F} . We carefully investigate how the Lie manifold structure and the associated data (metric, Sobolev spaces, differential operators) change with each blow-up. Our method applies also to higher-order differential operators, to certain classes of pseudodifferential operators, and to matrices of scalar operators.

Bernd Ammann bernd.ammann@mathematik.uni-regensburg.de

Jérémy Mougel j.mougel@hotmail.com

Victor Nistor victor.nistor@univ-lorraine.fr

- ¹ Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, Germany
- ² Mathematisches Institut Georg-August-Universität Göttingen, 37083 Göttingen, Germany
- ³ Université de Lorraine, CNRS, IECL, 57000 Metz, France
- ⁴ Institute of Mathematics of the Romanian Academy, PO Box 1-764, 014700 Bucharest, Romania

B.A. has been partially supported by SPP 2026 (Geometry at infinity) and the SFB 1085 (Higher Invariants), both funded by the DFG (German Science Foundation). V.N. has been partially supported by ANR-14-CE25-0012-01 (SINGSTAR) funded by ANR (French Science Foundation).

Keywords Schrödinger equation \cdot Regularity \cdot Eigenfunctions \cdot N-body problem \cdot Georgescu-Vasy compactification

Mathematics Subject Classification Primary 35J10; Secondary 35B65 · 35J47 · 58J05

Contents

1.1Basic notation and constructions31.2Statement of the main result and comments41.3Contents of the paper52Preliminaries62.1A few basic notations and definitions62.2Manifolds with corners and their submanifolds72.3Submanifolds of manifolds with corners82.4Clean intersections103Blow-ups113.1The blow-up along a closed p-submanifold113.2Iterated blow-ups133.3Admissible orders163.4A blow-up point of view on the spherical compactification194Distance functions and Sobolev spaces for blown-up spaces204.1Smoothed distance function to a p-submanifold214.2Smoothed distance functions to a family254.3Blow-ups and Lie manifolds285The Lie manifolds associated with the generalized N-body problem335.1.1The semilattices and blow-ups and the data on $X_{\rm GV}$ 355.1.3Interior blow-ups365.1.4Blow-ups and the data on $X_{\rm GV}$ 355.1.3Interior blow-ups396Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{T}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	1	Introduction	2
1.2Statement of the main result and comments41.3Contents of the paper52Preliminaries62.1A few basic notations and definitions62.2Manifolds with corners and their submanifolds72.3Submanifolds of manifolds with corners82.4Clean intersections103Blow-ups113.1The blow-up along a closed p-submanifold113.2Iterated blow-ups133.3Admissible orders163.4A blow-up point of view on the spherical compactification194Distance functions to a family254.1Smoothed distance functions to a family255The Lie manifolds285The Lie manifolds335.1.2Boundary blow-ups of the N-body problem335.1.2Boundary blow-ups and the data on X_{GV} 355.1.3Interior blow-ups365.2A lifting lemma and smoothed distance functions375.3More on smooth distance functions on blow-ups396Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix C. A splitting lemma49References514Broup as on compactifications of vector spaces47Appendix C. A splitting lemma49		1.1 Basic notation and constructions	3
1.3 Contents of the paper52 Preliminaries62.1 A few basic notations and definitions62.2 Manifolds with corners and their submanifolds72.3 Submanifolds of manifolds with corners82.4 Clean intersections103 Blow-ups113.1 The blow-up along a closed p-submanifold113.2 Iterated blow-ups133.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance functions to a family254.2 Smoothed distance functions to a family255 The Lie manifolds285 The Lie manifolds subt the blow-ups335.1.2 Boundary blow-ups of the N-body problem335.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 47Appendix C. A splitting lemma49References51		1.2 Statement of the main result and comments	4
$ \begin{array}{llllllllllllllllllllllllllllllllllll$		1.3 Contents of the paper	5
2.1 A few basic notations and definitions62.2 Manifolds with corners and their submanifolds72.3 Submanifolds of manifolds with corners82.4 Clean intersections103 Blow-ups113.1 The blow-up along a closed p-submanifold113.2 Iterated blow-ups133.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1.1 The semilattices and blow-ups335.1.2 Boundary blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix C. A splitting lemma49References51	2	Preliminaries	6
2.2Manifolds with corners and their submanifolds72.3Submanifolds of manifolds with corners82.4Clean intersections103Blow-ups113.1The blow-up along a closed p-submanifold113.2Iterated blow-ups133.3Admissible orders163.4A blow-up point of view on the spherical compactification194Distance functions and Sobolev spaces for blown-up spaces204.1Smoothed distance function to a p-submanifold214.2Smoothed distance functions to a family254.3Blow-ups and Lie manifolds285The Lie manifolds associated with the generalized N-body problem335.1.1The semilattices and the blow-ups335.1.2Boundary blow-ups and the data on X_{GV} 355.1.3Interior blow-ups365.2A lifting lemma and smoothed distance functions375.3More on smooth distance functions375.4Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix C. A splitting lemma49References51		2.1 A few basic notations and definitions	6
2.3Submanifolds of manifolds with corners82.4Clean intersections103Blow-ups113.1The blow-up along a closed p-submanifold113.2Iterated blow-ups133.3Admissible orders163.4A blow-up point of view on the spherical compactification194Distance functions and Sobolev spaces for blown-up spaces204.1Smoothed distance function to a p-submanifold214.2Smoothed distance functions to a family254.3Blow-ups and Lie manifolds285The Lie manifolds associated with the generalized N-body problem335.1.1The semilattices and the blow-ups335.1.2Boundary blow-ups and the data on X_{GV} 355.1.3Interior blow-ups365.2A lifting lemma and smoothed distance functions375.3More on smooth distance functions on blow-ups396Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		2.2 Manifolds with corners and their submanifolds	7
2.4 Clean intersections103 Blow-ups113.1 The blow-up along a closed p-submanifold113.2 Iterated blow-ups133.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions375.4 Nore on smooth distance functions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		2.3 Submanifolds of manifolds with corners	8
3Blow-ups113.1The blow-up along a closed p-submanifold113.2Iterated blow-ups133.3Admissible orders133.3Admissible orders163.4A blow-up point of view on the spherical compactification194Distance functions and Sobolev spaces for blown-up spaces204.1Smoothed distance function to a p-submanifold214.2Smoothed distance functions to a family254.3Blow-ups and Lie manifolds285The Lie manifolds associated with the generalized N-body problem335.1Semilattices and blow-ups for the N-body problem335.1.2Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3Interior blow-ups365.2A lifting lemma and smoothed distance functions375.3More on smooth distance functions on blow-ups396Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		2.4 Clean intersections	10
3.1 The blow-up along a closed p-submanifold113.2 Iterated blow-ups133.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1 Semilattices and blow-ups for the N-body problem335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	3	Blow-ups	11
3.2 Iterated blow-ups133.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1 Semilattices and blow-ups for the N-body problem335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		3.1 The blow-up along a closed p-submanifold	11
3.3 Admissible orders163.4 A blow-up point of view on the spherical compactification194 Distance functions and Sobolev spaces for blown-up spaces204.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1 Semilattices and blow-ups for the N-body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		3.2 Iterated blow-ups	13
3.4 A blow-up point of view on the spherical compactification 19 4 Distance functions and Sobolev spaces for blown-up spaces 20 4.1 Smoothed distance function to a p-submanifold 21 4.2 Smoothed distance functions to a family 25 4.3 Blow-ups and Lie manifolds 28 5 The Lie manifolds associated with the generalized N-body problem 33 5.1 Semilattices and blow-ups for the N-body problem 33 5.1.1 The semilattices and the blow-ups 33 5.1.2 Boundary blow-ups and the data on X_{GV} 35 5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		3.3 Admissible orders	16
4 Distance functions and Sobolev spaces for blown-up spaces 20 4.1 Smoothed distance function to a p-submanifold 21 4.2 Smoothed distance functions to a family 25 4.3 Blow-ups and Lie manifolds 28 5 The Lie manifolds associated with the generalized N-body problem 33 5.1 Semilattices and blow-ups for the N-body problem 33 5.1.1 The semilattices and the blow-ups 33 5.1.2 Boundary blow-ups and the data on X_{GV} 35 5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		3.4 A blow-up point of view on the spherical compactification	19
4.1 Smoothed distance function to a p-submanifold214.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized N-body problem335.1 Semilattices and blow-ups for the N-body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	4	Distance functions and Sobolev spaces for blown-up spaces	20
4.2 Smoothed distance functions to a family254.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized <i>N</i> -body problem335.1 Semilattices and blow-ups for the <i>N</i> -body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		4.1 Smoothed distance function to a p-submanifold	21
4.3 Blow-ups and Lie manifolds285 The Lie manifolds associated with the generalized <i>N</i> -body problem335.1 Semilattices and blow-ups for the <i>N</i> -body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51		4.2 Smoothed distance functions to a family	25
5 The Lie manifolds associated with the generalized <i>N</i> -body problem 33 5.1 Semilattices and blow-ups for the <i>N</i> -body problem 33 5.1.1 The semilattices and the blow-ups 33 5.1.2 Boundary blow-ups and the data on X_{GV} 35 5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		4.3 Blow-ups and Lie manifolds	28
5.1 Semilattices and blow-ups for the <i>N</i> -body problem335.1.1 The semilattices and the blow-ups335.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 355.1.3 Interior blow-ups365.2 A lifting lemma and smoothed distance functions375.3 More on smooth distance functions on blow-ups396 Regularity results for eigenfunctions40Appendix A. The equivalence of ρ and $\delta \mathcal{F}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	5	The Lie manifolds associated with the generalized <i>N</i> -body problem	33
5.1.1 The semilattices and the blow-ups 33 5.1.2 Boundary blow-ups and the data on X_{GV} 35 5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.1 Semilattices and blow-ups for the <i>N</i> -body problem	33
5.1.2 Boundary blow-ups and the data on $X_{\rm GV}$ 35 5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.1.1 The semilattices and the blow-ups	33
5.1.3 Interior blow-ups 36 5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.1.2 Boundary blow-ups and the data on X_{GV}	35
5.2 A lifting lemma and smoothed distance functions 37 5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.1.3 Interior blow-ups	36
5.3 More on smooth distance functions on blow-ups 39 6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.2 A lifting lemma and smoothed distance functions	37
6 Regularity results for eigenfunctions 40 Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45 Appendix B. Group actions on compactifications of vector spaces 47 Appendix C. A splitting lemma 49 References 51		5.3 More on smooth distance functions on blow-ups	39
Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$ 45Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	6	Regularity results for eigenfunctions	40
Appendix B. Group actions on compactifications of vector spaces47Appendix C. A splitting lemma49References51	A	Appendix A. The equivalence of ρ and $\delta_{\mathcal{F}}$	
Appendix C. A splitting lemma 49 References 51	A	Appendix B. Group actions on compactifications of vector spaces	
References	A	Appendix C. A splitting lemma	
	Re	References	

1 Introduction

We obtain regularity estimates for the eigenfunctions of a class of elliptic differential operators with singular coefficients. Our results cover Schrödinger-type operators whose potentials have inverse square-type singularities and uniform radial limits at infinity. Weaker singularities, such as Coulomb-type singularities, are included in our results; in particular, the "usual" *N*-body Hamiltonians [7, 13, 15] are covered by our results. Our results are obtained by combining and extending the results of our previous papers [1] and [6], which will be two basic references in what follows.

1.1 Basic notation and constructions

Here is first a minimum of notation needed to state our results for the case of Coulombtype singularities, Theorem 1.1. This notation was explained in more detail in [6]. Let us fix from now on a *finite-dimensional*, Euclidean (real) vector space X. We shall let \mathbb{S}_X denote its *sphere at infinity* (more precisely: we define \mathbb{S}_X as the set of rays in X emanating form 0) and, then, we let $\overline{X} = X \sqcup \mathbb{S}_X$, called the *spherical compactification* of X, as described, for instance, in [6, Section 5.1] (see also [37, 38, 46, 47]). Note that here \sqcup denotes disjoint union as sets, but \overline{X} also carries a differential structure that makes \overline{X} diffeomorphic to a closed disk, see again [6, Section 5.1]. In particular, we have a diffeomorphism

$$\overline{X} \setminus \{0\} \simeq \mathbb{S}_X \times (0, \infty]. \tag{1}$$

Moreover, the Euclidean metric on *X* defines a natural diffeomorphism from \mathbb{S}_X to the unit sphere in *X*.

Let us assume throughout the paper that \mathcal{F} is a finite, non-empty set of linear subspaces of *X*. We need to define various distance functions in terms of \mathcal{F} . First, by $dist(x, y) = dist_{eucl}(x, y)$, we shall denote the Euclidean distance on *X* and, for any subset $Z \subset X$, we let

$$d_Z(x) := \operatorname{dist}(x, Z) := \inf_{z \in Z} |x - z|$$
 (2)

denote the Euclidean distance from *x* to *Z*. Let then $\bigcup \mathcal{F} := \bigcup_{Y \in \mathcal{F}} Y$ denote the union of the elements of \mathcal{F} and

$$\delta_{\mathcal{F}}(x) := \min_{Y \in \mathcal{F}} \{ d_Y(x), 1 \} = \min\{ \operatorname{dist}(x, \bigcup \mathcal{F}), 1 \}.$$
(3)

This function will be the weight used in the definition of the weighted Sobolev spaces that appear (sometimes only implicitly) in the statements of our results.

We shall assume throughout the paper, as in, for instance, [9, 13], that \mathcal{F} is stable under intersections. More precisely, let 2^A denote the set of all subsets of some set A. We shall say that $S \subset 2^A$ is a *semilattice*.¹ if $A, B \in S$ implies $A \cap B \in S$. Thus, from now on, we shall assume that \mathcal{F} is a *finite semilattice* of linear subspaces of Xsuch that $X \notin \mathcal{F}$ but $\{0\} \in \mathcal{F}$. (Mathematically the assumption that $X \notin \mathcal{F}$ is not a very important assumption, but it allows us to simplify the statement of Theorem 1.1, for instance.)

If Y is a linear subspace of X, its closure \overline{Y} in \overline{X} coincides with the spherical compactification of Y, so there is no danger of confusion. As in [6], to the semilattice \mathcal{F} we will associate the semilattices of *spherical compactifications* and, respectively, *spheres at infinity* of subspaces in \mathcal{F} :

$$\overline{\mathcal{F}} := \{ \overline{Y} \mid Y \in \mathcal{F} \} \text{ and } \mathbb{S}_{\mathcal{F}} := \{ \mathbb{S}_Y \mid Y \in \mathcal{F} \}.$$
(4)

¹ In the literature, what we call a semilattice is often called a *meet-semilattice*.

Finally, we can now introduce the main spaces considered in our main results as iterated blow-ups:

$$X_{\rm GV} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \quad \text{and} \quad X_{\mathcal{F}} := [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}]. \tag{5}$$

The notions of blow-up and iterated blow-up will be recalled in Sect. 3. We will see in Sect. 5.1.1 that $X_{\mathcal{F}} \simeq [X_{\text{GV}} : \widehat{\mathcal{F}}]$, where $\widehat{\mathcal{F}}$ denotes the lift of $\overline{\mathcal{F}}$ to X_{GV} , see Sect. 5.1 for details.

1.2 Statement of the main result and comments

The following result for eigenfunctions is formulated, for simplicity, just for the usual Coulomb-type singularities (see Theorems 6.1 and 6.6 for more general versions). There is no loss of generality to assume $X = \mathbb{R}^n$, and we shall do that when convenient, for instance, in the next theorem. Let $\partial_i := \frac{\partial}{\partial x^i}$ and $\partial^{\alpha} := \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_n^{\alpha_n}$, as usual. The Laplacian will always have the "analytic" sign convention, i.e., $\Delta = \sum_i \partial_i^2$.

Theorem 1.1 Let $\mathcal{F} \not\supseteq X$ be a finite semilattice of linear subspaces of $X := \mathbb{R}^n$, $\{0\} \in \mathcal{F}$. Let $X_{\mathcal{F}} := [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}]$ and let $d_Y(x) := \text{dist}(x, Y)$ and $\delta_{\mathcal{F}}(x) := \min\{\text{dist}(x, \bigcup \mathcal{F}), 1\}$ be the distance functions introduced in the previous subsection. For every $Y \in \mathcal{F} \cup \{X\}$, let $a_Y \in \mathcal{C}^\infty(X_{\mathcal{F}})$ and

$$V(x) := \sum_{Y \in \mathcal{F}} a_Y(x) d_Y(x)^{-1} + a_X(x).$$
(6)

Let us assume that $u \in L^2(X)$ satisfies $(\Delta + V)u = \lambda u$ in distribution sense on $X \setminus \bigcup \mathcal{F}$, for some $\lambda \in \mathbb{C}$. Then, for all multi-indices $\alpha \in \mathbb{N}^n$, we have

$$\delta_{\mathcal{F}}^{|\alpha|} \partial^{\alpha} u \in L^2(X).$$

Define $\rho(x):= \text{dist}_{\overline{g}}(x, \bigcup \mathcal{F})$, where \overline{g} is a (true) metric on X_{GV} (that is, a metric on the compact manifold with corners X_{GV} that is smooth up to the boundary). Then, we can use ρ instead of $\delta_{\mathcal{F}}$, see "Appendix A." That is, for *u* as in the theorem above and for all multi-indices $\alpha \in \mathbb{N}^n$, we have

$$\rho^{|\alpha|}\partial^{\alpha} u \in L^2(X).$$

Theorem 1.1 is, as we have already mentioned, a particular case of Theorem 6.6 (which, in turn, follows from Theorem 6.1). In particular, that theorem covers also the case of inverse square potentials, which have been studied recently in [14, 16, 19, 32, 33, 41] and in many other papers. We also consider higher-order uniformly strongly elliptic operators. This result strengthens a similar regularity result in [1]; compared to this previous result we will obtain better decay rates at infinity. Our results generalize immediately to systems. Regularity results for the eigenfunctions of Schrödinger operators have been obtained in many papers. See, for instance, [1, 11,

17, 18, 20–25, 28, 31, 34, 44, 45, 48, 49] and the references therein. Many methods in this article may be generalized to nonlinear equations, e.g., to semi-linear equations of the form

$$\Delta u + V|u|^s u = \lambda u$$

with V as in (6), $\lambda \in \mathbb{C}$ and $0 < s \le 2/(n-2)$, see Remark 6.7 for details.

1.3 Contents of the paper

The paper is essentially self-contained and consists of two parts. The first part contains mostly background material, but presented in a novel way. It consists of Sects. 2 and 3. In Sect. 2, we review manifolds with corners, their submanifolds, clean intersections of submanifolds, and some other basic concepts needed in the paper. We also recall in this section some results from [6] and from some earlier papers, including [1, 37, 1]42]. In Sect. 3, we recall and study the blow-ups and their iteration with respect to one or more suitable subsets. In particular, we extend the definition of an admissible order to a k-tuple of closed subsets of M and explain that the iterated blow-up with respect to a clean semilattice with an admissible order is defined and independent of the chosen admissible order [1, 6]. We also explain that the action of a Lie group on (M, S) extends to an action on the iterated blow-up [M : S]. The second part of the paper begins with Sect. 4, where we study metric aspects of the iterated blow-ups. We start by introducing the concept of a "smoothed distance function" to a p-submanifold $P \subset M$, which is a function that behaves like the distance to P close to P, but is smooth outside P. We study then the smoothed distance function to a blow-up suitable k-tuple, in general, and to a clean semilattice, in particular. Then, we recall how the metrics, the Sobolev spaces, and the differential operators change if one conformally changes the metric using a smoothed distance function. We apply these results to blowups (including the iterated ones). We distinguish here the case of a blow-up along a submanifold contained in the boundary and the case of a manifold not contained in the boundary. The main technical result of this section is the behavior of the smoothed distance functions when performing iterated blow-ups, Proposition 4.15. We also recall here the regularity result for the natural elliptic operators on Lie manifolds [3], which will then provide the regularity result of this paper. (The main work of this paper is to position ourselves to be able to use the regularity theorem for Lie manifolds.) In Sect. 5, we introduce the relevant semilattices and Lie manifolds needed to deal with the *N*-body problem and we particularize the constructions and results of Sect. 4 to this setting. Section 6 explains how to obtain our regularity result, Theorem 1.1 and some of its generalizations from the results of the other sections. Finally, the paper contains three appendices. In "Appendix A," we prove the equivalence of the functions ρ and $\delta_{\mathcal{F}}$ used in Theorem 1.1 and right after. (In fact, they are both equivalent to $\rho_{\mathcal{F}}$, the smoothed distance function to \mathcal{F} , which is the function that is actually used in the proofs, but is more difficult to define.) Appendices B and C contain technical details used in the article. The preprint version (arXiv 2012.13902) contains two additional appendices.

2 Preliminaries

We now recall a few concepts and results needed to understand the main results of this paper. For the concepts and results not recalled here, we refer to [1] or [6], on which this paper is heavily based.

2.1 A few basic notations and definitions

We need to complete the notation introduced in the Introduction as follows. We let

$$\mathbb{R}^n_k := [0,\infty)^k \times \mathbb{R}^{n-k} \text{ and } \mathbb{S}^{n-1}_k := \mathbb{S}^{n-1} \cap \mathbb{R}^n_k, \tag{7}$$

where \mathbb{S}^{n-1} is the unit sphere of \mathbb{R}^n for the Euclidean norm. The space \mathbb{S}_k^{n-1} will be called a *(generalized) orthant of the sphere*, for instance, \mathbb{S}_1^{n-1} is a half-sphere. It will be used in order to help us understand the smooth structure on the spherical compactification \overline{X} . More precisely, let us assume that $X = \mathbb{R}^n$ and let us consider the bijective map $\Theta_n : \overline{X} \to \mathbb{S}_1^n$

$$\begin{cases} \Theta_n(x) := \frac{1}{\sqrt{1 + \|x\|^2}} (1, x) \in \mathbb{S}_1^n & \text{if } x \in X, \\ \Theta_n(\mathbb{R}_+ v) := \frac{1}{\|v\|} (0, v) \in \mathbb{S}_1^n & \text{if } \mathbb{R}_+ v \in \mathbb{S}_X. \end{cases}$$
(8)

and its inverse $\Theta_n^{-1} : \mathbb{S}_1^n \to \overline{X}$

$$\Theta_n^{-1}(y_0, y_1, \dots, y_n) \longmapsto \begin{cases} \frac{1}{y_0}(y_1, \dots, y_n) \in \mathbb{R}^n & \text{if } y_0 \neq 0\\ \mathbb{R}_+(y_1, \dots, y_n) \in \mathbb{S}_X & \text{if } y_0 = 0 \end{cases}$$
(9)

We define the smooth structure on \overline{X} by the requirement that the map Θ_n be a diffeormphism as in [37, 42, 46]. For general X, we define the smooth structure on \overline{X} using a linear isomorphism $X \simeq \mathbb{R}^n$, for suitable *n*.

Remark 2.1 Let us assume that $Y = \mathbb{R}^p \subset \mathbb{R}^n = X$. Then, we have the following commutative diagram of smooth embeddings

$$\begin{array}{l} Y \hookrightarrow \overline{Y} \xrightarrow{\Theta_p} \mathbb{S}_1^p \\ \downarrow \qquad \downarrow \qquad \downarrow^n \\ X \hookrightarrow \overline{X} \xrightarrow{\Theta_n} \mathbb{S}_1^n \end{array}$$
(10)

which is a basic functoriality property of the spherical compactification. In particular, this shows that the closure of Y in \overline{X} can be identified as a differential manifold with \overline{Y} , the spherical compactification of Y, so there is no danger of confusion.

Let |A| denote the number of elements of a set A. We continue with several basic definitions.

Definition 2.2 Let $I \subset \{1, \ldots, n\}$ and

$$L_{I,k}^{n} := \{ x = (x_1, \dots, x_n) \in \mathbb{R}_k^{n} := [0, \infty)^k \times \mathbb{R}^{n-k} \mid x_i = 0 \text{ if } i \in I \}.$$
(11)

Then, $b_{I,k}^n := |I \cap \{1, \dots, k\}|$ will be called the *boundary depth* of $L_{I,k}^n$ in \mathbb{R}_k^n and $b_x := |\{i \in \{1, \dots, k\} \mid x_i = 0\}|$ will be called the *boundary depth* of $x = (x_1, \dots, x_n)$ in \mathbb{R}_k^n . Clearly, c := |I| is the *codimension* of L_I in \mathbb{R}_k^n and d := n - c is its *dimension*.

Obviously $b_{I,k}^n := \min\{b_x \mid x \in L_{I,k}^n\}$. The role of the sets $L_{I,k}^n$ is to serve as models for p-submanifolds [42, Definition 1.7.4]. A p-submanifold is a suitable submanifold of a manifold with corners, a concept recalled next. The letter "p" comes from the fact that for such a p-submanifold and any of its points, a local chart can be found around that point such that the submanifold is a factor of a *product*.

2.2 Manifolds with corners and their submanifolds

Let us now summarize a few basic definitions related to manifolds with corners [4, 35–38, 43]. Recall that a *manifold with corners* M of dimension n is a topological space locally modeled on $\mathbb{R}_k^n := [0, \infty)^k \times \mathbb{R}^{n-k}$ with smooth transition functions. The spaces \mathbb{R}_k^n and \mathbb{S}_k^n introduced in (7) provide simple examples of manifolds with corners. Given a point x in a manifold with corners M, the (*boundary*) depth of x in M is defined locally using Definition 2.2. More precisely, the boundary depth of x in M is the number of nonnegative coordinate functions vanishing at p in any local coordinate chart at p. It is the least k such that, for all $x \in M$, there exists a chart $\phi : U \to \mathbb{R}_k^n$ defined on an open neighborhood of x in M. For further reference, let us formalize the following concept in a definition.

Definition 2.3 The *boundary depth* of $P \subset M$ is the minimum over all boundary depths of points $x \in P$.

Let $(M)_k$ be the set of points of M of boundary depth k. Its connected components are called the *open* boundary faces of codimension (or boundary depth) k of M. A *boundary face* of boundary depth k is the closure of an open boundary face of boundary depth k. A boundary face H of codimension one of M will be called a *boundary hyperface* of M. The union of the boundary hyperfaces H of M is denoted ∂M ; it is the *boundary* of M. It is the set of points of M of boundary depth ≥ 1 . In particular, a subset $A \subset M$ has boundary depth 0 if, and only if, it is not contained in ∂M , the boundary of M.

The smooth functions on a manifold with corners M are defined as on their counterparts without corners. This is conveniently done by embedding M into a smooth manifold without corners \widetilde{M} , called an *enlargement* of M. This enlargement can also be used to define the tangent spaces of M.

Definition 2.4 Let *H* be a hyperface of *M* or a union of hyperfaces of *M*. A *boundary defining function* of *H* (in *M*) is a function $0 \le x_H \in C^{\infty}(M)$ such that $H = x_H^{-1}(0)$ and $dx_H \ne 0$ on *H*.



Fig. 1 A teardrop domain as a subset in \mathbb{R}^2

Remark 2.5 The example of the teardrop domain, see Fig. 1, shows that not all hypersurfaces have a boundary defining function. However, each (connected) boundary face F of codimension k can *locally* (i.e., , in a sufficiently small neighborhood U of any given point x) be represented as

 $F \cap U = \{x_1 = x_2 = \ldots = x_k = 0\},\$

where x_j are boundary defining functions in U of the hyperfaces of U containing $F \cap U$. Then, k is the boundary depth of F.

In order to define the blow-up, we need the concept of an *inward-pointing normal* bundle of a p-submanifold P of M. We first recall the concept of an inward-pointing tangent space to a manifold with corners M, a concept used also in [3, 4].

Definition 2.6 Let *M* be a manifold with corners. We let the *inward-pointing tangent* space T_x^+M to be defined as the set of derivatives

$$T_x^+ M := \{ \gamma'(0) \mid \gamma : [0, \epsilon) \to M \text{ smooth}, \quad \gamma(0) = x \}.$$

Its elements are called *inward-pointing tangent vectors* to $T_x M$.

We stress that, according to this definition, a tangent vector to M that is tangent to the boundary will automatically be an inward pointing tangent vector.

2.3 Submanifolds of manifolds with corners

The following concept introduced in [42] will play a crucial role in what follows. (Recall that |A| denotes the number of elements of a set *A*.)

Definition 2.7 A subset *P* of a manifold with corners *M* is a *p*-submanifold if, for every $x \in P$, there exists a chart $\phi : U \to \mathbb{R}^n_k$, with *U* an open neighborhood of *x* in *M*, and $I \subset \{1, 2, ..., n\}$ such that

$$\phi(P \cap U) = L^n_{I,k} \cap \phi(U),$$

with $L_{I,k}^n$ as defined in Eq. (11). The number n - |I| (respectively, |I|, respectively, $|I \cap \{1, \ldots, k\}|$) will be called the *dimension* (respectively, the *codimension* of *P*

at *x*, respectively, the *boundary depth* of *P* at *x*). We allow p-submanifolds *Y* of non-constant dimension. We define dim(*Y*) as the maximum of the dimensions of the connected components of *Y* and dim(\emptyset) = $-\infty$.

In particular, \mathbb{S}_k^n is a closed p-submanifold of $S_{k+k'}^{n+n'}$ of codimension n' and boundary depth k'.

Definition 2.8 Let $P \subset M$ be a p-submanifold of the manifold with corners M.

- (i) The quotient bundle $N^M P := TM|_P / TP$ is called the *normal bundle of P in M*.
- (ii) The image N^M₊ P of T⁺M|_P in N^MP is called the *inward-pointing normal fiber* bundle of P in M. One can show that, in a neighborhood of x ∈ P, it is a fiber bundle with fiber ℝ^{n'}_{k'} where n' is the codimension of P in M at x and where k' is the boundary depth of P in M.
- (iii) The set $\mathbb{S}(N_+^M P)$ of rays (emanating from 0) in $N_+^M P$ is called the *inward-pointing spherical normal bundle of* P *in* M. It is a fiber bundle with fibers $\mathbb{S}_{k'}^{n'-1}$, where n' and k' are as above.

Remark 2.9 A (positive definite) scalar product on TM induces a scalar product on $N^M P$. The choice of a metric on M will thus define a natural diffeomorphism from $\mathbb{S}(N^M_+ P)$ to the set of unit vectors in $N^M_+ P$.

Besides p-submanifolds, we shall also need *weak submanifolds*, which are really just plain submanifolds (without any other qualification or condition). We now recall (and slightly reformulate) the definition of a weak submanifold from [6]; the reformulation provides a definition equivalent to [6, Definition 2.10], but avoids introducing the concept of "submanifolds in Melrose's sense," [6, Definition B.1] and its simplified version [6, Definition 2.9]

Definition 2.10 A subset *S* of a manifold with corners *M* is a *weak submanifold of M* if, for every $p \in S$, there are

- natural numbers $k = k_p, m = m_p \in \{0, ..., n\}$ and $\ell = \ell_p \in \{0, ..., m\}$,
- a chart $\phi = \phi_p : U \xrightarrow{\sim} V \subset \mathbb{R}^n_k$ on M with V open in \mathbb{R}^n_k ,
- a diffeomorphism $\psi = \psi_p : \widetilde{V} \xrightarrow{\sim} W$ with \widetilde{V} and W open in \mathbb{R}^n

such that

(1) $p \in U$, (2) $V = \widetilde{V} \cap \mathbb{R}^n_k$, (3) $\psi(\phi(S \cap U)) = W \cap \mathbb{R}^m_\ell$.

If *S* is a weak submanifold of a manifold with corners *M*, then $\{\psi_p \circ \phi_p \mid p \in S\}$ provides an atlas for a manifold with corners structure on *S* such that the inclusion map $S \hookrightarrow M$ is an injective immersion and a homeomorphism onto its image. In conclusion, a weak submanifold of *M* is a manifold with corners on its own. Conversely, if *S* is an abstract manifold with corners, together with an injective immersion $\iota : S \to M$ that defines a homeomorphism from *S* to $\iota(S)$, then [6, Proposition 2.13] states that *S* is weak submanifold of *M*. Moreover, any p-submanifold of *M* is a weak submanifold of *M*.

The number m_p is the dimension of *S* at *p* and thus locally constant. (Again we do not require the function dim_{*p*}(*S*) to be constant on *S*.) The number ℓ_p is the (boundary) depth of *p* in *S*, and might by smaller, larger or equal to k_p , the boundary depth of *p* in *M*.

Note that Definition 2.10 is weaker than the notion of a submanifold in Melrose's sense [6, Definition B.1], which explain the word "weak" in the above definition.

2.4 Clean intersections

In order to study the iterate the blow-up construction in the next section, we will need *clean intersections*, which we recall next. For simplicity, we discuss only the case of p-submanifolds.

Definition 2.11 Let *P* and *Q* be two p-submanifolds of a manifold with corners *M* such that $P \cap Q$ is also a p-submanifold of *M*. We say that *P* and *Q* intersect cleanly or that they have a clean intersection if, for every $x \in P \cap Q$, we have $T_x(P \cap Q) = T_x P \cap T_x Q$.

For example, if P is a p-submanifold of M and F is a boundary face of M, then F and P have a clean intersection. This is not, however, the case, in general, if P is just a *weak submanifold* of M. This explains, in particular, the need to consider p-submanifolds. We next recall how to extend the definition of clean intersection to semilattices of p-submanifolds.

Definition 2.12 Let S be a finite semilattice of p-submanifolds of M. We call S a *clean semilattice of p-submanifolds (of M)* if any $P, Q \in S$ intersect cleanly.

Remark 2.13 If S is a clean semilattice of p-submanifolds then, for all $P_1, \ldots, P_k \in S$ and all $x \in \bigcap_{j=1}^k P_j$, we obtain that

$$T_x\left(\bigcap_{j=1}^k P_j\right) = \bigcap_{j=1}^k T_x P_j.$$
 (12)

In particular, our definition of clean semilattices coincides with Definition 2.7 in [1]. This property does not hold anymore, if S is not a semilattice, as seen in the next example.

Example 2.14 Assume that S is a set of p-submanifolds with k elements, such that each pair $\{P_i, P_j\} \subset S$ intersections cleanly, then we cannot conclude, in general, that we have the property expressed in Eq. (12) for $k \ge 3$. Indeed, let us consider the family consisting of the following three surfaces P_1 , P_2 and P_3 in \mathbb{R}^3 (i.e., two-dimensional submanifolds of \mathbb{R}^3):

$$P_{1} := \left\{ (x, y, z)^{T} \in \mathbb{R}^{3} \mid y = 0 \right\}, \qquad P_{2} := \left\{ (x, y, z)^{T} \in \mathbb{R}^{3} \mid z = 0 \right\},$$
$$P_{3} := \left\{ (x, y, z)^{T} \in \mathbb{R}^{3} \mid y + z = x^{2} \right\}.$$

🖄 Springer

Each pair intersects cleanly in a one-dimensional submanifold. However, $P_1 \cap P_2 \cap P_3 = \{0\}$, but $T_0P_1 \cap T_0P_2 \cap T_0P_3 = \mathbb{R}(1, 0, 0)^T$. Thus, the triple (P_1, P_2, P_3) does not intersect cleanly.

The following Lemma will provide the needed examples of pairs of cleanly intersecting p-submanifolds.

Lemma 2.15 Let Y, Z be two linear subspaces of X. Then, all subsets of $\{\overline{Y}, \overline{Z}, \mathbb{S}_Y, \mathbb{S}_Z\}$ intersect cleanly.

The proof is a direct verification. We include it as an appendix in the arxiv version of this article.

3 Blow-ups

In this section, we gather the needed results on blow-ups, following the approach of [6]. Most of the results in this section were discussed in a similar form in the literature, see [1, 6, 28, 29, 37, 42] for more details. In particular, the basic construction of the blowup along a p-submanifold coincides with the one in these references. However, the lifting (or pullback) of submanifolds differs in certain cases from the one used in, for instance, [30, 37, 42]. As a consequence, our notion of iterated blow-up exhibits some subtle differences to the iterated blow-up discussed in the aforementioned papers. Our slightly different approach to the iterated blow-up has better locality and functorial properties, in particular it is compatible with restriction to open subsets, and thus avoids discussing additional special cases. However, our modified approach requires additional attention when citing the existing literature: for example [30, Lemma 7.2 (b)] does not hold using our definitions. As a consequence, the current section will be written in a self-contained way, both in order to provide reliable foundations and for better readability. The subtle difference in the approaches, however, will finally have no effect in our applications: in these applications, our iterated blow-up spaces coincide with the iterated blow-up spaces in the work of Melrose and in the related work cited above.

3.1 The blow-up along a closed p-submanifold

We now recall the definition of the blow-up [M : P] of a manifold with corners M along a closed p-submanifold P of M. See [6, 37] for further details and references. We begin by specifying the underlying set of the blow-up [M : P]. Its topology and smooth structure will be defined shortly after that. If A and B are disjoint, we sometimes denote $A \sqcup B := A \cup B$ their union.

Definition 3.1 (see [6, Definition 3.1]) Let *M* be a manifold with corners and *P* be a *closed* p-submanifold of *M*. We let $\mathbb{S}(N^M_+ P)$ denote the inward pointing spherical normal bundle of *P* in *M* (Definition 2.8). Then, as a set, the *blow-up of M along P* is the disjoint union

$$[M:P] := (M \smallsetminus P) \sqcup \mathbb{S}(N^M_+ P).$$

The blow-down map $\beta = \beta_{M,P} : [M : P] \to M$ is defined as the identity map on $M \setminus P$ and as the fiber bundle projection $\mathbb{S}(N^M_+ P) \to P$ on the complement.

If $P \subset \partial M$ in the above definition, we shall say that [M : P] is a *boundary blow-up*. If no component of *P* is contained in the boundary of *M*, then we shall say that [M : P] is an *interior blow-up*. See [6, 37, 42], for instance, for the definition of the topology and smooth structure on the blow-up [M : P]. Let us say, nevertheless, that the smooth structure on [M : P] is defined using Euclidean model spaces and is such that it induces the given smooth structures on $M \setminus P$ and on $\mathbb{S}(N^M_+ P)$. We also remark that the topology on [M : P] is such that *M* has the quotient topology with respect to $\beta_{M,P}$. The following extreme cases of blow-ups deserve a special treatment.

Remark 3.2 If *P* is a union of connected components of *M*, then $[M : P] = M \setminus P$. In particular, we have $[M : \emptyset] = M$ and $[M : M] = \emptyset$.

We shall need the following Proposition from [6]. In that proposition a *lift* denotes a smooth map j^{β} that yields a commutative diagram



Proposition 3.3 [6, Proposition 3.14] Let P and Q be closed p-submanifolds of M intersecting cleanly. Then, the inclusion $j : Q \to M$ lifts uniquely to a smooth map

$$\underbrace{(Q \smallsetminus (P \cap Q)) \sqcup \mathbb{S}(N^Q_+(P \cap Q))}_{[Q:P \cap Q] :=} \xrightarrow{j^\beta} \underbrace{(M \smallsetminus P) \sqcup \mathbb{S}(N^M_+P)}_{[M:P] :=}.$$

This map is an injective immersion, a homeomorphism onto its image, and the image of j^{β} is a p-submanifold. Moreover

$$\overline{\beta_{M,P}^{-1}(Q \smallsetminus P)} = j^{\beta}([Q : P \cap Q]).$$

In view of the above proposition, we shall write $[Q : P \cap Q] \subset [M : P]$, by abuse of notation.

We have the following useful factorization lemma due (essentially) to Kottke [37]. It deals with a particular, but important case of the iterated blow-up. We shall need to recall the formulation of the first factorization Lemma from [6], i.e., [6, Lemma 4.7], which deals with the setting $Q \subset P \subset M$ (see also [1]).

Lemma 3.4 Let us assume that Q is a p-submanifold of P and that P is a p-submanifold of M. Then, Q is a p-submanifold of M. Moreover, there exists a smooth, canonical map

$$\zeta_{M,Q,P} : [M:Q,P] := \left[[M:Q] : [P:Q] \right] \rightarrow [M:P]$$

that restricts to the identity on $M \setminus P$.

3.2 Iterated blow-ups

In the applications of the blow-up, we typically have to blow up several subsets. This subsection deals with some of the intricacies of this procedure. The first thing to discuss is the *pullback* of a subset in M to the blow-up [M : P] following [6, 37, 38].

Definition 3.5 Let *P* be a p-submanifold of *M* and *Q* be a closed subset of *M*. The *pullback* or *lifting* $\beta^*_{M-P}(Q)$ of *Q* to [M:P] is defined by

$$\beta_{M,P}^*(Q) := \overline{\beta^{-1}(Q \setminus P)} \tag{13}$$

In the case $Q \subset P$, our definition of $\beta_{M,P}^*(Q)$ is different from the one given by Melrose in [42, Chapter 5, Section 7], see [6, Remark 3.16] for details.

The following factorization lemma is similar in spirit, only easier (see [1, 6, 37]).

Lemma 3.6 Let us assume that P and Q are closed, disjoint p-submanifolds of M. We have $\beta_{M,Q}^*(P) = P$ and similarly for P and Q switched. With these identifications, we then have $[[M : Q] : P] \simeq [[M : P] : Q]$ canonically, and hence there exists a smooth, canonical map

$$\zeta_{M,Q,P}: \left[[M:Q]:P \right] \to [M:P]$$

that restricts to the identity on $M \setminus (P \cup Q)$.

In order to introduce the iterated blow-up with respect to an ordered family (*n*-tuple) of closed subsets, we first introduce the families with respect to which we can define the blow-up. We found it convenient to allow repetitions in these families. The following is Definition 3.8 in [6] and Definition 2.9 in [1], but see also [37] and [42].

Definition 3.7 Let *M* be a manifold with corners and let $\mathcal{P}:=(P_i)_{i=1}^k$ be a *k*-tuple of closed subsets of *M*, $k \ge 1$. If P_1 is a *closed p-submanifold* of *M* and k > 1, we define the *pullback* of \mathcal{P} to be the (k - 1)-tuple

$$\mathcal{P}' := \beta_{M,P_1}^* (\mathcal{P} \setminus \{P_1\}) := \left(\beta_{M,P_1}^* (P_i)\right)_{i=2}^k.$$

Then, by induction on k, we say that \mathcal{P} is *blow-up-suitable (in M)* if:

- (i) P_1 is a closed *p*-submanifold of *M* and
- (ii) if k > 1, the pullback \mathcal{P}' is blow-up-suitable in $[M : P_1]$.

Of course, $k = |\mathcal{P}|$. If k = 0, that is, if $\mathcal{P} = \emptyset$, then we also say that \mathcal{P} is blow-up-suitable in M.

In what follows, we will often use the pullback \mathcal{P}' of various *k*-tuples \mathcal{P} . We now define the iterated blow-up with respect to blow-up-suitable families.

Definition 3.8 We use the notation introduced in Definition 3.7, in particular, $\mathcal{P}:=(P_i)_{i=1}^k$ is a *k*-tuple of closed subsets of *M* and $\mathcal{P}':=\beta_{M,P_1}^*(\mathcal{P}\setminus\{P_1\}):=(\beta_{M,P_1}^*(P_i))_{i=2}^k$ (the pullback of \mathcal{P}). If \mathcal{P} is blow-up-suitable, then we define by induction the *iterated* blow-up $[M:\mathcal{P}]$ by

$$[M:\mathcal{P}] = [M:(P_i)_{i=1}^k] := \begin{cases} [M:P_1] & \text{if } k = 1, \\ [[M:P_1]:\mathcal{P}'] & \text{if } k > 1. \end{cases}$$

The blow-down map $\beta_{M,\mathcal{P}} : [M : \mathcal{P}] \to M$ is defined by induction on k as the composition of the blow-down maps $[M : \mathcal{P}] := [[M : P_1] : \mathcal{P}'] \to [M : P_1] \to M$.

It is not difficult to see, by induction, that Definition 3.8 makes sense. This is the point of introducing Definition 3.7. We shall also use $[M : P_1, P_2, ..., P_k]$ as an alternative notation for $[M : \mathcal{P}]$. Let $\mathcal{E} := \emptyset$, (which is regarded as a *k*-tuple with k = 0) and define $[M : \mathcal{E}] = M$ and $\mathcal{P}' = \mathcal{E}$ for if \mathcal{P} has only one element. Then, the relation $[M : \mathcal{P}] = [[M : P_1] : \mathcal{P}']$ remains true also for k = 1. This may be useful for proofs by induction, for instance in the proof of Lemma 4.10.

Remark 3.9 Of course, a *k*-tuple \mathcal{P} without repetitions is the same thing as a linearly ordered set. We want the \mathcal{P} in the definition of the iterated blow-up $[M : \mathcal{P}]$ to be a *k*-tuple rather than a linearly ordered finite set of p-submanifolds. That is, we want to allow repetitions in \mathcal{P} . The reason for this choice is that, even if \mathcal{P} does not have repetitions, its pullback $\mathcal{P}' := (\beta^*(P_i))_{i=2}^k$ might have repetitions. An example is provided by $\mathcal{P} = (A, B, A \cup B)$, where A and B are disjoint closed p-submanifolds of M, in which case $\mathcal{P}' = (B, B)$. We shall often consider semilattices with an additional total order (other than the order given by inclusion).

In the next remark we will explain how to eliminate repetitions. For this purpose, the following proposition is a useful technical result. Recall that $\bigcup \mathcal{P} := \bigcup_{P \in \mathcal{P}} P$.

Proposition 3.10 We use the notations of Definition 3.8 and we assume that \mathcal{P} is blow-up-suitable (and hence that $[M : \mathcal{P}]$ is defined). Then,

(i) $M \setminus \bigcup \mathcal{P} \subset [M : \mathcal{P}]$ and $\beta = id$ on $M \setminus \bigcup \mathcal{P}$ and (ii) $M \setminus \bigcup \mathcal{P}$ is open and dense in $[M : \mathcal{P}]$.

Proof The first part follows from the definitions of the blow-up and of the iterated blow-up. To prove the second part, we proceed by induction on k, the number of elements of \mathcal{P} . The case k = 1 follows from the definition (and was discussed also in [6]). For the induction step, we notice that

$$M \smallsetminus \bigcup \mathcal{P} = (M \smallsetminus P_1) \smallsetminus \bigcup \mathcal{P}' \subset [M:P_1] \smallsetminus \bigcup \mathcal{P}'$$

is dense in $[M : P_1] \setminus \bigcup \mathcal{P}'$ by the case k = 1, since $\bigcup \mathcal{P}'$ is closed. Since $[M : P_1] \setminus \bigcup \mathcal{P}'$ is dense in $[M : \mathcal{P}] := [[M : P_1] : \mathcal{P}']$ by the case k - 1 (the induction hypothesis), the result follows.

Given two blow-ups of M, we shall say that they are *canonically diffeomorphic* if there exists a diffeomorphism that is the identity outside the sets that are blown up. In view of the above proposition, this entails a uniqueness property for the canonical diffeomorphisms. We now explain how we can eliminate the repetitions in \mathcal{P} .

Remark 3.11 We use the notation introduced in Definition 3.8 and the results of Proposition 3.10. Let \mathcal{P}_{red} be obtained from \mathcal{P} by removing repetitions by keeping only the *first* appearance of a p-submanifold. Thus, if $\mathcal{P} = (\emptyset, A, B, A)$, then $\mathcal{P}_{red} = (\emptyset, A, B)$. Then, $[M : \mathcal{P}]$ is defined if, and only if, $[M : \mathcal{P}_{red}]$ is defined. Moreover, these two iterated blow-ups are canonically diffeomorphic (when defined), in the sense that:

- (i) $M \setminus \bigcup \mathcal{P} = M \setminus \bigcup \mathcal{P}_{red};$
- (ii) $M \setminus \bigcup \mathcal{P}$ is dense in $[M : \mathcal{P}]$ and $M \setminus \bigcup \mathcal{P}_{red}$ is dense in $[M : \mathcal{P}_{red}]$, by Proposition 3.10; and
- (iii) the identity map $M \setminus \bigcup \mathcal{P} \to M \setminus \bigcup \mathcal{P}_{red}$ extends to a diffeomorphism $[M : \mathcal{P}] \to [M : \mathcal{P}_{red}]$, which is unique by the density properties of ii.

In addition to removing repetitions, we could as well remove entries $P_j = \emptyset$ from \mathcal{P} with the same effect. However, we found it convenient for exposition purposes to do exactly the opposite, that is, to usually assume \emptyset to be the first entry of \mathcal{P} , especially if \mathcal{P} is a semilattice. We have the following properties.

Remark 3.12 We use the notation introduced in Definition 3.8. In what follows, the pullback operation $\mathcal{P} \mapsto \mathcal{P}'$ introduced in

- (i) We have [M : P]:=[[M : P₁], P'] = [[M : P₁] : P'_{red}], which may be useful if one wants to deal only with *reduced* tuples (which is the same as linearly ordered finite sets).
- (ii) Let us introduce the *iterated pullbacks* of \mathcal{P} by $\mathcal{P}^{(1)}:=\mathcal{P}'$ and $\mathcal{P}^{(k+1)}:=(\mathcal{P}^{(k)})'$. Then,

$$[M:\mathcal{P}] = [[M:P_1, P_2, \dots, P_k]:\mathcal{P}^{(k+1)}].$$

- (iii) $(\mathcal{P}'_{red})'_{red} = \mathcal{P}''_{red}$ (where ' always comes before _{red}, meaning that $\mathcal{P}'_{red} := (\mathcal{P}')_{red}$).
- (iv) A special situation arises if P_1 , the first element of \mathcal{P} , is $P_1 = \emptyset$, which is, in fact, the norm when dealing with semilattices. Then, in our definition of the blow-up, the first step, the blow-up with respect to P_1 is trivial (it does not change our sets, except that it removes P_1 from the list). In particular, $\mathcal{P} = (\emptyset, \mathcal{P}')$. It is the next blow-up that may be interesting. For us, it will then be useful to introduce the following notation:

$$\widetilde{\mathcal{P}} := (\emptyset, \mathcal{P}'') := (\emptyset, \beta_{M;P_2}^*(P_3), \beta_{M;P_2}^*(P_4), \dots, \beta_{M;P_2}^*(P_k)).$$

(Note that $\beta^*_{M:P_2}(P_2) = \emptyset$.) We then have the relation

$$[M:\mathcal{P}] = [M:\mathcal{P}'] = [[M:P_2]:\mathcal{P}''] = [[M:P_2]:\tilde{\mathcal{P}}]$$

= [[M:P_2]: $\tilde{\mathcal{P}}_{red}$].

We also notice that, if \mathcal{P}_{red} is a clean semilattice, then $\widetilde{\mathcal{P}}_{red} = (\widetilde{\mathcal{P}_{red}})_{red}$ is also a clean semilattice; here the semilattice property is obvious and the cleanness was proved in [1, Theorem 2.8]. (In this context we define again $\widetilde{\mathcal{P}}_{red} := (\widetilde{\mathcal{P}})_{red}$, that is, the "tilde" comes before "reduced.")

(v) To wrap up the list of needed properties, let us notice that

$$(\emptyset, \mathcal{P})' = \mathcal{P}$$
 and $[M : (\emptyset, \mathcal{P})] = [M : \mathcal{P}].$

In particular, if we define $\mathcal{P}_0 := \mathcal{P}^{(0)} := \mathcal{P}, \mathcal{P}_{k+1} := \widetilde{\mathcal{P}}_k$ and $\mathcal{P}^{(k+1)} = (\mathcal{P}^{(k)})'$, then $\mathcal{P}_k = (\emptyset, \mathcal{P}^{(k+1)})$ for $k \ge 0$. Hence, the iteration of the tilde operation can be expressed in terms of the iteration with respect to the prime operation and the addition of the empty set.

The following results generalize to the iterated blow-up [6, Lemma 3.9] on the compatibility of the blow-up with products.

Proposition 3.13 Let M and M_1 be two manifolds with corners and $\mathcal{P} = (P_1, P_2, ..., P_k)$ be a blow-up-suitable (in M) k-tuple of closed subsets of M. Then, $\mathcal{P} \times M_1 := (P_1 \times M_1, P_2 \times M_1, ..., P_k \times M_1)$ is a blow-up-suitable (in $M \times M_1$) k-tuple of closed subsets of $M \times M_1$ and there exists a canonical diffeomorphism $[M \times M_1 : \mathcal{P} \times M_1] \simeq [M : \mathcal{P}] \times M_1$ such that the following diagram commutes:

Proof If k = 1 (that is, \mathcal{P} consists of a single set), then the result was proved in [1] (it can be found also in [6]). In general, it follows by induction, using again the result from [1] (the case k = 1) and using also that $\beta^*(P) \times M_1 = \beta^*(P \times M_1)$, where β is an appropriate blow-down map.

3.3 Admissible orders

A natural question when we do iterated blow-ups of a manifold with corners M along an ordered family $\mathcal{P} = (P_0, P_1, \dots, P_k)$, is to decide how the order of the blow-up influences the final space [37]. In particular, a related question is whether the iterated blow-up is defined for a given order. The aim of this subsection is to recall the results of [1, 6, 37] that give a positive answer to these questions if "admissible orders" are used. Before stating the main result from [6], we first introduce admissible orders and graph blow-ups, which will be needed for the statement of the theorem.

If, in the definition of a blow-up-suitable k-tuple \mathcal{P} , we further require P_1 to be minimal for inclusion, we obtain the notion of an "admissible" k-tuple. Let us state this explicitly.

Definition 3.14 Let *M* be a manifold with corners and let $\mathcal{P}:=(P_i)_{i=1}^k$ be a *k*-tuple of closed subsets of $M, k \ge 1$. By induction on *k*, we say that \mathcal{P} is *admissible (in M)* if:

- (i) P_1 is a closed *p*-submanifold of M,
- (ii) there is no i > 1 such that $P_i \subsetneq P_1$, and
- (iii) if k > 1, \mathcal{P}' is admissible in $[M : P_1]$, where $\mathcal{P}' = (\beta_{M,P_1}^*(P_j))_{j=2}^k$ is the pullback of \mathcal{P} as before (in particular, β_{M,P_1}^* is as defined in (13)).

Of course, $k \ge |\mathcal{P}|$. If k = 0, that is, if $\mathcal{P} = \emptyset$, then we also say that \mathcal{P} is admissible in M.

This definition of and admissible k-tuple is more general than the one in [6].

Remark 3.15 We notice the following

- (1) A k-tuple with an admissible order is, in particular, also blow-up-suitable.
- (2) \mathcal{P} is admissible if, and only if, (\emptyset, \mathcal{P}) is admissible.
- (3) Assume the (k + 1)-tuple S = (P₀, P₁,..., P_k) is a clean semilattice, P₀ = Ø. Then, S̃:=(Ø, β^{*}_{M,P1}(P₂), β^{*}_{M,P1}(P₃), ..., β^{*}_{M,P1}(P_k)) is also a clean semilattice by the results of [1], see Remark 3.12 (iv). Moreover, S (with the indicated order) is admissible if, and only if, S̃ (with the induced order) is admissible. This explains why we sometimes consider S̃ and do not work exclusively with the iterated pullbacks S', S", ..., S^(j).

We now consider a different type of blow-up with respect to a *k*-tuple of p-submanifolds that is immediately seen not to depend on the choice of the order. To define it, we introduce the *multi-diagonal map*. For a semilattice S endowed with an admissible order, we let $U:=M \setminus \bigcup S$, which is a dense subset of [M : S]. The multi-diagonal map is the map

$$\delta: U \to \prod_{P \in \mathcal{S}} [M:P], \quad x \mapsto (x, x, \dots, x).$$
(15)

Definition 3.16 Let $\mathcal{P} = (P_i)$ be a *k*-tuple of closed p-submanifolds of the manifold with corners *M* and let δ be the multi-diagonal map defined in (15). Then, the *graph blow-up* $\{M : \mathcal{P}\}$ of *M* along \mathcal{P} is defined by

$$\{M:\mathcal{P}\} := \overline{\delta(M \setminus \bigcup \mathcal{P})} = \overline{\{(x, x, \dots, x) \mid x \in M \setminus \bigcup \mathcal{P}\}} \subset \prod_{i \in I} [M:P_i].$$

For the graph blow-up, we can also remove the repetitions.

Remark 3.17 The graph blow-up $\{M : \mathcal{P}\}$ is a weak submanifold of $Q := \prod_{i \in I} [M : P_i]$. The notion of a "weak submanifold" was introduced in [6, Subsec. 2.3.1]; however, the only aspect we need to know about it here is that, if one restricts the sheaf of smooth functions on Q to $\{M : \mathcal{P}\}$, then this restriction defines the structure of a manifold with corners on $\{M : \mathcal{P}\}$ which is compatible with the topology induced from Q.

Theorem 4.19 of [6] shows that, if S is a semilattice endowed with an admissible order, then the iterated blow-up and the graph blow-ups of M with respect to S are canonically diffeomorphic. The following statement combines the statement of that theorem with part of Remark 4.13 and with Theorem 4.21 of that paper.

Theorem 3.18 Let $S \ni \emptyset$ be a clean semilattice of closed p-submanifolds of M with an admissible order on its elements (Definition 3.14). Then, the iterated blow-up [M : S] is defined and $\{M : S\}$ is a manifold with corners that is canonically diffeomorphic to [M : S]. If G is a Lie group acting smoothly on M such that G maps S to itself, then G acts by diffeomorphisms on [M : S].

In the theorem "canonically diffeomorphic" means that the multi-diagonal map defined in (15) has a unique smooth extension which provides this diffeomorphism.

The iterated blow-up does not depend on the order of the sets (up to a canonical diffeomorphism). Hence, an immediate consequence of the last theorem is that the iterated blow-up [M : S] does not depend on the choice of the *admissible* order on S.

The reader may wonder, at this time, whether admissible orders exist on a given k-tuple. Note that, at this time, it is not clear even that a blow-up-suitable order exists on a given k-tuple of closed p-submanifolds of M. The following proposition gives a positive answer to this question if our k-tuple is a *clean semilattice*. This result and the previous theorem motivates the use of semilattices in our work.

Remark 3.19 Let S be a finite, clean semilattice of closed p-submanifolds of M, $\emptyset \in S$. Let us assume construct an order (P_0, P_1, \ldots, P_k) on the elements of $S = \{P_0, P_1, \ldots, P_k\}$ by choosing $P_j \in S$ by induction on $0 \le j \le k = |S| - 1$ as follows.

- (i) Let us choose an arbitrary initial order on the elements of S, to be able to talk about pullbacks. With each choice of P_j , we modify this order by moving P_j on the (j + 1)-position. (So, after choosing P_j , the first (j + 1)-elements of the modified order will be the chosen elements (P_0, P_1, \ldots, P_j) .)
- (ii) We begin by choosing $P_0:=\emptyset$, which implies that the pullback S' is given by $S' = S \setminus \{\emptyset\}$ (after which we change the order on S according to the previous point).
- (iii) We let P_1 to be an arbitrary minimal element of S' for inclusion.

(iv) Let

$$\widetilde{\mathcal{S}}:=(\emptyset, \mathcal{S}''):=(\emptyset, \beta_{M, P_1}^*(\mathcal{S} \smallsetminus \{\emptyset, P_1\})) = \beta_{M, P_1}^*(\mathcal{S} \smallsetminus \{\emptyset\}).$$

Then, the elements of \widetilde{S} form again a clean semilattice of closed p-submanifolds of M [1, Theorem 2.8] and we choose $P_2 \in S \setminus \{P_0, P_1\}$ such that the lift $\beta_{M,P_1}^*(P_2)$ of P_2 is minimal element of S'' for inclusion. In particular, $\beta_{M,P_1}^*(P_2)$ will be a closed p-submanifold of $[M : P_1]$ by the aforementioned result in [1].

(v) We then iterate this construction with \widetilde{S} in place of S. More precisely, recall from Remark 3.12(v) that $S_j := \widetilde{S}_{j-1} = (\emptyset, S^{(j+1)})$, where $S^{(j+1)} = (S^{(j)})'$ and $S_0 = S^{(0)} = S$. Assume P_0, P_1, \ldots, P_j were chosen. Then, we choose $P_{j+1} \in S \setminus \{P_0, P_1, \ldots, P_j\}, j+1 \le k$, to correspond to a minimal element

of $S^{(j+1)}$ for inclusion. This satisfies the desired condition that the lift of P_{j+1} in $S^{(j+1)}$ be a p-submanifold.

Proposition 3.20 We use the notation introduced in Remark 3.19. First, the procedure of that remark is well defined in the sense that it yields an order (P_0, P_1, \ldots, P_k) on the elements of S. Most importantly, the resulting order is admissible. Let $\emptyset \neq Y \in S$. With suitable choices in the procedure of that remark, we obtain an admissible order on S such that all elements that precede Y in this order are contained in Y.

Proof The procedure of Remark 3.19 is well defined since, at each step, the resulting tuples $\widetilde{S} := (\emptyset, S'')$ and $(\emptyset, S^{(j+1)}) := (\widetilde{\emptyset}, S^{(j)}), k \ge 1$, are clean semilattices, by [1], so the procedure stops only after we have chosen P_k (i.e., after having ordered all elements). (See Remark 3.12(v) for the notation.) The resulting order on S is admissible, by the definition of an admissible order and by induction on the number of elements of S. Finally, given $Y \in S$, we are going to successively choose the sets P_k to be either Y or a set contained in Y, if possible (that is, unless Y has already been chosen). This procedure will yield the desired admissible order on S.

3.4 A blow-up point of view on the spherical compactification

This subsection is not needed for the main results of the article. We will show the following proposition which might add a helpful perspective for future research. In particular, it shows that if we blow up at infinity the one-point compactification X_{∞} of a finite-dimensional vector space X obtained by stereographic projection, then we obtain a compactification canonically diffeomorphic to the spherical compactification \overline{X} .

For introducing the one-point compactification X_{∞} , we use the scalar product $\langle \cdot, \cdot \rangle$ on *X*, and we consider the (unit) sphere $\mathbb{S}_{\mathbb{R}\times X}$ in $\mathbb{R} \times X$. We define the *south pole* $S:=(-1, 0) \in \mathbb{S}_{\mathbb{R}\times X}$. The *stereographic projection* is the map

$$\sigma : \mathbb{S}_{\mathbb{R} \times X} \smallsetminus \{S\} \to X,$$
$$\mathbb{R} \times X \ni \begin{pmatrix} \cos \theta \\ \sin(\theta) \cdot y \end{pmatrix} \mapsto \tan\left(\frac{\theta}{2}\right) \cdot y, \quad \theta \in (0, \pi], \quad y \in \mathbb{S}_X.$$

After formally defining $\infty := \sigma(S)$, there is a unique topology and a unique smooth structure on $X_{\infty} := X \cup \{\infty\}$ such that σ is a diffeomorphism. This manifold X_{∞} is called the *one-point-compactification* of X.

We also consider a new map Θ_X , which is a slightly modified version of the map Θ_n from (8). For that purpose, let

$$\begin{split} \mathbb{S}'_{+} &:= \left\{ x + S \mid x \in \mathbb{S}_{[0,\infty) \times X} \right\} = \left\{ x - (1,0) \mid x = (x^{0}, x'), \ \|x\| = 1, \ x^{0} \ge 0 \right\}, \\ \Theta_{X} &: \overline{X} \to \mathbb{S}'_{+}, \quad \begin{cases} \Theta_{X}(x) &:= \frac{1}{\sqrt{1 + \langle x, x \rangle}} (0, x) \in \mathbb{S}'_{+} & \text{if } x \in X, \\ \Theta_{X}(\mathbb{R}_{+}v) &:= \frac{1}{\|v\|} (-1, v) \in \mathbb{S}'_{+} & \text{if } \mathbb{R}_{+}v \in \mathbb{S}_{X}. \end{cases} \end{split}$$

We write $S_X := \overline{X} \setminus X$ for the sphere at infinity of *X*, as usual.



Fig. 2 One-point and disk compactification. The stereographic projection σ maps the dot on the circle to the dot on the line, and the map Θ_X maps the dot on the line to the dot on the half-circle

Lemma 3.21 Let $\Psi: \overline{X} \to X_{\infty}$ be the map with $\Psi|_X = id_X$ and $\Psi(\mathbb{S}_X) = \{\infty\}$. This is a smooth map from the spherical compactification \overline{X} to the one-pointcompactification X_{∞} , extending the identity id_X . There is a diffeomorphism $\Phi: \overline{X} \to [X_{\infty}: \{\infty\}]$ extending id_X , and thus $\Psi \circ \Phi^{-1}$ is the blow-down map $\beta_{X_{\infty}, \{\infty\}}$.

Proof We define $\psi := \sigma^{-1} \circ \Theta_X^{-1} : \mathbb{S}'_+ \to \mathbb{S}_{\mathbb{R} \times X}$. Then, $\Psi = \sigma \circ \psi \circ \Theta_X$. By construction, σ and Θ_X are diffeomorphisms. To prove the lemma, one thus has to show that ψ is that there is a diffeomorphism $\phi : \mathbb{S}'_+ \to [\mathbb{S}_{\mathbb{R} \times X} : \{S\}]$ such that $\psi \circ \phi^{-1} = \beta_{\mathbb{S}_{\mathbb{R} \times X}, \{S\}}$. The construction of such a ϕ is an easy exercise, if one uses the following formula for ψ , which is apparent in view of Fig. 2, setting $\theta := 2\alpha$ above.

$$\psi\Big(\big((\cos\alpha)-1,\,(\sin\alpha)y\big)\Big) = \big((\cos2\alpha),\,(\sin2\alpha)y\big), \qquad \alpha \in [0,\,\pi/2], \ Y \in \mathbb{S}_X.$$

The function $\cos \alpha$ is a boundary defining function for \mathbb{S}'_+ and a smoothed distance function for *S* in $\mathbb{S}_{\mathbb{R}\times X}$. This completes the proof.

4 Distance functions and Sobolev spaces for blown-up spaces

We now investigate how several geometric quantities (metrics, distance functions, Sobolev spaces, natural differential operators, ...) change when performing a blow-up. The manifolds M we consider have a complete metric in the interior M_0 described precisely in terms of Lie manifolds, Definition 4.20. One important case will be that when M_0 is a Euclidean vector space X with its spherical compactification $M = \overline{X}$, which was described in the Introduction. Some other times, M will be a blow-up of \overline{X} .

We distinguish here the case of a blow-up along a submanifold *contained in the boundary* (the easy case) and the case of a manifold *not contained in the boundary* (the difficult case, but treated already in [1] and in other papers). In the first case, the *boundary case*, we will additionally require that our vector fields are tangent to the submanifold contained in the boundary. In that case, the metric in the interior remains the same, only the compactification is altered, which makes many investigations much easier. In the second case, the *interior case*, the metric will be changed conformally by multiplication with r^{-2} , where *r* is a "smoothed distance function," (a concept that will be defined in this section). The main technical result of this section is the behavior of "smoothed distance functions" when performing iterated blow-ups, Proposition 4.15. We also recall in this section the needed regularity result on Lie manifolds from [3].

4.1 Smoothed distance function to a p-submanifold

Let *M* denote a manifold with corners, as before. We assume that $\mathcal{P} \subset 2^M$ —equipped with a suitable ordering—is a blow-up-suitable *k*-tuple of subsets of *M* (so that $[M : \mathcal{P}]$ is defined). As always, we shall write

$$\bigcup \mathcal{P} := \bigcup_{P \in \mathcal{P}} P.$$

We shall need the following simple concept of "equivalent functions" on $[M : \mathcal{P}]$.

Definition 4.1 Let M be a manifold with corners, let \mathcal{P} be a blow-up-suitable k-tuple of p-submanifolds of M (so, in particular, $[M : \mathcal{P}]$ is defined), and let $k \in \mathbb{N} \cup \{0, \infty\}$. Assume that we have two continuous functions $f_i : M \to [0, \infty), i = 0, 1$, such that the functions f_i are \mathcal{C}^k on $M \setminus \bigcup \mathcal{P}$ and nowhere vanishing. We shall say that f_0 and f_1 are \mathcal{C}^k -equivalent (on $[M : \mathcal{P}]$) or that f_0 is \mathcal{C}^k -equivalent (on $[M : \mathcal{P}]$) to f_1 if

$$\frac{f_1|_{M\smallsetminus \bigcup \mathcal{P}}}{f_0|_{M\smallsetminus \bigcup \mathcal{P}}}$$

extends to a nowhere vanishing C^k function on [M : P]. In the case $k = \infty$, we shall say that f_0 is *smoothly equivalent* or just *equivalent* to f_1 and write $f_0 \sim f_1$. In the case k = 0, we say f_0 and f_1 are *continuously equivalent*.

The family \mathcal{P} as well as the space $[M : \mathcal{P}]$, which we assumed to be defined, will sometimes be understood, so we may occasionally not mention them. Recall that two functions $f_1, f_2 : U \to [0, \infty)$ are called *Lipschitz equivalent* if there exists C > 0 such that

$$C^{-1}f_1 \le f_2 \le Cf_1. \tag{16}$$

Two functions $f_1, f_2 : M \to [0, \infty)$ are called *locally Lipschitz equivalent* if any point in *M* has an open neighborhood *U* such that $f_1|_U$ and $f_2|_U$ are Lipschitz equivalent. Remark 4.2 Let us record a few easy consequences of the definition:

- (i) Clearly, the \mathcal{C}^k -equivalence of functions on $[M : \mathcal{P}]$ is an equivalence relation.
- (ii) Since $M \setminus \bigcup \mathcal{P}$ is dense in $[M : \mathcal{P}]$ (Proposition 3.10) the extension of the quotient $\frac{f_1|_{M \setminus \bigcup \mathcal{P}}}{f_0|_{M \setminus \bigcup \mathcal{P}}}$ to $[M : \mathcal{P}]$ by continuity is unique (when it exists).
- (iii) Obviously C^k -equivalence implies C^ℓ -equivalence for $\ell < k$.
- (iv) If two functions $f_i : M \to [0, \infty)$ as above are continuously equivalent, then they are locally Lipschitz equivalent. As a consequence, for *M* compact, C^0 equivalence implies Lipschitz equivalence.
- (v) If *H* is a boundary hyperface of *M* with at least one defining function, then any two defining functions of *H* will be equivalent on $[M : H] \simeq M$.

Let us now discuss the type of metrics that we will use. Let M be a manifold with corners. In the following, we shall make use of two kinds of metrics on M. The first kind of metrics will consist of what we are calling "true Riemannian metrics" on M. A *true Riemannian metric* on M is, by definition, nothing but a smooth, fiberwise positive definite symmetric section of $T^*M \otimes T^*M \to M$ (the usual kind of metrics on M). The second kind of metrics will consist of the so-called *compatible metrics*, which will be defined below, Definition 4.20, and whose definition requires some additional data (a structural Lie algebra of vector fields on M). Compatible metrics are Riemannian metrics on $M_0:=M \setminus \partial M$ and do not extend to a true metric on M, unless $\partial M = \emptyset$.

Let *P* be a closed p-submanifold of *M*. Recall that $\beta_{M,P} : [M : P] \to M$ denotes the blow-down map (see Definition 3.1).

Definition 4.3 Let *M* be a manifold with corners and $P \subset M$ be a closed p-submanifold. A function $r_P : M \to [0, \infty)$ will be called *a smoothed distance function to P* (*in M*) if its lift $\beta_{M,P}^*(r_P) := r_P \circ \beta_{M,P}$ is a boundary defining function (Definition 2.4) for $\mathbb{S}N^M_+ P = \beta_{M,P}^{-1}(P)$, the hyperface (or union of hyperfaces) of [M : P] obtained by blowing up *M* along *P*.

We also remark that a smoothed distance function to P in M is continuous since it lifts to a continuous function on [M : P] and M has the quotient topology. The following remark explains the name "smoothed distance function to P" in M for r_P .

Remark 4.4 The function $f : M \to [0, 1]$ is a smoothed distance function to P in M if, and only if, it satisfies the following conditions:

- (i) it is continuous on *M* and smooth on $M \setminus P$,
- (ii) $f^{-1}(\{0\}) = P$, and most importantly,
- (iii) there is a neighborhood V of P such that f is (smoothly) equivalent on [V : P] to the distance to P with respect to some suitable true metric on M.

We omit details here, as this fact will not be used in our article.

Remark 4.5 In general, a smoothed distance to P in M function r_P will *not* be smooth on M; in fact, smoothness on M will only hold if P is empty, or a union of connected components of M. However, it will be continuous on M and smooth on [M : P]. This is one of the main reasons for considering the blow-up [M : P].

Remark 4.6 Given a closed p-submanifold $P \subset M$, we see that a smoothed distance to P in M (Definition 4.3) is *not uniquely determined*. However, any two such smoothed distance functions are equivalent on [M : P], and hence they are equivalent on any other iterated blow-up [M : P] as well, as long as $P \in P$. This is, in fact, our motivation for introducing the notion of equivalence of such functions. For this reason, we will sometimes talk about *the* smoothed distance to P, although we will really mean the equivalence class of the smoothed distances to P in M.

Remark 4.7 In the case that $M = \overline{X}$ is the spherical compactification of some Euclidean vector space X as explained in Sect. 1.1 and if Y is a linear subspace of X, we may consider the Euclidean distance function $d_Y := \min\{||x-y|| \mid y \in Y\}$ introduced in Eq. (2). If $r_{\overline{Y}} : \overline{X} \to [0, \infty)$ is a *smoothed* distance function to \overline{Y} , then the function $r_{\overline{Y}}$ is clearly not Lipschitz equivalent to d_Y , as the first one is bounded and the second one is not. One can also show that $r_{\overline{Y}}$ is also not bi-Lipschitz to arctan $\circ d_Y$, as these two functions behave differently close to \mathbb{S}_X . The precise behavior of $r_{\overline{Y}}$ is that it is continuously equivalent and Lipschitz equivalent to $x \mapsto \arctan(||x||^2 + 1)^{-1/2} d_Y(x))$. Furthermore, $r_{\overline{Y}}$ is continuously equivalent and thus locally Lipschitz equivalent to $x \mapsto (||x||^2 + 1)^{-1/2} d_Y(x)$.

We now consider as in [6] and [37] pairs (P, Q) of closed p-submanifolds of M. As in those papers, we need to consider the cases $Q \subset P$ and $Q \cap P = \emptyset$. We begin with the first case.

Lemma 4.8 Let $P \subset M$ be a closed p-submanifold of the manifold with corners M, let $Q \subset P$ be a closed p-submanifold of P, and let r_P (respectively, r_Q) be a smoothed distance function to P (respectively, to Q) in M. Then,

$$(r_Q^{-1}r_P)\Big|_{M\smallsetminus P}$$

extends to a smoothed distance function to $\beta_{M,Q}^*(P) := \overline{\beta_{M,Q}^{-1}(P \setminus Q)}$ in [M : Q].

Of course, in the above lemma, we have $\beta_{M,Q}^*(P) := \beta_{M,Q}^{-1}(P \setminus Q) \simeq [P : Q]$ see Proposition 3.3. Here \simeq denotes the existence of a diffeomorphism extending the identity on $P \setminus Q$. Also, for the simplicity of the presentation, we may and will assume in the following that $\beta_{M,Q}^*(P) = [P : Q]$.

Remark 4.9 Note that the assumptions in the lemma allow the case when, for all $q \in Q$, we have $\dim_q(Q) = \dim_q(P)$. In this case $\beta_{M,Q}^*(P) = P \setminus Q$ is a union of some connected components of P, and thus for any $q \in Q$ we have $(\beta_{M,Q})^{-1}(q) \cap [P : Q] = \emptyset$. If U is an open neighborhood of q, then any positive function defined on $\beta_{M,Q}^{-1}(U)$ is thus a smoothed distance function to $[P : Q] \cap \beta_{M,Q}^{-1}(U) = \emptyset$.

On the other hand, in this special case, $r_Q^{-1}r_P$, defined on $U \setminus \{q\}$, where U is an open neighborhood of q with $U \cap P = Q$, extends to a positive smooth function on $\beta_{[M:Q],[P:Q]}^*(\beta_{M,Q}^*(U)) = \beta_{M,Q}^*(U)$. The latter statement follows as $r_P|U$ is a smooth distance function to Q in U and using Remark 4.5. These arguments provide a proof of the lemma in this exceptional case. This special case is also included in the following proof, if we use the convention $\mathbb{S}^{-1} = \emptyset$ (here \mathbb{S}^{-1} is the unit sphere in \mathbb{R}^{0}).

Proof of Lemma 4.8 We need to prove that the function $r_Q^{-1}r_P : M \setminus P \to (0, \infty)$, extends to a smooth function on

$$[M:Q,P] := [[M:Q]:[P:Q]]$$

and that this extension of $r_O^{-1}r_P$ is a defining function for the hyperface

$$P' := \mathbb{S}(N_{+}^{[M:Q]}[P:Q]) = \beta_{[M:Q],[P:Q]}^{-1}([P:Q])$$

of [M:Q,P]. Let

$$\beta := \beta_{M,Q,P} := \beta_{M,Q} \circ \beta_{[M:Q],[P:Q]} : [M:Q,P] \to M$$

be the blow-down map and $z \in P'$ (see Definition 3.8). Then, $\beta(z) \in P$.

Recall that by the definition of r_P , we have that $r_P \circ \beta_{M,P}$ is a boundary defining function $r_P \circ \beta_{M,P}$: $[M : P] \to [0, \infty)$ of $\mathbb{S}(N^M_+P) = \beta^{-1}_{M,P}(P) \subset [M : P]$ as a hyperface of [M : P]. The map $[M : Q, P] \to [M : P]$, defined by Lemma 3.4 is a diffeomorphism outside the preimage of Q. Thus, if $\beta(z) \notin Q$, the function $r_Q^{-1}r_P : M \setminus P = [M : Q] \setminus [P : Q] \to [0, \infty)$ extends locally—i.e., in a neighborhood of *z*—to a defining function for $P' \subset [M : Q, P]$, since $r_Q > 0$ at and near *z* and since r_P is a defining function of $\mathbb{S}(N^M_P)$, the pullback of *P* in [M : P], as we have just explained.

On the other hand, if $\beta(z) \in Q$, we use (again) the fact that our problem is local. Thus, by choosing a suitable chart around $\beta(z)$, we can reduce the lemma to the special case

$$\begin{cases} M = \mathbb{R}_{\ell}^{k} \times \mathbb{R}_{\ell'}^{k'} \times \mathbb{R}_{\ell''}^{k''} \ni (x, x', x'') \\ P = \mathbb{R}_{\ell}^{k} \times \mathbb{R}_{\ell'}^{k'} \times \{0\} \ni (x, x', 0) \\ Q = \mathbb{R}_{\ell}^{k} \times \{0\} \times \{0\} \ni (x, 0, 0) \\ \beta(z) = (0, 0, 0). \end{cases}$$

Assume first that $\ell' = \ell'' = 0$. In this very special case, we have

$$[M:Q] = \mathbb{R}_{\ell}^{k} \times \mathbb{S}^{k'+k''-1} \times [0,\infty) \ni (x,\xi,r),$$
$$[P:Q] = \mathbb{R}_{\ell}^{k} \times \mathbb{S}^{k'-1} \times [0,\infty),$$

where $r = \sqrt{(x')^2 + (x'')^2}$ and $\xi = (x', x'')/r$ away from Q. Furthermore, the inclusion $[P : Q] \subset [M : Q]$ is given by the inclusion of $\mathbb{S}^{k'-1} \subset \mathbb{S}^{k'+k''-1}$ on the first k' components of $\mathbb{S}^{k'+k''-1}$ and by the identity on the other factors (that is, on \mathbb{R}^k_ℓ and on $[0, \infty)$). For $\xi \in \mathbb{S}^{k'+k''-1}$, let $\theta(\xi)$ be the length of the shortest geodesic from ξ to

 $\mathbb{S}^{k'-1}$, unless k' = 0. In the case k' = 0, we set $\theta \equiv \pi/2$, thus $\sin \circ \theta \equiv 1$. Then,

$$\hat{r} := \sin \circ \theta \colon \mathbb{S}^{k' + k'' - 1} \to [0, 1]$$

is a smoothed distance function to $\mathbb{S}^{k'-1}$ in $\mathbb{S}^{k'+k''-1}$, thus—by pullback— \hat{r} is also a smoothed distance function to [P:Q] in [M:Q]. Smoothed distance functions r_Q and r_P (for Q and P in M) are then given by

$$r_Q(x, \eta, r) = \sqrt{\|x'\|^2 + \|x''\|^2} = r$$
 and $r_P = \|x''\| = r\sin(\theta(\xi)).$

Since, obviously, $\hat{r} = r_Q^{-1} r_P$, we obtain the desired statement if $\ell' = \ell'' = 0$. The general case follows by replacing $\mathbb{S}^{k'-1}$ with $\mathbb{S}_{\ell'}^{k'-1}$ and $\mathbb{S}^{k'+k''-1}$ with $\mathbb{S}^{k'+k''-1} \cap (\mathbb{R}_{\ell'}^{k'} \times \mathbb{R}_{\ell''}^{k''})$.

We now turn to the second case, that when P and Q are disjoint. For later use, we prove a more general statement.

Lemma 4.10 Let M be a compact manifold with corners, let $P \subset M$ be a closed p-submanifold and $Q = (Q_1, Q_2, ..., Q_k)$ be a blow-up-suitable k-tuple of closed subsets of M disjoint from P. Let $\beta : [M : Q] \to M$ be the blow-down map and let r_P be a smoothed distance to P (in M). Let $\widehat{P} := \beta^{-1}(P) = \beta^*(P)$. Then, $r_P \circ \beta$ is a smoothed distance function to \widehat{P} in [M : Q].

Proof Let us prove our result by induction on k. For k = 0 (i.e., for $Q = \emptyset$) there is nothing to prove according to our conventions for the blow-up with respect to an empty family. Let us write $Q = Q_1$, for the simplicity of the notation. Let us prove our result for k = 1. The blow-down $[M : Q, P] \rightarrow [M : P]$ induces a diffeomorphism $[M : Q, P] \setminus \beta^{-1}(Q) \rightarrow [M \setminus Q : P]$ (see Lemma 3.6). The function $r_P \circ \beta_{M,Q}$ is a smoothed distance function to $\beta^{-1}(P)$ on $M \setminus Q \subset [M : Q]'$ (since it coincides with r_P there). Moreover, $r_P \circ \beta_{M,Q}$ is smooth everywhere on the iterated blow-up [M : Q, P] and vanishes only on the preimage of P (in particular, it is > 0 on the preimage of Q, which is disjoint from the preimage of P). Hence, $r_P \circ \beta_{M,Q}$ is a smoothed distance function to $\beta^*_{M,Q}(P) = \beta^{-1}_{M,Q}(P)$ in [M : Q]. The induction step is always to perform an additional blow-up, so it reduces to the case k = 1 just proved.

4.2 Smoothed distance functions to a family

We need to extend the definition of the smoothed distance function to families. (The reader may want to review Definition 3.8 at this point.)

Definition 4.11 Let $\mathcal{P}:=(P_i)_{i=1}^k$, $P_i \subset M$ be a blow-up-suitable *k*-tuple (that is, an ordered family of closed subsets of M such that the iterated blow-up $[M : P_1, P_2, \ldots, P_k]$ is defined, Definition 3.7). Let $\beta : [M : P_1] \to M$ be the blow-down map and $\mathcal{P}':=(\beta^*(P_2), \beta^*(P_3), \ldots, \beta^*(P_k))$, which, we recall, is a family of

closed p-submanifolds of $[M : P_1]$. We then define a smoothed distance function $\rho_{\mathcal{P}}$ to \mathcal{P} in M to be a function $\rho_{\mathcal{P}} : M \to [0, \infty)$ given by induction on k by the formula

$$\rho_{\mathcal{P}}(x) := \begin{cases} r_{P_1}(x) & \text{if } k = 1\\ r_{P_1}(x)\rho_{\mathcal{P}'}(y) & \text{if } k > 1 \text{ and } y \in \beta^{-1}(\{x\}), \end{cases}$$

where $r_{P_1}: M \to [0, \infty)$ is a smoothed distance function to P_1 in M (Definition 4.3) and $\rho_{\mathcal{P}'}$ is a smoothed distance function to \mathcal{P}' in $[M : P_1]$. (We note that the last expression is well defined since, for $x \notin P_1$, y is unique whereas, for $x \in P_1$, we have $r_{P_1}(x) = 0$.)

Again, we use the notation introduced in Definition 3.8 (which is, in turn, the same as the one in [6, Definition 4.1]). We have then, similarly to [6, Remark 4.3].

Remark 4.12 We have that $\rho_{\mathcal{P}}$ is continuous on M and that $\rho_{\mathcal{P}}^{-1}(\{0\}) = \bigcup \mathcal{P} = \bigcup_{j=1}^{k} P_j$. Let $\gamma_0 = \text{id}, \gamma_1 := \beta_1^* \text{ and } \gamma_j := \beta_j^* \circ \gamma_{j-1} = \beta_j^* \circ \dots \circ \beta_1^*$, where

$$\beta_k := \beta_{[M:P_1,\dots,P_{k-1}],[P_k:P_1,\dots,P_{k-1}]} : [M:P_1,\dots,P_k] \to [M:P_1,\dots,P_{k-1}].$$

If $x \notin \bigcup \mathcal{P}$, then

$$\rho_{\mathcal{P}}(x) = r_{\gamma_0^*(P_1)}(x)r_{\gamma_1^*(P_2)}(x)\dots r_{\gamma_{k-1}^*(P_k)}(x).$$

In particular, it follows that the pullback of $\rho_{\mathcal{P}}$ to $[M : \mathcal{P}]$ is smooth, or equivalently, suppressing all pullbacks from the notation, $\rho_{\mathcal{P}} \in C^{\infty}([M : \mathcal{P}])$. However, note that in contrast to the non-iterated case, $\rho_{\mathcal{P}} \in C^{\infty}([M : \mathcal{P}])$ is no longer a boundary defining function for some boundary hypersurface of $[M : \mathcal{P}]$.

Remark 4.13 As in Remark 4.6, a smoothed distance to \mathcal{P} function $\rho_{\mathcal{P}}$ (see Definition 4.11) is not unique, but it is unique *up to equivalence* on $[M : \mathcal{P}]$. Indeed, this follows from Remark 4.12 since all the factors $r_{\gamma_{k-1}^*}(P_k)$ are uniquely determined up to equivalence on $[M : \mathcal{P}]$ and the equivalence is compatible with products and preserved when we increase \mathcal{P} .

The following result extends to smoothed distance functions the compatibility with products of Proposition 3.13.

Proposition 4.14 Let M and M_1 be two manifolds with corners and $\mathcal{P} = (P_i)_{i=1}^k$ be a blow-up-suitable k-tuple of subsets of M. Then,

$$\mathcal{P} \times M_1 := (P_j \times M_1)_{j=1}^k$$

is a blow-up-suitable k-tuple of subsets of $M \times M_1$. Let $r_{\mathcal{P}}$ be a smoothed distance to \mathcal{P} in M and $p: M \times M_1 \to M$ the projection. Then, $r_{\mathcal{P}} \circ p$ is a smoothed distance function to $\mathcal{P} \times M_1$ in $M \times M_1$.

Proof If *H* is a hyperface of *M* and r_H is a defining function for *H* in *M*, then $r_H \circ p$ is a defining function for the hyperface $H \times M_1$ in $M \times M_1$. The result follows from definitions by induction on *k* using repeatedly this observation. Indeed, there is nothing to prove if k = 0. The case k = 1 follows from Proposition 3.13 (used also for k = 1) and the observation about hyperfaces. The induction step is completely similar to the case k = 1 and it reduces to that one, as in the proof of Proposition 3.13, using that proposition, the relation $\beta^*(P \times M_1) = \beta^*(P) \times M_1$, and the definition of smoothed distance functions.

From now on, we shall assume that our *k*-tuple is a clean semilattice of closed p-submanifolds of M, endowed with an admissible order. Also, from now on we shall denote our *k*-tuple with S instead of P in order to stress that it is stable for intersections (i.e., that S is a semilattice). (Recall that we can choose any admissible order on S.) The following proposition will play a crucial role in our application to *N*-body-type problems.

Proposition 4.15 Let S be a clean semilattice of closed p-submanifolds of a connected manifold with corners M and $\emptyset \neq Y \in S$. Let r_Y be a smoothed distance function to Y in M (Definition 4.3) and ρ_S be a smoothed distance function to S in M (Definition 4.11). Then, the function $\rho_S/r_Y : M \setminus \bigcup S \to (0, \infty)$ extends to a smooth function on [M : S].

Note that in the statement of the proposition, it is important that r_Y is a smoothed distance in M and not in some blow-up of M. The proposition would be a trivial consequence of the recursive definition of ρ_S if we replaced r_Y by a smooth distance function in some suitable blow-up of M.

Proof There is no loss of generality to assume that $\emptyset \in S$, so we will do so for brevity. (The case $\emptyset \notin S$ is completely similar.) We shall prove the statement by induction on the number of elements of S.

If S has only one non-trivial element, that is, if $S = (\emptyset, Y)$, then we have $\rho_S \simeq r_Y$, by the definition of ρ_S , as explained in Remarks 4.6 and 4.13.

Now, for the induction step, let us arrange $S = (P_0 = \emptyset, P_1, P_2, \dots, P_k)$ in an admissible order. Let us blow up along $P_1 \in S$ (which is a minimal element of $S \setminus \{\emptyset\}$, by the definition of an admissible order). We use the notation introduced in Definition 4.11, in particular S' consists of the lifts to $[M : P_1]$ of the p-submanifolds P_2, \dots, P_k . If $Y = P_1$, then the result follows from the formula for ρ_S in Remark 4.12. Let us assume therefore that $Y \neq P_1$. Since P_1 is a minimal (non-empty) element of S, we have two possibilities: either $P_1 \subset Y$ or $P_1 \cap Y = \emptyset$.

Let us assume that we are in the case $P_1 \subset Y$. We choose then as a smoothed distance for $[Y : P_1]$ in $[M : P_1]$ the function $r'_{[Y:P_1]} := r_Y/r_{P_1}$, which is possible in view of Lemma 4.8. By the induction hypothesis, we know that $(r_Y/r_{P_1})^{-1}\rho_{S'}$ extends to a smooth function on $[[M : P_1] : S']$.

Since $[[M : P_1] : S'] = [M : S]$, by the definition of the iterated blow-up, by suppressing pullbacks in notation we obtain that

$$r_{Y}^{-1}\rho_{\mathcal{S}} = r_{Y}^{-1}r_{P_{1}}\rho_{\mathcal{S}'} = (r_{Y}/r_{P_{1}})^{-1}\rho_{\mathcal{S}'} \in \mathcal{C}^{\infty}([M:\mathcal{S}]).$$
(17)

🖄 Springer

This equation should be understood in the sense of functions $M \setminus \bigcup S \to \mathbb{R}_+$ that extend to smooth functions on [M : S]. Thus, we have obtained the desired relation, so the proof is complete in the case $P_1 \subset Y$.

Let us assume now that $P_1 \cap Y = \emptyset$. The proof is then an immediate application of Lemma 4.10 for k = 1.

The following remark recalls [1, Lemma 3.17].

Remark 4.16 Recall that in this and in the following sections, S is a clean semilattice of closed p-submanifolds of M. Let $\rho(x) := \operatorname{dist}_{\overline{\rho}}(x, \mathcal{S}) := \operatorname{dist}_{\overline{\rho}}(x, \bigcup \mathcal{S})$ be the distance to S in some true metric \overline{g} on M. (That is, a metric on TM that extends smoothly to the boundary of M, unlike a "compatible metric," a concept that will be introduced shortly and which is really only a metric on $M \setminus \partial M$ with some additional properties.) One can show that the functions ρ_{S} and ρ are continuously equivalent. A special case of this statement which is in fact the only case needed in the present article was proved in [1, Lemma 3.17] based on preliminary work in [8]. In [1] it is additionally assumed that all p-submanifolds in S are *interior submanifolds*. By definition, an interior submanifold is a p-submanifold of positive boundary depth, i.e., no component is contained in ∂M . However, a proof may be extended to the full statement claimed above; the result extends to p-submanifolds that are not interior, either by adapting the proofs or by embedding the manifold M into a Riemannian manifold N without corners and boundary, such that all boundary hypersurfaces of M are totally geodesic. The function ρ , however, is not smooth on $M \setminus \bigcup S$, in general. This justifies the notations ρ and ρ_S , since they are very similar in scope. See Lemma A.2 for an extension of this remark.

4.3 Blow-ups and Lie manifolds

We now look at a class of differential operators that will "desingularize" the *N*-body Hamiltonian (with Coulomb or inverse square potentials). Let thus *M* be a manifold with corners and $\mathcal{V} \subset \mathcal{C}^{\infty}(M; TM)$ be a subspace of smooth vector fields on *M* that is stable for the Lie bracket and for multiplication with functions in $\mathcal{C}^{\infty}(M)$. We then let $\text{Diff}_{\mathcal{V}}^m(M)$ denote the space of differential operators of order $\leq m$ on *M* generated by $\mathcal{C}^{\infty}(M)$ and by \mathcal{V} and

$$\operatorname{Diff}_{\mathcal{V}}(M) := \bigcup_{m \in \mathbb{N}_0} \operatorname{Diff}_{\mathcal{V}}^m(M) = \left\{ \sum a V_1 V_2 \dots V_k \mid a \in \mathcal{C}^{\infty}(M), V_j \in \mathcal{V} \right\}.$$
(18)

Furthermore, we will assume that (M, \mathcal{V}) is a "Lie manifold," a concept from [4] that we now recall.

Definition 4.17 Let *M* be a smooth, compact manifold with corners and $\mathcal{V} \subset \mathcal{C}^{\infty}(M; TM)$ be a subspace of smooth vector fields on *M*. In addition to the relations $\mathcal{C}^{\infty}(M)\mathcal{V} = \mathcal{V}$ and $[\mathcal{V}, \mathcal{V}] \subset \mathcal{V}$, let us also assume that:

(i) $\mathcal{C}^{\infty}_{c}(M \smallsetminus \partial M; TM) \subset \mathcal{V};$

(ii) all vector fields in \mathcal{V} are tangent to the boundary; and

(iii) \mathcal{V} has a $\mathcal{C}^{\infty}(M)$ -local basis near each point.

Then, we shall say that $(M; \mathcal{V})$ is a *Lie manifold*; the space \mathcal{V} will be called its associated *structural Lie algebra of vector fields*.

Remark 4.18 The condition (iii) in the above definition means that any $p \in M$ has a closed neighborhood U such that the restriction of \mathcal{V} to U is a free $\mathcal{C}^{\infty}(U)$ -module. An even better way of expressing condition (iii) is to assume the existence of a vector bundle $A \to M$ together a vector bundle morphism $\varrho : A \to TM$ over id_M with $\varrho(\Gamma(A)) = \mathcal{V}$. The morphism ϱ is called an *anchor* map, see [4, 40] and the references therein. Then, Condition (i) is equivalent to saying that ϱ restricts to a vector bundle isomorphism on $M_0 := M \setminus \partial M$. Similarly, Condition (ii) is equivalent to saying that, for any $p \in \partial M$ of boundary depth 1, we have $\varrho(A_p) \subset T_p(\partial M) \subsetneq T_pM$. This condition then implies analogous tangency conditions for points of boundary depth ≥ 2 .

The following simple and basic example(s) will play a crucial role in our applications to the *N*-body problem.

Example 4.19 We continue to assume that X is a finite-dimensional real vector space and that \overline{X} is its spherical compactification. We shall take \overline{X} for our ambient manifold (thus \overline{X} plays the role of the manifold typically called M so far). Then, X identifies with the space of constant vector fields on itself. Corollary B.3 tells us that these vector fields extend to vector fields on \overline{X} and that these extended vector fields vanish on \mathbb{S}_X . We shall consider then the structural Lie algebra of vector fields $\mathcal{V}:=\mathcal{C}^{\infty}(\overline{X})X$. Then, the pair $(\overline{X}; \mathcal{V})$ is a Lie manifold (Definition 4.17). Indeed, \mathcal{V} is a free $\mathcal{C}^{\infty}(\overline{X})$ module since any basis of X, viewed as a vector space, also provides a basis of \mathcal{V} as a $\mathcal{C}^{\infty}(\overline{X})$ -module. The resulting algebra of differential operators $\text{Diff}_{\mathcal{V}}(\overline{X})$ is the algebra of differential operators on X with coefficients in $\mathcal{C}^{\infty}(\overline{X})$.

To a Lie manifold $(M; \mathcal{V})$, there is canonically associated a class of metrics on the interior of M, called \mathcal{V} -compatible metrics on $M_0 = M \setminus \partial M$, as follows:

Definition 4.20 Let (M, \mathcal{V}) be a Lie manifold, $n = \dim(X)$. A \mathcal{V} -compatible metric on M is defined as a Riemannian metric on M_0 with the following property: each $x \in M$ has a closed neighborhood U_x such that the restriction of \mathcal{V} to U_x has a $\mathcal{C}^{\infty}(U_x)$ -basis (v_1, v_2, \ldots, v_n) that is orthonormal in the sense

 $\forall y \in M_0 \cap U_x$: (v_1, v_2, \dots, v_n) is an orthonormal basis of $(T_y M, g_y)$.

(Of course, this is a true restriction on g and \mathcal{V} only at the boundary of M).

A \mathcal{V} -compatible metric on M will never be a true metric on M since the induced metric on TM_0 will not extend continuously to M. For instance, for Example 4.19 (and the related Example 5.2 relevant for the N-body problem), any Euclidean metric on X will be a \mathcal{V} -compatible metric on M.

Remark 4.21 An alternative way to define a \mathcal{V} -compatible is as follows. We briefly recall this, as it will be used in "Appendix A." See [4] for a proof of the equivalence of the definitions. Let (M, \mathcal{V}) be a Lie manifold with anchor map $\varrho : A \to TM$ as defined above. A compatible metric may be defined as a metric on A, which is by definition a symmetric positive definite section $\gamma \in \Gamma(A^* \otimes A^*)$. The anchor map ϱ induces a map $\rho_* = \rho \otimes \rho : A \otimes A \to TM \otimes TM$, and on M_0 the inverse of the anchor map, namely $(\varrho|_{M_0})^{-1}$, may be dualized to a pullback map $((\varrho|_{M_0})^{-1})^* : A^*|_{M_0} \to T^*M_0$. Then, $g := ((\varrho|_{M_0})^{-1})^* \otimes ((\varrho|_{M_0})^{-1})^* (\gamma)$ is a Riemannian metric on M_0 . A Riemannian metric on M_0 is \mathcal{V} -compatible if, and only if it arises this way.

If $B \in V^* \otimes V^*$ is a symmetric bilinear form on a finite-dimensional vector space V, then we denote $\flat_B : V \to V^*$, $X \mapsto B(X, \cdot)$ the associated canonical homomorphism. If B is non-degenerate, \flat_B is invertible and one defines $\sharp_B := \flat_B^{-1}$. In this case we can define ${}^{\natural}B^{\sharp} := B(\sharp_B(\cdot), \sharp_B(\cdot)) \in V \otimes V$ whose matrix with respect to some basis of V is the inverse of the matrix of B with respect to its dual basis. Now, if γ is a metric on the vector bundle A, we map apply this construction for any $q \in M$ to $\gamma|_q \in A_q^* \otimes A_q^*$, and we obtain a smooth section ${}^{\natural}\gamma^{\sharp} \in \Gamma(A \otimes A)$. Then, $G := (\rho \otimes \rho)({}^{\natural}\gamma^{\sharp}) \in \Gamma(TM \otimes TM)$ is a well-defined smooth tensor. One can easily show that G is the unique continuous extension of ${}^{\natural}g^{\sharp}$ for the g defined above. We will use this description to proof in "Appendix A" for a true metric \overline{g} and a compatible metric g we have, see Lemma A.2, a constant C with $\overline{g} \leq Cg$ in the sense of symmetric forms, i.e., $Cg - \overline{g}$ is positive semi-definite.

Any two \mathcal{V} -compatible metrics g and \tilde{g} are Lipschitz equivalent, i.e., there exists a constant C > 0 with $C^{-1}\tilde{g} \le g \le C\tilde{g}$, again in the sense of symmetric forms. Thus, the volume forms are Lipschitz equivalent as well, and hence the definition of the space $L^p(M, g) = L^p(M; \mathcal{V})$ only depends on \mathcal{V} , and not on the metric. This allows us to define the associated *Sobolev spaces* on X (or M) as

$$W^{k,p}(M;\mathcal{V}) = \left\{ u \mid V_1 \dots V_j u \in L^p(M;\mathcal{V}), \forall V_1, \dots, V_j \in \mathcal{V}, \ j \in \{0, 1, \dots, k\} \right\} (19)$$

where $k \in \mathbb{N}_0$ and $1 \le p \le \infty$. Obviously, they do not depend on the choice of \mathcal{V} compatible metric. These Sobolev spaces coincide with the standard Sobolev spaces
on the complete Riemannian manifold associated with any compatible metric.

The main reason we are interested in Lie manifolds is that there is also a notion of an associated pseudodifferential calculus [5], a notion of ellipticity in $\text{Diff}_{\mathcal{V}}(M)$, and, most importantly for this paper, an elliptic regularity result [3]. Explicitly, an operator in $\text{Diff}_{\mathcal{V}}^m(M)$ is *elliptic* in $\text{Diff}_{\mathcal{V}}^m(M)$ if its principal symbol is uniformly elliptic in one (equivalently, any) \mathcal{V} -compatible metric.

The following theorem is the special case s = 0 and $m \in \mathbb{Z}_+$ of [3, Theorem 8.7].

Theorem 4.22 (Ammann–Ionescu–Nistor) Let $(M; \mathcal{V})$ be a Lie manifold. Let $k, j \in \mathbb{Z}$, $m \in \mathbb{Z}_+, 1 , and <math>P \in \text{Diff}_{\mathcal{V}}^m(M)$ be elliptic. Let $u \in W^{k,p}(M; \mathcal{V})$ be such that $Pu \in W^{j,p}(M; \mathcal{V})$. Then, $u \in W^{j+m,p}(M; \mathcal{V})$.

Let us mention also that for in Example 4.19 (and the related Example 5.2 relevant for the *N*-body problem), an operator $P \in \text{Diff}_{\mathcal{V}}(M)$ is elliptic in that algebra if, and only if, it is uniformly elliptic (in the usual, Euclidean metric of *X*). In particular, the above regularity theorem (Theorem 6.1) can also be obtained from the regularity result in [2].

Let us now provide an alternative, more general method to obtain a Lie manifold structure on the Georgescu–Vasy space X_{GV} . (See the Introduction or Sect. 5 (22) for the definition of X_{GV} .)

Remark 4.23 Let $(M; \mathcal{V})$ be a Lie manifold and $Y \subset M$ be a closed p-submanifold. We assume $Y \subset \partial M$, i.e., Y is everywhere of positive boundary depth. We also assume that all $V \in \mathcal{V}$ satisfy $V|_Y \in \Gamma(TY)$. The latter condition implies (due to a straightforward modification of [1, Proposition 3.2]) that any $V \in \mathcal{V}$ has a *lift* \tilde{V} , that is, a vector field on [M : Y] such that

commutes. As a consequence, the inclusion $\mathcal{V} \hookrightarrow \Gamma(TM)$ "lifts" to a map $\mathcal{V} \hookrightarrow \Gamma(T[M : Y])$, both maps are injective Lie algebra homomorphisms and $\mathcal{C}^{\infty}(M)$ -linear. We obtain a Lie manifold structure on [M : Y] by taking as structural Lie algebra of vector fields

$$\mathcal{W}_{\text{bdry}} := \mathcal{C}^{\infty}([M:Y]\mathcal{V}.$$
(20)

Let us record here how the various quantities associated with the Lie manifold (M, \mathcal{V}) change when going to the blow-up Lie manifold $([M : Y], \mathcal{W}_{bdry})$:

- The interior smooth manifold is the same: $M_0 = M \setminus \partial M = [M : Y] \setminus \partial [M : Y]$.
- If a metric on M_0 is \mathcal{V} -compatible, then it is also \mathcal{W}_{bdry} -compatible.
- $W^{k,p}(M; \mathcal{V}) = W^{k,p}([M:Y]; \mathcal{W}_{bdry}).$
- $\operatorname{Diff}_{\mathcal{W}_{\operatorname{bdry}}}([M:Y]) = \mathcal{C}^{\infty}([M:Y]) \operatorname{Diff}_{\mathcal{V}}(M).$

The last equality is based on the fact that the lift of \mathcal{V} acts on $\mathcal{C}^{\infty}([M : Y])$ by derivations. By iterating this construction, we can obtain similarly a Lie manifold structure on $[M : \mathcal{Y}]$, where \mathcal{Y} is a clean semilattice of p-submanifolds of ∂M . Indeed, let us assume that \mathcal{V} is tangent to each manifold $Y \subset \mathcal{Y}$, then \mathcal{V} will lift to all intermediate blow-ups leading to $[M : \mathcal{Y}]$. At each step, by density, the lifting of \mathcal{V} will be tangent to the manifold with respect to which we blow up.

One typically describes a Lie manifold via the local behavior of the vector fields near the boundary points. This is not what we did in our case (in the remark right above and the one following next). See [10, 12, 39] for similar Lie manifold structures that are described locally.

Example 4.24 Recall from the Introduction that \mathcal{F} is a finite semilattice of linear subspaces of X such that $\{0\} \in \mathcal{F}$ and $X \notin \mathcal{F}$. The Lie manifold structure on X_{GV} will be obtained starting from \overline{X} by taking $\mathcal{Y}:=\mathbb{S}_{\mathcal{F}}:=\{\mathbb{S}_{Y} \mid Y \in \mathcal{F}\}$. Since the action of X on the boundary \mathbb{S}_{X} of \overline{X} is trivial, see Remark 4.19 and "Appendix C," the vector fields in $\mathcal{V}:=\{f V \mid f \in \mathcal{C}^{\infty}(\overline{X}), V \in X\}$ are tangent to all the submanifolds of $\mathbb{S}_{\mathcal{F}}$.

The situation described in the last remark changes, however, dramatically if Y has boundary depth 0, i.e., if no connected component of Y is contained of the boundary. In that case, there is, in principle, more work to be done, but that has already been completed to a large extent in [1]. Let us summarize some required results from that article.

Remark 4.25 We use the notation introduced in Remark 4.23. Unlike there, however, we shall consider, this time, the case when Y is connected and has boundary depth 0 (that is, it is not contained in the boundary ∂M of M). Let r_Y be a smoothed distance function to Y, Definition 4.3. We then obtain a Lie manifold structure on [M : Y] as in [1] by taking as structural Lie algebra of vector fields

$$\mathcal{W}_{\text{int}} := r_Y \mathcal{C}^{\infty}([M:Y]) \mathcal{V}.$$
(21)

See [1, Lemma 3.8 and Theorem 3.10] where it was proved that all vector fields in $r_Y \mathcal{V}$ lift canonically to smooth vector fields on [M : Y]. (This justifies the additional factor r_Y , since without it, that would not cover our case.) Let us record here how the various quantities associated with the Lie manifold (M, \mathcal{V}) change when going to the blow-up Lie manifold $([M : Y], \mathcal{W}_{int})$:

• The interior smooth manifold of the blow-up is, this time:

$$M_1 := [M : Y] \setminus \partial [M : Y] = M \setminus (\partial M \cup Y).$$

- If g is a \mathcal{V} -compatible metric on M, then $\tilde{g} := r_Y^{-2} \beta_{M,Y}^* g$ is a \mathcal{W}_{int} -compatible metric on [M : Y].
- The volume forms for g and \tilde{g} are related by the formula $dvol^{\tilde{g}} = r_Y^{-n}\beta^*_{[M:Y]} dvol^g$, $n = \dim(M)$. As a consequence we have

$$u \in L^p([M:Y]; \mathcal{W}_{int}) \iff r_Y^{-n/p} u \in L^p(M; \mathcal{V})$$

 The Sobolev spaces defined by W_{int} are "weighted Sobolev spaces" in the old metric (compare to Equation (19) and notice the factor r^j_V):

$$W^{k,p}([M:Y], \mathcal{W}_{int}) = \left\{ u \mid r_Y^j V_1 \dots V_j u \in L^p([M:Y]; \mathcal{W}_{int}) \\ \forall V_1, \dots, V_j \in \mathcal{V}, \ j \in \{0, 1, \dots, k\} \right\}$$

• Let $m \in \mathbb{Z}_+$, then $\text{Diff}_{\mathcal{W}_{\text{int}}}^m([M:Y])$ is the linear span of differential monomials of the form $r_Y^j a V_1 V_2 \dots V_j$, with $a \in \mathcal{C}^{\infty}([M:Y]), V_1, V_2, \dots, V_j \in \mathcal{V}$, and $0 \le j \le m$. Moreover, if $D \in \text{Diff}_{\mathcal{V}}^{m}(M)$, then $r_{Y}^{m}D \in \text{Diff}_{\mathcal{W}_{\text{int}}}^{m}([M:Y])$ and, if D is elliptic in $\text{Diff}_{\mathcal{V}}^{m}(M)$, then $r_{Y}^{m}D$ is elliptic in $\text{Diff}_{\mathcal{W}_{\text{int}}}^{m}([M:Y])$.

The last two statements are also obtained using that $V(r_Y) \in C^{\infty}([M : Y])$ for all $V \in \mathcal{V}$. (We note in passing that $V(r_Y) \notin C^{\infty}(M)$.)

As in Remark 4.23, we can iterate the construction of Remark 4.25, to obtain similarly a Lie manifold structure on $[M : \mathcal{Y}]$, where \mathcal{Y} is a clean semilattice (i.e., a cleanly intersecting semilattice) of p-submanifolds of M. All the resulting objects on $[M : \mathcal{Y}]$ will be independent on the admissible order chosen on \mathcal{Y} .

5 The Lie manifolds associated with the generalized N-body problem

We now investigate how the constructions of the previous section particularize to the case of *N*-body problems. Let us recall our standing conventions that *X* is a finite-dimensional, real vector space and that \mathcal{F} is a finite semilattice of *linear* subspaces of *X* satisfying $\{0\} \in \mathcal{F}$ and $X \notin \mathcal{F}$. (We stress, however, that the assumptions $\{0\} \in \mathcal{F}$ and $X \notin \mathcal{F}$ are not essential, since the general case of a finite semilattice \mathcal{F} easily reduces to this case.)

5.1 Semilattices and blow-ups for the N-body problem

We begin by introducing the semilattices that we will use for the study of the *N*-body problem. These semilattices will then be used to introduce the associated blow-ups and Lie manifolds.

5.1.1 The semilattices and the blow-ups

We denote by {0} the vector space consisting of just $0 \in X$. We agree that $\mathbb{S}_{\{0\}} = \emptyset$, and hence $\overline{\{0\}} = \{0\}$. As in [1, 6], we shall consider the semilattices $\overline{\mathcal{F}} := \{\overline{Y} \mid Y \in \mathcal{F}\}$ and $\mathbb{S}_{\mathcal{F}} := \{\mathbb{S}_Y \mid Y \in \mathcal{F}\}$, recalled earlier in Eq. (4). Note that $\emptyset \in \mathbb{S}_{\mathcal{F}}$ because $0 \in \mathcal{F}$.

Proposition 5.1 The sets $\overline{\mathcal{F}}$, $\mathbb{S}_{\mathcal{F}}$, $\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$ are clean semilattices of *p*-submanifolds of \overline{X} .

Proof In view of Remark 2.13, this result is obtained by a direct application of Lemma 2.15. \Box

We fix an admissible order on $\mathbb{S}_{\mathcal{F}}$, but the constructions below do not depend on this choice. For instance, the *Georgescu–Vasy space*

$$X_{\rm GV} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \tag{22}$$

does not depend on the choice of an admissible order on $\mathbb{S}_{\mathcal{F}}$. (This space is the same as the one introduced in [6] and in Eq. (5).) We now introduce the Lie manifold structure on X_{GV} .

Example 5.2 Let *X* act on \overline{X} by translations, as in Example 4.19. Let also S be a finite, clean semilattice of closed submanifolds of $\mathbb{S}_X = \overline{X} \setminus X$. The Lie group action of *X* on \overline{X} acts trivially on \mathbb{S}_X , thus it preserves S, and therefore we obtain an action of *X* on the blown-up space $[\overline{X} : S]$ (where the blow-up is defined using any admissible order on S). As explained in Lemma B.4, this implies that any constant vector field on *X* extends to a smooth vector field on $M := [\overline{X} : S]$ that is *tangent* to the boundary. Let $W_{bdry} := \mathcal{C}^{\infty}(M)X$.

As in Example 4.19, then $(M; W_{bdry})$ is a Lie manifold (Definition 4.17). Indeed, again, W_{bdry} is a free $C^{\infty}(M)$ -module, a basis being given by extensions of the canonical basis, viewed as constant vector fields. In this case, the resulting algebra of differential operators $\text{Diff}_{W_{bdry}}(M)$ is the algebra of differential operators on X with coefficients in $C^{\infty}(M)$. So the blow-up of \overline{X} by S leads to a larger algebra of differential operators than the one associated with the Lie algebroid $(\overline{X}, V), V := C^{\infty}(\overline{X})X$, considered in Example 4.19. The class of differential operators to which our regularity results apply, however, is even larger than $\text{Diff}_{W_{bdry}}(M)$ (it allows also for interior blow-ups).

For us, the relevant choice is $S = S_F$, which yields $[\overline{X} : S_F] =: X_{GV}$. (Note that X_{GV} was defined using only boundary blow-ups.) We thus have obtained a *Lie* manifold structure on the Georgescu–Vasy space X_{GV} . This class of examples appears implicitly (but prominently) in Georgescu's work [26, 27].

We now proceed to further blow-up X_{GV} with respect to $\overline{\mathcal{F}}$, i.e., the lifts of elements of \mathcal{F} .

Lemma 5.3 Let us arrange both $\mathbb{S}_{\mathcal{F}}$ and $\overline{\mathcal{F}}$ in admissible orders (see Definition 3.14). We then use the resulting order to obtain an order relation on $\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$ in which all the elements of $\mathbb{S}_{\mathcal{F}}$ precede the ones of $\overline{\mathcal{F}}$. Then, the resulting order on the union $\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$ is an admissible order.

Proof Let $\overline{Y} \in \overline{\mathcal{F}}$, where $Y \in \mathcal{F}$. Let also $\mathcal{P} \subset \mathbb{S}_{\mathcal{F}}, \mathcal{P} \neq \mathbb{S}_{\mathcal{F}}$, and $\beta : [\overline{X} : \mathcal{P}] \to \overline{X}$ be the blow-down map. Then, $\beta^*(\overline{Y})$ will contain *Y* and hence cannot be a minimal element of $(\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}) \setminus \mathcal{P}$ since it is not contained in any element of $\mathbb{S}_{\mathcal{F}} \setminus \mathcal{P}$. \Box

We shall need for the following result the following notation for the lifts of \mathcal{F} to X_{GV} . Let $\beta_{\text{GV}} : X_{\text{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \to \overline{X}$ be the blow-down map. If $Y \in \mathcal{F}$, we let

$$\widehat{Y} := \beta_{\rm GV}^* \overline{Y} \subset X_{\rm GV} \quad \text{and} \\ \widehat{\mathcal{F}} := \{\widehat{Y} \mid Y \in \mathcal{F}\} = \beta_{\rm GV}^* \overline{\mathcal{F}}.$$
(23)

(In general, later on, we shall denote by \widehat{Y} the lift of \overline{Y} to any intermediate blow-up between $X_{\text{GV}} := [X : \mathbb{S}_{\mathcal{F}}]$ and $[\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}]$.)

The independence of the blow-up on the admissible order [6] (a result reminded in Theorem 3.18) gives the following:

Corollary 5.4 We have canonical diffeomorphisms

$$X_{\mathcal{F}} := [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}] \simeq \left[[\overline{X} : \mathbb{S}_{\mathcal{F}}] : \widehat{\mathcal{F}} \right] = [X_{\mathrm{GV}} : \widehat{\mathcal{F}}],$$

where $\widehat{\mathcal{F}}$ is the lift of $\overline{\mathcal{F}}$ to $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$.

Remark 5.5 Let us assume that we have an order $\mathcal{F} = (Y_0 = \{0\}, Y_1, \dots, Y_k)$ on the linear subspaces of \mathcal{F} such that $Y_i \subsetneq Y_j \Rightarrow i < j$ (that is, we have a "size-order" on \mathcal{F} , in the sense of Kottke [37]). Then, this is an admissible order on \mathcal{F} and the induced orders on $\mathcal{S}_{\mathcal{F}}, \overline{\mathcal{F}}$, and $\widehat{\mathcal{F}}$ are again admissible.

So $X_{\mathcal{F}}$ is obtained from X_{GV} using only *interior* blow-ups. By contrast, in [1], the process of blowing up along the collision planes $\overline{\mathcal{F}}$ started from \overline{X} to arrive at $[\overline{X} : \overline{\mathcal{F}}]$. This had the disadvantage that the smooth functions on $\overline{X/Y}$ do not lift to smooth functions on $[\overline{X} : \overline{\mathcal{F}}]$. Our paper thus fixes this issue from [1] and provides uniform estimates at infinity.

5.1.2 Boundary blow-ups and the data on X_{GV}

Let us now see how the constructions of Lie manifolds and of their geometric objects of the previous section particularize to our setting, taking into account the two types of blow-ups: boundary blow-up (along a p-submanifold contained in the boundary) and interior blow-up (along a p-submanifold **not** contained in the boundary). We begin with the blow-up of \overline{X} with respect to $\mathbb{S}_{\mathcal{F}}$, which, we recall, is obtained using a sequence of boundary blow-ups. We will let \mathcal{W}_{eucl} denote the structural Lie algebra of vector fields on \overline{X} .

Remark 5.6 We will now define a Lie manifold structure on \overline{X} , as well as on any of the intermediate blow-ups leading all the way up to $X_{GV}:=[\overline{X}:\mathbb{S}_{\mathcal{F}}]$. At first, the Lie manifold structure on \overline{X} is defined by the action of X on itself and on \overline{X} as described in Example 4.19, namely, the structural Lie algebra of vector fields \mathcal{W}_{eucl} on \overline{X} is $\mathcal{W}_{eucl}:=\mathcal{C}^{\infty}(\overline{X})X$. We have already described in [6] and in then in the Introduction of this paper how to get from \overline{X} to X_{GV} by iterated blow-ups. On any of these intermediate blow-ups \widehat{X} Theorem 3.18 defines an action of X (as a Lie group) on \widehat{X} extending smoothly the translation action. Deriving this Lie group action yields a Lie algebra action of X on \widehat{X} , given by a linear map $E: X \to \Gamma(\widehat{X})$, which in fact provides smooth extensions of all constant vector fields on X to the compactification \widehat{X} . In general E(X) will not vanish at the boundary, but it will be tangent to the boundary, see below for further discussion.

On all these blow-ups leading to the Georgescu–Vasy space X_{GV} , the Euclidean metric will be compatible with the Lie manifold structure on the (intermediate) blow-ups, since we are performing only boundary blow-ups.

An alternative way to obtain the Lie manifold structure on all the iterated blowups \widehat{X} , including X_{GV} is to use Remark 4.23 to define the Lie manifold structure inductively, starting from the Lie manifold structure on \overline{X} . It is easy to check by induction over the iterated blow-ups that this construction yields the same Lie manifold structures as above.

We stress that, as a consequence of Remark 4.23, the resulting Sobolev spaces on all the blow-ups between \overline{X} and X_{GV} will be the same. Namely, they are the *usual Sobolev spaces* on X. Recall that we identify X with constant vector fields on itself. The algebra of differential operators, however, will change with each blow-up (along

some $\mathbb{S}_Y, Y \in \mathcal{F}$), eventually leading on $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$ to the structural Lie algebra of vector fields $\mathcal{W}_{bdry}:=\mathcal{C}^{\infty}([\overline{X} : \mathbb{S}_{\mathcal{F}}])X$. Therefore,

$$\operatorname{Diff}_{\mathcal{W}_{\operatorname{bdry}}}(X_{\operatorname{GV}}) = \mathcal{C}^{\infty}(X_{\operatorname{GV}})\mathbb{C}[X],$$

where $\mathbb{C}[X]$ denotes the algebra of polynomials on X and where a monomial of degree d is interpreted as a differential operator of degree d with constant coefficients.

5.1.3 Interior blow-ups

As we have seen already, the situation becomes more complicated once we start performing blow-ups along (lifts of) elements of $\overline{\mathcal{F}}$.

Remark 5.7 We use the notation introduced in Example 5.2 and Lemma 5.3. In particular, let $\widehat{\mathcal{F}}$ be the lifting of $\overline{\mathcal{F}}$ to X_{GV} . Let $\rho_{\widehat{\mathcal{F}}}$ be a smoothed distance function to $\widehat{\mathcal{F}}$ in $X_{\text{GV}}:=[\overline{X}:\mathbb{S}_{\mathcal{F}}]$. By iterating Remark 4.25, we obtain a Lie manifold structure on

$$X_{\mathcal{F}} := [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}] = [[\overline{X} : \mathbb{S}_{\mathcal{F}}] : \widehat{\mathcal{F}}] = [X_{\mathrm{GV}} : \widehat{\mathcal{F}}]$$

(see Corollary 5.4) by taking

$$\mathcal{W}_{\mathcal{F}} := \rho_{\widehat{\mathcal{F}}} \mathcal{C}^{\infty}(X_{\mathcal{F}}) \mathcal{W}_{\text{int}} = \rho_{\widehat{\mathcal{F}}} \mathcal{C}^{\infty}(X_{\mathcal{F}}) \mathcal{W}_{\text{eucl}} = \rho_{\widehat{\mathcal{F}}} \mathcal{C}^{\infty}(X_{\mathcal{F}}) X.$$

Let us record here what are the various quantities associated with the Lie manifold $(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}})$, again by iterating Remark 4.25 (starting from $M := X_{\text{GV}}$ and successively blowing up with respect to the lifts of $Y \in \mathcal{F}$):

• The interior smooth manifold of the blow-up is this time:

$$X_{\mathcal{F}} \smallsetminus \partial(X_{\mathcal{F}}) = X \smallsetminus \bigcup \mathcal{F}.$$

- If g is a W_{eucl}-compatible metric on X_{GV}, then ğ̃:=ρ⁻²_Fβ^{*}g is a W_F-compatible metric on X_F:=[X_{GV} : F̂]. Here β : X_F → X_{GV} is the associated blow-down map.
- The volume forms for g and \tilde{g} are related by the formula $dvol^{\tilde{g}} = \rho_{\hat{\mathcal{F}}}^{-n} \beta^* dvol^g$, $n = \dim(M) = \dim(X)$. As a consequence, we have

$$u \in L^{p}(X_{\mathcal{F}}; \mathcal{W}_{\mathcal{F}}) \iff \rho_{\widehat{\mathcal{F}}}^{-n/p} u \in L^{p}(X_{\mathrm{GV}}; \mathcal{W}_{\mathrm{int}})$$
$$\iff \rho_{\widehat{\mathcal{F}}}^{-n/p} u \in L^{p}(X; \operatorname{dvol}_{g}).$$

The Sobolev spaces defined by W_F are "weighted Sobolev spaces" in the old metric g (compare to Equation (19) and notice the factor ρ^j_τ):

$$W^{k,p}(X_{\mathcal{F}},\mathcal{W}_{\mathcal{F}})$$

🖄 Springer

$$= \left\{ u \mid \rho_{\widehat{\mathcal{F}}}^{j} v_{1} \dots v_{j} u \in L^{p}(X_{\mathcal{F}}; \mathcal{W}_{\mathcal{F}}), \forall v_{1}, \dots, v_{j} \in X, 0 \leq j \leq k \right\}$$
$$= \left\{ u \mid \rho_{\widehat{\mathcal{F}}}^{j} V_{1} \dots V_{j} u \in L^{p}(X_{\mathcal{F}}; \mathcal{W}_{\mathcal{F}}), \forall V_{1}, \dots, V_{j} \in \mathcal{W}_{\text{int}}, 0 \leq j \leq k \right\}.$$

• Diff_{\mathcal{W}_{\mathcal{F}}}^m(X_{\mathcal{F}}) is the linear span of differential monomials of the form $a\rho_{\widehat{\mathcal{F}}}^j V_1 V_2 \dots V_j$, with $a \in \mathcal{C}^{\infty}(X_{\mathcal{F}}), V_1, V_2, \dots, V_j \in \mathcal{W}_{\text{eucl}}, 0 \leq j \leq m$.

In order to obtain the last statement, we use, in particular, that $\rho_{\widehat{\mathcal{F}}}$ is the product of the smoothed distance functions to the blow-ups of \widehat{Y} , where $\widehat{Y} \in \widehat{\mathcal{F}}$, see Remark 4.12. It is important to keep in mind that $\rho_{\widehat{\mathcal{F}}}$ is a smooth distance functions in X_{GV} and that it is a product of smoothed distance functions in blow-ups of X_{GV} ; the statements would be different for smoothed distance functions in \overline{X} and the corresponding blow-ups thereof.

5.2 A lifting lemma and smoothed distance functions

We continue to let *X* be a finite-dimensional Euclidean vector space. Let *Y* be a linear subspace of *X*. In this subsection, we prove a technical lemma (a lifting lemma) and apply it to the study of smoothed distance functions, leading to some results that will be used in the next subsection. Recall that the projection map $\pi_{X/Y} : X \to X/Y$ extends canonically to a smooth map $\psi_Y : [\overline{X} : \mathbb{S}_Y] \to \overline{X/Y}$, see [6, Proposition 5.2] and the proof of [37, Theorem 4.1]. See also [42].

Lemma 5.8 There exists a diffeomorphism

$$\Psi_Y = (\psi_Y, \chi) : [\overline{X} : \mathbb{S}_Y] \to \overline{X/Y} \times \overline{Y}$$
(24)

such that $\chi(y) = y$ if $y \in Y$ and $\beta_{\overline{X}, \mathbb{S}_Y} \circ \Psi_Y^{-1}$ restricts on $\overline{X/Y} \times \mathbb{S}_Y$ to the projection $\overline{X/Y} \times \mathbb{S}_Y \to \mathbb{S}_Y$.

Proof This follows from [6, Lemma 4.14] for k = 1 and k' = 0, see Proposition C.2 in "Appendix C" for details.

Remark 5.9 In order to compare the above well-known lemma to the literature, we mention that it says that the map ψ_Y is the projection maps of a trivial fibration with fiber \overline{Y} . In this context, a smooth map $f : A \to B$ between manifolds with corners is the projection of a trivial fibration with fiber *C*, if and only if, *f* is the first component of a diffeomorphism $A \to B \times C$. This is a special case of a fibration in the sense of [42, Subsection 2.4], a concept whose definition will not be recalled here, as it will not be required further.

Continuing our idea to extend the map Ψ_Y to a case with more than one blow-up, let us introduce some notation. Let $Y \in \mathcal{F}$ and choose an admissible order ($P_0 = \emptyset, P_1, \ldots, P_k$) on $\mathbb{S}_{\mathcal{F}}$ (which is the same as choosing an admissible order on \mathcal{F}). According to the last part of Proposition 3.20, we can choose this admissible order such that $\mathbb{S}_Y = P_t$ for some $t \leq k$ and $\mathbb{S}_{P_j} \subset \mathbb{S}_Y$ for all j < t. Admissibility implies $\mathbb{S}_{P_j} \not\subset \mathbb{S}_Y$ for j > t. Let Ψ_Y be the diffeomorphism defined in Lemma 5.8. Let $\mathcal{P}:=(P_0, P_1, \ldots, P_{t-1}), \widehat{X}:=[\overline{X}: (\mathcal{P}, \mathbb{S}_Y)].$ **Corollary 5.10** Let \widehat{Y} be the lift of \overline{Y} to $\widehat{X} := [\overline{X} : (\mathcal{P}, \mathbb{S}_Y)]$. Using the notation we have just introduced, we have that the diffeomorphism $\Psi_Y : [\overline{X} : \mathbb{S}_Y] \to \overline{X/Y} \times \overline{Y}$ of Lemma 5.8 lifts to a diffeomorphism

$$\widehat{\Psi}_{Y}: \widehat{X} := [\overline{X}: (\mathcal{P}, \mathbb{S}_{Y})] \to \overline{X/Y} \times [\overline{Y}: \mathcal{P}].$$
⁽²⁵⁾

This diffeomorphism maps \widehat{Y} to $\{0_{X/Y}\} \times [\overline{Y} : \mathcal{P}]$ diffeomorphically. In particular, a smoothed distance function to $\{0_{X/Y}\}$ in $\overline{X/Y}$ pulls back via $\widehat{\Psi}_Y$ to a smoothed distance function to \widehat{Y} in \widehat{X} .

Proof Let $\beta := \beta_{\overline{X}, \mathbb{S}_Y}$ and $\beta^{-1}(\mathcal{P}) := (\beta^{-1}(P_0), \beta^{-1}(P_1), \dots, \beta^{-1}(P_{t-1}))$. (Notice that $\beta^{-1}(P_0) = \beta^{-1}(\emptyset) = \emptyset$.) Thanks to the properties of the family \mathcal{F} and the fact that \mathbb{S}_Y is a maximum of $\{P_0, P_1, \dots, P_t\}$, the two orders $(P_0, P_1, \dots, P_t = \mathbb{S}_Y)$ and $(P_0, \mathbb{S}_Y, P_1, \dots, P_{t-1})$ are both "intersection orders" in the sense of [37] for the semilattice $(P_i)_{i=1}^t$. Corollary 3.5 of [37] gives the first diffeomorphism of the following equation:

$$\begin{split} [\overline{X} : (\mathcal{P}, \mathbb{S}_Y)] &\simeq \left[[\overline{X} : \mathbb{S}_Y], \beta^{-1}(\mathcal{P}) \right] \\ &\simeq [\overline{X/Y} \times \overline{Y} : \Psi_Y(\beta^{-1}(\mathcal{P}))] \\ &\simeq [\overline{X/Y} \times \overline{Y} : \overline{X/Y} \times \mathcal{P}] \\ &\simeq \overline{X/Y} \times [\overline{Y} : \mathcal{P}]. \end{split}$$

The second diffeomorphism of this equation is obtained by using the diffeomorphism Ψ_Y of Lemma 5.8. Using $P_j \subset \mathbb{S}_Y$, Lemma 5.8 implies, $\Psi_Y(\beta^{-1}(P_j)) = \overline{X/Y} \times P_j$, and hence $\Psi_Y(\beta^{-1}(\mathcal{P})) = \overline{X/Y} \times \mathcal{P}$. This provides the third diffeomorphism. The last diffeomorphism follows then from Proposition 3.13.

We now turn to the second statement of our result, namely, that $\widehat{\Psi}_Y$ maps \widehat{Y} to $\{0_{X/Y}\} \times [Y:\mathcal{P}]$ diffeomorphically. By definition, the lift \widehat{Y} of \overline{Y} to \widehat{X} is the closure of $\overline{Y} \setminus \mathcal{P}$ in \widehat{Y} . In view of the diffeomorphism of the last equation and of Lemma 5.8, this lift is the closure of $\{0\} \times Y$ in $\overline{X/Y} \times [\overline{Y}:\mathcal{P}]$, and hence diffeomorphic to $[\overline{Y}:\mathcal{P}]$. (The diffeomorphism $\widehat{Y} \simeq [\overline{Y}:\mathcal{P}]$ also follows by iterating Proposition 3.3.)

The final statement about distances is then an immediate consequence of this diffeomorphism. (It is also a consequence of the more general result proved in Proposition 4.14.)

We will use Corollary 5.10 to define a canonical map $b : X_{\text{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \to \overline{X/Y}$. This map coincides with the composite map considered also in [6, Proposition 5.10]. Combining with Lemma 4.10, we obtain the following.

Corollary 5.11 Using the notation introduced in Corollary 5.10, let r_0 be a smoothed distance function to $0 \in \overline{X/Y}$. Let $\beta_{\text{GV}} : [\overline{X} : \mathbb{S}_{\mathcal{F}}] \to \overline{X}$ be the blow-down map. Let b be the composite map

$$X_{\mathrm{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \to [\overline{X} : (\mathcal{P}, \mathbb{S}_{Y})] \to \overline{X/Y}$$
(26)

obtained from Corollary 5.10. Then, $r_0 \circ b$ is a smoothed distance function to $\beta^*_{\text{GV}}(\overline{Y}) \subset [\overline{X} : \mathbb{S}_{\mathcal{F}}]$ in $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$.

Proof This follows from the properties of the map b and the aforementioned Lemma 4.10 and Corollary 5.10 as follows. Let \mathcal{P} and \widehat{Y} be as in Corollary 5.10. The remaining spheres at infinity yield $\mathcal{P}_{\text{rem}} := \beta_{\overline{X}, (\mathcal{P}, \mathbb{S}_Y)}(\mathbb{S}_{\mathcal{F}} \setminus \{\mathcal{P}, \mathbb{S}_Y\})$ which consists of p-submanifolds disjoint from \widehat{Y} . We first have that $r_0 \circ b$ is a distance function to \widehat{Y} in $\widehat{X} := [\overline{X} : (\mathcal{P}, \mathbb{S}_Y)]$. Then, this lifts to a distance function to $\beta^*(\overline{Y})$ in $X_{\text{GV}} := [[\overline{X} : (\mathcal{P}, \mathbb{S}_Y)] : \mathcal{P}_{\text{rem}}]$ by Lemma 4.10.

5.3 More on smooth distance functions on blow-ups

As before let $\beta_{\text{GV}} : X_{\text{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}] \to \overline{X}$ be the blow-down map from the Georgescu-Vasy compactification to the ball compactification. Again for $Y \in \mathcal{F}$, we write $\beta_{\text{GV}}^*(Y)$ for the lift of $\overline{Y} \subset \overline{X}$ to $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$, i.e., it is the closure of Y in $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$. In this subsection we use the notation $\widehat{Y} := \beta_{\text{GV}}^*(Y)$. (Note that now \widehat{Y} only has this specific meaning, whereas it denoted a more general class of lifts previously.)

The goal of the current subsection is to compare a smoothed distance to \widehat{Y} in X_{GV} to the Euclidean distance to Y. More precisely, we show in the following lemma that functions similar to $\operatorname{arctan}(d_Y) : X \to \mathbb{R}$, where d_Y is the Euclidean distance function to X, extend to a smoothed distance function to \widehat{Y} in X_{GV} . This is mostly a question about the asymptotic of d_Y at infinity.

The reader should be aware that a smoothed distance to \overline{Y} in \overline{X} will not pull back to a smoothed distance function to \widehat{Y} in X_{GV} ; thus, it is important to keep in mind with respect to which of the two compactifications, \overline{X} or \widehat{X} , the smoothed distance functions is defined. Again, we let $\widehat{\mathcal{F}} := \beta_{\text{GV}}^*(\overline{\mathcal{F}}) = \{\widehat{Y} \mid Y \in \mathcal{F}\}$ and recall from Corollary 5.4 that $[X_{\text{GV}} : \widehat{\mathcal{F}}] = [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}] =: X_{\mathcal{F}}$.

Lemma 5.12 Let $\phi_0 : [0, \infty) \to [0, 1]$ be a smooth, non-decreasing function satisfying $\phi_0(t) = t$ for $t \le 1/2$ and $\phi_0(t) = 1$ for $t \ge 1$. Let $Y \in \mathcal{F}$ and $d_Y(x) := \text{dist}(x, Y)$ be the distance function from $x \in X$ to Y with respect to the Euclidean distance on X, as before. Then, $\phi_0 \circ d_Y$ extends to a smoothed distance function $r_{\widehat{Y}}$ to \widehat{Y} in $X_{\text{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}]$ such that $r_{\widehat{Y}}/d_Y$ extends to a smooth function on X_{GV} .

It follows trivially from the lemma that $r_{\hat{Y}}/d_Y$ extends to a smooth function on $X_{\mathcal{F}}:=[X_{\text{GV}}:\widehat{\mathcal{F}}].$

Proof Let $r_0(z):=\phi_0(|z|) = \phi_0(\operatorname{dist}(z, 0))$, where $z \in X/Y$. Then, r_0 extends to a smooth distance function to 0 in $\overline{X/Y}$. Let *b* be the map of Corollary 5.11. Then, that corollary gives that $r_{\widehat{Y}} = r_0 \circ b$ is a smoothed distance function to \widehat{Y} in X_{GV} that coincides with $\phi_0 \circ d_Y$ on *X*.

Next, the function $q(z):=r_0(z)/|z| = \phi_0(|z|)/|z|$ extends to a smooth function on $\overline{X/Y}$. Hence, $q \circ b$ is smooth on X_{GV} . This function is the desired smooth extension of $r_{\widehat{Y}}/d_Y$.

Remark 5.13 Note that $d_Y|_X$ is a smoothed distance function to Y in X, but not in \overline{X} as it does not extend to a continuous (or smooth) function $\overline{X} \to [0, \infty)$.

The purpose of the framework developed in Sect. 4.1 was to prove the following result.

Proposition 5.14 Let $Y \in \mathcal{F}$ and $d_Y(x) := \operatorname{dist}(x, Y) = \inf\{||x - y|| | y \in Y\}$ be the Euclidean distance function from $x \in X$ to Y and $\widehat{\mathcal{F}}$ be the lift of $\overline{\mathcal{F}}$ to X_{GV} , as before. Let $\rho_{\widehat{\mathcal{F}}} : X_{\mathrm{GV}} \to [0, \infty)$ be a smoothed distance function to $\widehat{\mathcal{F}}$ in X_{GV} . Then, $\rho_{\widehat{\mathcal{F}}} d_Y^{-1} : X \setminus Y \to [0, \infty)$ extends smoothly to a function in $\mathcal{C}^{\infty}([X_{\mathrm{GV}} : \widehat{\mathcal{F}}]) = \mathcal{C}^{\infty}(X_{\mathcal{F}})$, denoted again by $\rho_{\widehat{\mathcal{F}}} d_Y^{-1}$.

Proof We use the notation introduced in Lemma 5.12, in particular, $r_{\widehat{Y}} := r_0 \circ b$ extends $\phi_0 \circ d_Y$. We thus have that $r_{\widehat{Y}}/d_Y \in \mathcal{C}^{\infty}(X_{\mathrm{GV}}) \subset \mathcal{C}^{\infty}(X_{\mathcal{F}})$, by Lemma 5.12. Moreover, $r_{\widehat{Y}}$ is a smoothed distance to \widehat{Y} in $X_{\mathrm{GV}} = [\overline{X} : \mathbb{S}_{\mathcal{F}}]$, by the same lemma. Hence, $\rho_{\widehat{\mathcal{F}}}/r_{\widehat{Y}}$ extends to a smooth function on $[[\overline{X} : \mathbb{S}_{\mathcal{F}}] : \widehat{\mathcal{F}}] = [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}] = X_{\mathcal{F}}$, by Proposition 4.15 for $M = [\overline{X} : \mathbb{S}_{\mathcal{F}}]$ and $\mathcal{S} = \widehat{\mathcal{F}}$. Hence,

$$\rho_{\widehat{\mathcal{F}}} d_Y^{-1} = \frac{\rho_{\widehat{\mathcal{F}}}}{r_{\widehat{Y}}} \cdot \frac{r_{\widehat{Y}}}{d_Y} \in \mathcal{C}^\infty(X_{\mathcal{F}}).$$

The proof is complete.

We fix in what follows $\rho_{\widehat{\mathcal{F}}}$ to be a smoothed distance to $\widehat{\mathcal{F}}$ in $[\overline{X} : \mathbb{S}_{\mathcal{F}}]$. Our constructions and reasoning will not depend on the particular choice of $\rho_{\widehat{\mathcal{F}}}$.

6 Regularity results for eigenfunctions

We now formulate and prove our main regularity results for certain differential operators with singular coefficients on X. These results apply, in particular, to Schrödinger operators with "inverse square potentials," a class of potentials which will be defined below and that includes the classical Schrödinger operator with Coulomb potential, which are used in physics and chemistry. The more general class of Schrödinger operators with inverse square potentials became of renewed interest [14, 16, 32, 33, 41]. To summarize our approach, we use the method in [1], but starting with X_{GV} instead of \overline{X} . This improvement leads to regularity statements which are even uniform close to infinity. The Lie manifold structure on X_{GV} is obtained from the action of X, as explained in Example 5.2. (See also Remarks 4.23 and 5.6.)

Recall that, throughout this paper, \mathcal{F} is a finite semilattice of linear subspaces of X with $0 \in \mathcal{F}, X \notin \mathcal{F}$, and $\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$ is the associated clean semilattice, as in Equation (4). There is no loss of generality to assume that $X = \mathbb{R}^n$, when convenient. As before, we let

- $X_{\text{GV}} := [X : \mathbb{S}_{\mathcal{F}}]$, the Georgescu–Vasy space;
- $X_{\mathcal{F}}:=[\overline{X}:\mathbb{S}_{\mathcal{F}}\cup\overline{\mathcal{F}}]=[[X:\mathbb{S}_{\mathcal{F}}]:\widehat{\mathcal{F}}];$ the blow-up along $\mathbb{S}_{\mathcal{F}}\cup\overline{\mathcal{F}};$
- $\rho_{\widehat{\mathcal{F}}}: X_{\text{GV}} \to [0, \infty)$, a smoothed distance to $\widehat{\mathcal{F}}$ in X_{GV} , see Definition 4.11;
- $\tilde{W}_{\mathcal{F}}:=\rho_{\hat{\mathcal{F}}}\mathcal{C}^{\infty}(X_{\mathcal{F}})X$, the structural Lie algebra of vector fields on $X_{\mathcal{F}}$, discussed in more detail in Remark 5.7.

Theorem 6.1 We use the usual notation, recalled for instance in the last paragraph. Let $m \in \mathbb{N}$ and D_0, D_1, \ldots, D_m be differential operators on X with coefficients in $\mathcal{C}^{\infty}(X_{\mathcal{F}}), D_k$ of order $\leq k$, and $D := D_m + \rho_{\widehat{\mathcal{F}}}^{-1} D_{m-1} + \ldots + \rho_{\widehat{\mathcal{F}}}^{-m} D_0$. Then,

- (i) $\rho_{\widehat{\tau}}^m D \in \text{Diff}_{\mathcal{W}_{\tau}}^m(X_{\mathcal{F}}).$
- (ii) If D_m is uniformly elliptic on X, then $\rho_{\widehat{T}}^m D$ is elliptic in $\text{Diff}_{W_{\mathcal{T}}}^m(X_{\mathcal{F}})$.
- (iii) For each boundary hyperface H of $X_{\mathcal{F}}$, let x_H be a defining function and $\mu_H \in \mathbb{R}$. Let $\chi := \prod_H x_H^{\mu_H}$ and $1 . We assume that <math>\rho_{\mathcal{F}}^m D$ is elliptic in $\operatorname{Diff}_{\mathcal{W}_{\mathcal{F}}}^m(X_{\mathcal{F}})$ and that $u \in \chi L^p(X_{\mathcal{F}})$ satisfies $Du = \lambda u$ on $X_{\mathcal{F}} \setminus \partial X_{\mathcal{F}} \subset X$ for some $\lambda \in \mathbb{R}$. Then, $u \in \chi W^{\ell, p}(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}})$ for all $\ell \in \mathbb{N}$.

Before proceeding to the proof, let us make a few comments on the setting.

Remark 6.2 All of X_{GV} , $X_{\mathcal{F}}$, $\rho_{\widehat{\mathcal{F}}}$, and $\mathcal{W}_{\mathcal{F}}$ are defined using an inductive procedure based on an ordering of $\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$. This ordering is not important, as long as it is an admissible order (see Definition 3.14). In our case, however, it is convenient to use an admissible order that puts first the elements of $\mathbb{S}_{\mathcal{F}}$ and then the elements of $\overline{\mathcal{F}}$, as many objects are defined on the intermediary blow-up $X_{\text{GV}} := [\overline{X} : \mathbb{S}_{\mathcal{F}}]$. We note also that $L^p(X_{\text{GV}}) = L^p(X)$ since

$$X_{\rm GV} \smallsetminus \partial X_{\rm GV} = \overline{X} \smallsetminus \partial \overline{X} = X \smallsetminus \bigcup \mathcal{F}, \tag{27}$$

with the same induced measure. However, the measure on the interior of $X_{\mathcal{F}}, X_{\mathcal{F}} \sim \partial X_{\mathcal{F}} = X \smallsetminus \bigcup \mathcal{F}$ differs from the measure on *X* by the factor $\rho_{\widehat{\mathcal{F}}}^{-n}$, where $n = \dim M$.

Also, we mention that this theorem generalizes Theorem 4.2 in [1], but note that, in the statement (ii) of that theorem, the radial compactification (denoted S in that paper) needs to be replaced with the Georgescu–Vasy space X_{GV} .

Proof

- Ad (i): This follows from Remarks 5.6 and 5.7 and from the definitions of $X_{\mathcal{F}}$ and $\mathcal{W}_{\mathcal{F}}$.
- Ad (ii): This follows from (i) just proved by combining with Remarks 4.25, 5.6, and 5.7.
- Ad (iii): Let $u \in \chi L^p(X_{\mathcal{F}})$ be such that $Du = \lambda u$. Then, $Q := \rho_{\widehat{\mathcal{F}}}^m D \lambda \rho_{\widehat{\mathcal{F}}}^m$ is an elliptic operator in Diff_{$\mathcal{W}_{\mathcal{F}}}(X_{\mathcal{F}})$, by (ii) just proved. It satisfies Qu = 0. The result is hence a direct consequence of the regularity result in [3] (this result was recalled in Theorem 4.22).}

Let us give now a more concrete application. We fix a Euclidean metric on X, $n = \dim(X)$. First, let us note that it follows from Remarks 4.25 and 5.6 that

$$W^{k,p}(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}}) := \{ u : X \to \mathbb{C} \mid \rho_{\widehat{\mathcal{F}}}^{|\alpha|} \partial^{\alpha} u \in L^{p}(X, \mu_{\mathcal{F}}), \ |\alpha| \le k \}$$
$$= \{ u : X \to \mathbb{C} \mid \rho_{\widehat{\mathcal{F}}}^{|\alpha| - (n/p)} \partial^{\alpha} u \in L^{p}(X, \mu_{\text{eucl}}), \ |\alpha| \le k \}$$
(28)

where we used the standard Lebesgue measure μ_{eucl} and the measure $\mu_{\mathcal{F}} = \rho_{\widehat{\mathcal{F}}}^{-n} \mu_{\text{eucl}}$ associated with $g_{\mathcal{F}} = \rho_{\widehat{\mathcal{F}}}^{-2} g_{\text{eucl}}$ on *X*.

🖄 Springer

Definition 6.3 Let $X, \mathcal{F}, \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}$, and $X_{\mathcal{F}} := [\overline{X} : \mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}]$ be as above (as in Theorem 1.1, for instance). Let $d_Y(x) := \operatorname{dist}(x, Y)$ denote the distance from $x \in X$ to $Y \in \mathcal{F}$. An *inverse square potential with singularities in* \mathcal{F} is a function $V : X \setminus \bigcup \mathcal{F} \to \mathbb{C}$ of the form

$$V(x) := \sum_{Y \in \mathcal{F}} \left(\frac{a_Y(x)}{d_Y(x)^2} + \frac{b_Y(x)}{d_Y(x)} \right) + c(x),$$

where $a_Y, b_Y, c \in \mathcal{C}^{\infty}(X_{\mathcal{F}})$.

Note that, since $C^{\infty}(\overline{X}) \subset C^{\infty}(X_{\mathcal{F}})$, our inverse square potentials are rather general and include the usual inverse square potentials. The following lemma justifies our definition of inverse square potentials.

Lemma 6.4 The set of inverse square potentials is a complex vector space. Let V be an inverse square potential, then $\rho_{\widehat{\tau}}^2 V \in \mathcal{C}^{\infty}(X_{\mathcal{F}})$.

Proof From the definition above, it is clear that the set of inverse square potentials is a complex vector space. Proposition 5.14 gives $\rho_{\widehat{\mathcal{F}}} d_Y \in \mathcal{C}^{\infty}(X_{\mathcal{F}})$. The result then follows since $\rho_{\widehat{\mathcal{F}}} \in \mathcal{C}^{\infty}(X_{\mathcal{F}})$.

Example 6.5 (The Schrödinger operator in quantum physics) The Schrödinger operator of an atom with heavy nucleus and with *N* electrons, studied in physics, is the operator

$$u(x) \mapsto (\mathcal{H}u)(x) := \Delta u(x) + V(x)u(x),$$

regarded as an unbounded, densely defined operator in $L^2(\mathbb{R}^n)$, n = 3N. We write $x = (x_1, x_2, \dots, x_N) \in \mathbb{R}^{3N}$ with $x_i \in \mathbb{R}^3$. The potential V is given by

$$V(x) := \sum_{1 \le j \le N} \frac{b_j}{\|x_j\|} + \sum_{1 \le i < j \le N} \frac{c_{ij}}{\|x_i - x_j\|}.$$

The potential V is an inverse square potential with singularities on the collision planes (more precisely, on the semilattice generated by the collision planes). We thus may apply Theorem 6.1 to eigenfunctions of the differential operator \mathcal{H} .

Moreover, the inverse square potentials considered in [14, 16, 32, 33, 41] are also inverse square potentials in our sense.

Theorem 6.6 We use the notation introduced in Theorem 6.1. Let D be a constant coefficient elliptic operator on $X = \mathbb{R}^n$ and V be an inverse square potential. Let $\rho_{\widehat{\mathcal{F}}}$ be a smoothed distance function to $\widehat{\mathcal{F}}$, as above. Assume $u \in L^2(\mathbb{R}^n)$ is an eigenfunction of D + V, then

$$\rho_{\widehat{\mathcal{F}}}^{|\alpha|} \partial^{\alpha} u \in L^2(\mathbb{R}^n)$$

for all multi-indices α .

🖉 Springer

Proof Let $g_{\mathbb{R}^n}$ be the canonical Euclidean metric on \mathbb{R}^n . It is also a compatible metric g_{GV} on X_{GV} , as explained in Remark 5.6. Then, $g_{X_{\mathcal{F}}} = \rho_{\widehat{\mathcal{F}}}^{-2} g_{\mathbb{R}^n}$ is a compatible metric $g_{\mathcal{F}}$ on $X_{\mathcal{F}}$, as explained in Remark 5.7. Hence, we have that

$$L^{2}(\mathbb{R}^{n},g_{\mathbb{R}^{n}}) = L^{2}(X_{\mathrm{GV}},g_{\mathrm{GV}}) = \rho_{\widehat{\mathcal{F}}}^{-n/2}L^{2}(X_{\mathcal{F}},g_{\mathcal{F}}).$$

The function $\rho_{\widehat{\mathcal{F}}}$ is a product of defining functions of hyperfaces of $X_{\mathcal{F}}$, by the definition of a smoothed distance function to a semilattice (Definition 4.11), and hence $\rho_{\widehat{\mathcal{F}}}^{-3N/2} = \chi$, for some χ as in Theorem 6.1 (iii) with $\mu_H \equiv -n/2$. The result then follows from Theorem 6.1 (iii) and the description of Sobolev spaces in Eq. (28).

Of course, $D = -\Delta$ satisfies the assumptions of the above theorem, so our regularity estimates are valid for Schrödiner operators with inverse square or Coulomb-type singularities. To obtain Theorem 1.1 stated in the Introduction, recall from Remark 4.16 the following. Let $\rho(x) := \text{dist}_{\overline{g}}(x, \bigcup \widehat{\mathcal{F}})$ be the distance to $\bigcup \widehat{\mathcal{F}}$ in some *true* metric \overline{g} on X_{GV} . Then, the functions $\rho_{\widehat{\mathcal{F}}}$, ρ , and the function $\delta_{\mathcal{F}} := \min\{\text{dist}(x, \bigcup \mathcal{F}), 1\}$ are Lipschitz equivalent, see "Appendix A."

Remark 6.7 The methods in this article also generalize to nonlinear equations. As an example, we consider equations of the form

$$\Delta u + V|u|^s u = \lambda u \tag{29}$$

on \mathbb{R}^n , where $\lambda \in \mathbb{C}$, $0 < s \le 2/(n-2)$, and *V* is of Coulomb type (i.e., as in (6)). Let $\hat{g} := \rho_{\hat{\mathcal{F}}}^{-2} g_{\text{eucl}}$ be an adapted Riemannian metric on $(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}})$. We define the *Yamabe operator* $L^g := \Delta^g - \frac{n-2}{4(n-1)} \operatorname{scal}^g$, and one may find in any reference about the Yamabe problem or conformal geometry that

$$L^{\hat{g}}(\hat{u}) = (\rho_{\hat{\mathcal{F}}})^{(n+2)/2} L^{\text{eucl}} ((\rho_{\hat{\mathcal{F}}})^{-(n-2)/2} \hat{u}).$$

After multiplication with $(\rho_{\widehat{\mathcal{F}}})^{(n+2)/2}$, Eq. (29) transforms into

$$L^{\hat{g}}\hat{u} + \rho_{\hat{\mathcal{F}}}^{t}V|\hat{u}|^{s}\hat{u} = \rho_{\hat{\mathcal{F}}}^{2}\lambda\hat{u}$$

with t = ((n+2) - (s+1)(n-2))/2 = 2 - (s+1)(n-2)/2 and $\hat{u} := \rho_{\hat{\mathcal{F}}}^{(n-2)/2} u$. We have $t \ge 1$ if and only if $s \le 2/(n-2)$, and in this case the Coulomb assumption implies the boundedness of $\rho_{\hat{\mathcal{F}}}^t V$. Then, regularity theory yields for any $\eta \in \mathcal{C}^{\infty}(X_{\mathcal{F}})$: if

$$\hat{u} \in \eta L^2(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}}) \cap \eta^{1/(s+1)} L^{2/(s+1)}(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}})$$
(30)

then

$$\hat{u} \in \eta W^{2,2}(X_{\mathcal{F}}, \mathcal{W}_{\mathcal{F}}).$$
(31)

🖄 Springer

We may apply this for example to $\eta := \rho_{\widehat{\mathcal{F}}}^{-(1+\frac{ns}{2})}$. Then,

$$u \in L^2(X, g_{\text{eucl}}) \cap L^{2/(s+1)}(X, g_{\text{eucl}})$$
(32)

implies (30), and then (31) implies

$$\rho_{\widehat{\mathcal{F}}}^{|\alpha|+\frac{ns}{2}}\partial^{\alpha}u \in L^2(X, g_{\text{eucl}}) \text{ for all multi-indices } \alpha \text{ with } |\alpha| \le 2.$$
(33)

Thus, we have $(32) \Rightarrow (33)$, and many similar conclusions are possible.

Remark 6.8 The point of view of Sect. 3.4 yields an alternative way to construct the space $X_{\mathcal{F}}$. Recall that as a set we have $X_{\infty} = X \cup \{\infty\}$ with the differential structure given by stereographic projection. Instead of taking the closure of a $\{0\} \neq Y \in \mathcal{F} \setminus \{0\}$ in \overline{X} , we may take its closure in X_{∞} which is then $Y_{\infty} = Y \cup \{\infty\}$. For $Y = \{0\}$, we use $\{0\}_{\infty} = \{0, \infty\}$ instead of the closure. We define

$$\mathcal{F}_{\infty} := \{ Y_{\infty} \mid Y \in \mathcal{F} \}.$$

Since we have (always) assumed that \mathcal{F} is a finite semilattice of linear subspaces of X with $\{0\} \in \mathcal{F}$, we see that \mathcal{F}_{∞} is a clean semilattice of closed p-submanifolds of X_{∞} (i.e., a cleanly intersecting family of closed p-submanifolds and a semilattice), which we endow with an admissible ordering. Because of the diffeomorphisms $[X_{\infty} : \{\infty\}] \simeq \overline{X}$ and $[Y_{\infty} : \{\infty\}] = \overline{Y} \subset [X_{\infty} : \{\infty\}]$, we see that $[X_{\infty} : \mathcal{F}_{\infty}] = [\overline{X}, \overline{\mathcal{F}}]$. This is the blow-up used in [1]. However, it differs from the blow-up constructed in this article, which is $X_{\mathcal{F}} = [X_{\text{GV}} : \overline{F}]$.

Example 6.9 Consider $\mathcal{F} = \{\{0\}, Y\}$ then $\mathcal{F}_{\infty} = \{\{0\} \cup \{\infty\}, Y_{\infty}\}$. Using the result that $[M : P, Q] \simeq [[M : P] : [Q : P]]$ for $P \subset Q$, we obtain

$$[X_{\infty}:\mathcal{F}_{\infty}] = [[X_{\infty}:\{\infty,0\}, [Y_{\infty}:\{\infty,0\}]] = [[X_{\infty}:\{\infty\}]:\{0\}, [Y_{\infty}:\{\infty\}]:\{0\}]$$
$$= [\overline{X}:\{0\}, \overline{Y}] = [\overline{X}:\overline{\mathcal{F}}] \neq X_{\mathcal{F}} = [\overline{X}:\mathbb{S}_{\mathcal{F}} \cup \overline{\mathcal{F}}].$$

Acknowledgements V.N. thanks Chris Kottke for some useful references. The authors thank Daniel Grieser for useful discussions about iterated blow-ups and lifts of submanifolds.

Funding Open Access funding enabled and organized by Projekt DEAL.

Data availability On behalf of all authors, the corresponding author (Bernd Ammann) states that there are no data associated with this article.

Declarations

Conflict of interest On behalf of all authors, the corresponding author (Bernd Ammann) states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Appendix A. The equivalence of ho and $\delta_{\mathcal{F}}$

Let *X* and \mathcal{F} be as in the previous sections, that is, \mathcal{F} is a finite semilattice of linear subspaces of the real, finite-dimensional vector space *X*. For each $Y \in \mathcal{F}$ let \widehat{Y} be the closure of *Y* in X_{GV} (it coincides with the lift $\beta^* \overline{Y}$ of \overline{Y} to X_{GV}) and $\widehat{\mathcal{F}} := \{\widehat{Y} \mid Y \in \mathcal{F}\}$. Recall from the Introduction that $\rho(x) := \text{dist}_{\overline{g}}(x, \widehat{\mathcal{F}}) = \text{dist}_{\overline{g}}(x, \mathcal{F})$ be the distance to $\widehat{\mathcal{F}}$ in some *true* metric \overline{g} on X_{GV} . Let $\text{dist}(x, \bigcup \mathcal{F})$ be the distance from *x* to $\bigcup \mathcal{F}$ in the usual, Euclidean distance and set $\delta_{\mathcal{F}}(x) := \min\{\text{dist}(x, \bigcup \mathcal{F}), 1\}$, again as in the Introduction. The function ρ used to obtain our regularity results in the previous section is maybe not explicit enough, so we prove now that the functions ρ and $\delta_{\mathcal{F}}$ are Lipschitz equivalent. More precisely, we shall prove the following result.

Proposition A.1 Let \mathcal{F} be a semilattice of linear subspaces of the real, finitedimensional vector space X. The following functions are Lipschitz equivalent (as functions on X):

- (1) $\rho :=$ the distance function to $\bigcup \widehat{\mathcal{F}}$ in some true metric on X_{GV} ;
- (2) $\delta_{\mathcal{F}}(x) := \min\{\operatorname{dist}(x, \bigcup \mathcal{F}), 1\}; and$
- (3) $\rho_{\mathcal{F}} := a$ smoothed *distance function to* $\bigcup \widehat{\mathcal{F}}$.

We recall that it was proved in [1] that $\rho_{\mathcal{F}}$ and ρ are continuously equivalent and thus Lipschitz equivalent. (This result was discussed and reminded in Remark 4.16.) The rest of this section will then be devoted to proving that ρ and $\delta_{\mathcal{F}}$ are Lipschitz equivalent.

Lemma A.2 Let (M, V) be a Lie manifold (and hence M is compact, see Definition 4.17), let \overline{g} be a true metric on M, and let g be a V-compatible metric. Then, there exists C > 0 such that $\overline{g} \leq C^2 g$. In particular, for all $x, y \in M \setminus \partial M$,

$$\operatorname{dist}_{\overline{g}}(x, y) \leq C \operatorname{dist}_{g}(x, y).$$

Proof As explained in Remark 4.21, a compatible metric on M is given by a metric $\gamma \in \Gamma(A^* \otimes A^*)$ on the vector bundle (Lie algebroid) A, and then $G := (\varrho \otimes \varrho)({}^{\sharp}\gamma^{\sharp}) \in \Gamma(TM \otimes TM)$ is a well-defined smooth symmetric section. On M_0 , the section G is non-degenerate, and thus we have $G|_{M_0} = {}^{\sharp}g_0{}^{\sharp}$ for some \mathcal{V} -compatible Riemannian metric g_0 on M_0 . By definition, any \mathcal{V} -compatible Riemannian metric arises in this way for some γ . We thus have seen that ${}^{\sharp}g{}^{\sharp}$ extends smoothly to a symmetric tensor $G \in \Gamma(TM \otimes TM)$. The \overline{g} -unit cotangent bundle $\mathbb{S}^*_{\overline{g}}M \subset T^*M$ is compact. By continuity of G and compactness of $\mathbb{S}^*_{\overline{\varrho}}M$ the supremum $c_1 := \sup\{G(X, X) \mid X \in \mathbb{S}^*_{\overline{\varrho}}M \in \mathbb{S}^*_{\overline{\varrho}}M$.

 $\mathbb{S}_{\overline{g}}^* M \subset T^*M$ is bounded. We get $G \leq c_1 \, {}^{\sharp}\overline{g}^{\sharp}$ and thus also ${}^{\sharp}g^{\sharp} \leq c_1 \, {}^{\sharp}\overline{g}^{\sharp}$. This implies the statement by duality for $C := \sqrt{c_1}$.

We obtain the following corollary.

Corollary A.3 Let g be the Euclidean distance on \mathbb{R}^n and \overline{g} be any true metric on X_{GV} . Then, there exists C > 0 such that, for any $x \in X$, we have

$$\operatorname{dist}_{\overline{g}}(x,\bigcup\mathcal{F}) \leq C\operatorname{dist}_g(x,\bigcup\mathcal{F})$$

Proof Let $y \in \bigcup \mathcal{F}$ be such that $\operatorname{dist}_g(x, \bigcup \mathcal{F}) = \operatorname{dist}_g(x, y)$, which exists since $\bigcup \mathcal{F}$ is a closed subset of *X*. Also, let *C* be as in Lemma A.2. Then,

$$C \operatorname{dist}_{\varrho}(x, \bigcup \mathcal{F}) = C \operatorname{dist}_{\varrho}(x, y) \ge \operatorname{dist}_{\overline{\varrho}}(x, y) \ge \operatorname{dist}_{\overline{\varrho}}(x, \bigcup \mathcal{F}).$$

Let $L:=\{x \in X \mid \operatorname{dist}_g(x, \bigcup \mathcal{F}) \ge 1\}.$

Corollary A.4 There is C > 0 such that, for all $x \in X$, $\rho(x) \leq C\delta_{\mathcal{F}}(x)$.

Proof The function ρ is bounded by some $C_1 > 0$ (since X_{GV} is compact). Hence, if $x \in L$, $\rho(x) \leq C_1 = C_1 \delta_{\mathcal{F}}(x)$. On the other hand, with *C* as in Corollary A.3, if $x \notin L := \{x \in X \mid \text{dist}_g(x, \bigcup \mathcal{F}) \geq 1\}$, we have $\rho(x) = \text{dist}_{\overline{g}}(x, \bigcup \mathcal{F}) \leq C \text{ dist}_g(x, \bigcup \mathcal{F}) = C \delta_{\mathcal{F}}(x)$. So the desired *C* is the largest of C_1 and the *C* of Corollary A.3.

Recall that in [6], one of the main results, namely Theorem 4.19, states that X_{GV} is diffeomorphic to the closure of $\delta(X)$, where δ is the diagonal map from X to $\prod_{Y \in \mathcal{F}} \overline{X/Y}$ (we assume that $0 \in \mathcal{F}$). The diffeomorphism is the unique extension of the diagonal map $x \mapsto (x, \ldots, x)$.

Let $(Z)_r$ resp. $(Z)_r$ be the open resp. closed ball of radius r around 0 in a Euclidean vector space Z.

Lemma A.5 Let Z be a finite-dimensional Euclidean space with open unit ball $(Z)_1$. There exists a true metric g_Z on \overline{Z} such that on $(Z)_1$ it yields the same distances as the Euclidean metric.

Proof Let us define $[0, \infty]$ as the closure of $[0, \infty)$ in \mathbb{R} . Then $[0, \infty]$ inherits a smooth structure of \mathbb{R} , and is a compact manifold with boundary, and there is a diffeomorphism $\rho : [0, \infty] \to [0, 2]$ with $\rho(t) = t$ for $t \le 1$. We obtain a diffeomorphism $\theta : \overline{Z} \to \overline{(Z)}_2$ defined by: $0 \mapsto 0, Z \setminus \{0\} \ni z \mapsto \frac{\rho(\|z\|)}{\|z\|} z$, and for $\|z\| = 1$: $\mathbb{S}_Z \ni \mathbb{R}_+ z \mapsto 2z$. Then, $g_Z := \theta^* g_{\text{eucl}}$ is a suitable true metric.

Let *g* be the Euclidean metric on $X = \mathbb{R}^n$, which, we recall, is a compatible metric on \overline{X} .

We are ready now to prove Proposition A.1.

Proof of Proposition A.1 We have proved in Corollary A.4 one of the two desired inequalities $(C\delta_{\mathcal{F}} \ge \rho)$ needed to prove that $\delta_{\mathcal{F}}$ and ρ are equivalent. Let us prove

now the opposite inequality. To that end, we can choose any true metric on X_{GV} (they are all equivalent). We shall choose then on each $\overline{X/Y}$ the true metric g_Y defined in Lemma A.5, and on $\prod_{Y \in \mathcal{F}} \overline{X/Y}$ we shall choose the product metric of these metrics. On X_{GV} we shall choose the induced Riemannian metric \overline{g} provided by the (diagonal) embedding $X_{GV} = \overline{\delta(X)} \subset \prod_{Y \in \mathcal{F}} \overline{X/Y}$ which is again a true metric. For this particular choice of true metric, we will show

$$\rho(x) \ge \delta_{\mathcal{F}}(x). \tag{34}$$

Note that the Riemannian distance from x to y in X_{GV} is bounded from below by the Riemannian distance of the same points x and y in $\prod_{Y \in \mathcal{F}} \overline{X/Y}$. This implies

$$\rho(x)^2 \ge \sum_{Y \in \mathcal{F}} \operatorname{dist}_{g_Y}(\pi_Y(x), 0)^2 \ge \operatorname{dist}_{g_Z}(\pi_Z(x), 0)^2$$

for any $Z \in \mathcal{F}$. We also have

$$\delta_{\mathcal{F}}(x) = \min\{1, \min_{Y \in \mathcal{F}} \operatorname{dist}(x, Y)\} \le \min\{1, \operatorname{dist}(x, Z)\}.$$

Thus, (34) will follow once we have shown

$$\operatorname{dist}_{g_Z}(\pi_Z(x), 0) \ge \min\{1, \operatorname{dist}(x, Z)\}$$
(35)

for any $Z \in \mathcal{F}$. Note that dist(x, Z) coincides with the Euclidean norm of $\pi_Z(x)$. As the distance with respect to g_Z coincides with the Euclidean distance inside the unit ball of $\overline{X/Z}$, see Lemma A.5 inequality (35) follows immediately. This completes the proof of (34) and thus of the proposition.

Appendix B. Group actions on compactifications of vector spaces

Here, we include the details on how the constant vector fields on a vector space extend to various compactifications. In order for the paper to be self-contained, we also recall and extend here some facts about several compactifications of a vector space X. We will start by considering the spherical compactification. We use the definitions of \mathbb{R}_k^n and \mathbb{S}_k^n as in (7).

Lemma B.1 Let X be a finite-dimensional real vector space. Let $A \in GL(X)$ and $V \in X$. Then, the affine map $p \mapsto L_{A,V}(p) := Ap + V$ extends to a diffeomorphism of \overline{X} .

Proof Let us assume first that $V = \mathbb{R}^n$. As a first, step we want to construct a smooth map $f_{A,V} : \mathbb{S}_1^n \to \mathbb{S}_1^n$ such that $\Theta_n^{-1} \circ f_{A,V} \circ \Theta_n$ is an extension of the affine map $L_{A,V}$, where Θ_n is the map $\overline{X} \to \mathbb{S}_1^n$ given by (8). As a second step, we prove that it is a diffeomorphism.

At first, we consider the linear map $F_{A,V} : \mathbb{R}^{n+1} \to \mathbb{R}^{n+1}$, $(t, p) \mapsto (t, Ap + tV)$. Obviously $F_{A,V}$ maps \mathbb{R}^{n+1}_1 to itself and $F_{A,V}((1, p)) = (1, L_{A,V}(p))$. It is easy to check that the smooth map

$$f_{A,V}((t, p)) := \frac{F_{A,V}((t, p))}{\|F_{A,V}((t, p))\|}$$

has the extension property for $L_{A,V}$ described above. It is obvious to see that $F_{A,V} \circ F_{A',V'} = F_{AA',V+AV'}$ and as a consequence $f_{A,V} \circ f_{A',V'} = f_{AA',V+AV'}$. It follows that $f_{A^{-1},-A^{-1}V}$ is the inverse to $f_{A,V}$, and hence $f_{A,V}$ is a diffeomorphism. The case of a general vector space X is obtained by choosing a linear isomorphism $X \simeq \mathbb{R}^n$. In view of the results proved, the resulting smooth structure on \overline{X} and the smoothness of the affine maps on \overline{X} do not depend on the choice of the isomorphism $X \simeq \mathbb{R}^n$. \Box

Obviously, the diffeomorphism $f_{A,V} : \overline{X} \to \overline{X}$ restricts to diffeomorphisms of the boundary, namely the sphere at infinity $f_{A,V}|_{\mathbb{S}_X} : \mathbb{S}_X \to \mathbb{S}_X$. From the proof of the last lemma, we also get:

Corollary B.2 The extension constructed in Lemma B.1 yields a group homomorphism $Aff(X) \rightarrow Diff(\overline{X}), L_{A,V} \mapsto f_{A,V}$. For a translation $L_{1,V} : X \rightarrow X$ by V, we have

$$f_{1,V}|_{\mathbb{S}_X} = id_{\mathbb{S}_X}.$$

Proof The proof is immediate from the formula defining the extensions of affine maps to the spherical compactification (see the proof of Lemma B.1). \Box

The map in the corollary defines a Lie group action of Aff(X) on \overline{X} , and the derivative of this Lie group action yields a Lie algebra action of aff(X), the Lie algebra of the affine group of X on \overline{X} . We may restrict to the Lie group of translations, resp. to the corresponding Lie algebra. This Lie algebra is X with the trivial bracket. We thus have a Lie algebra action of X on \overline{X} . If a Lie algebra \mathfrak{g} acts on a manifold with corners M, then this is given by a Lie algebra homomorphism $\mathcal{L} : \mathfrak{g} \to \Gamma(TM)$. In the above situation $X = \mathfrak{g}$ and \mathcal{L} maps a vector of X to the corresponding constant vector field in $\Gamma(X)$. As the Lie algebra X acts on \overline{X} , any constant vector field on X extends to a vector field on \overline{X} . As the Lie group of translations acts by the identity on \mathbb{S}_X , the Lie group X acts trivially on \mathbb{S}_X which proves that the smooth extension of a constant vector field to \overline{X} vanishes on \mathbb{S}_X . We thus have proven the following.

Corollary B.3 For any $V \in X$, the vector field

$$W(p) := \begin{cases} V & \text{if } p \in X \\ 0 & \text{if } p \in \mathbb{S}_X, \end{cases}$$

is a smooth vector field on \overline{X} .

We now turn to more involved compactifications. Let S be a finite, clean semilattice of closed submanifolds of S_X . We consider the compactification, $M := [\overline{X} : S]$, an example being the Georgescu–Vasy compactification X_{GV} discussed in 5.1.1.

The action of the Lie group X on \overline{X} constructed above is the identity on any $P \in S$, as $P \subset S_X$. Thus, [6, Theorem 4.21] or Theorem 3.18 tells us that the Lie group action of X on \overline{X} lifts (uniquely) to an action of X on M. The action is by diffeomorphisms, and, as X is connected, the action of an element of X on M maps each boundary face of M to itself. The corresponding Lie algebra action yields a Lie algebra homomorphism $\mathcal{L}: X = \text{Lie}(X) \rightarrow \Gamma(TM)$. Its image consists of vector fields tangent to the boundary of M as the corresponding Lie group action preserves all boundary faces. We have thus obtained the following result:

Lemma B.4 Let X and \mathcal{F} be as above and $M := [\overline{X} : \mathcal{F}]$. Then, any constant vector field on X extends to a smooth vector field on M that is tangent to the boundary.

We will now discuss whether the smooth extension of a constant vector field to $M := [\overline{X} : S]$ still vanishes on the boundary.

Example B.5 Let X and $S = \{\mathbb{S}_Y\}$, $Y \subset X$, $Y \neq \{0\}$. Then, we have seen that the constant vector fields $v \in X$ extend to a vector fields on \overline{X} vanishing at the boundary $\mathbb{S}_{\underline{X}}$. One can show that the lift of v to $[\overline{X} : \mathbb{S}_Y]$ vanishes at all the boundary faces of $[\overline{X} : \mathbb{S}_Y]$ if, and only if $v \in Y$. If $v \notin Y$, then its lift does not vanish on the boundary hyperface emerging from \mathbb{S}_Y , but it still vanishes on the other boundary hyperface of M (respectively, in the exceptional case dim $Y = \dim X - 1$, on the other two boundary hyperfaces of $[\overline{X} : \mathbb{S}_Y]$).

Appendix C. A splitting lemma

We now include the promised details explaining how Lemma 5.8 follows from [6, Lemma 4.14]. In particular, we consider the relations between (blow-ups of) the (spherical) compactification of the vector space *X* and the compactification of linear subspaces, providing, in particular, the full details of the proof of Lemma 5.8. So let *Y* be a linear subspace of *X*. We have shown in [6, Proposition 5.2] that the quotient map $X \to X/Y$ extends uniquely to a smooth map $\psi_Y : [\overline{X} : S_Y] \to \overline{X/Y}$. We will show that ψ_Y is a trivial fibration, a trivialization will be provided by the choice of any complement; in other words: $[\overline{X} : S_Y]$ is a product of manifolds with corners, and ψ_Y the projection to one of its factors.

In order to work on the space $[\overline{X} : S_X]$, it is convenient to have a better understanding of its structure, including of its smooth structure. The following result due to Melrose (see [6, Lemma 4.14] and the references therein) provide the needed initial results.

In the following lemma, we write elements $\mathbb{R}_{k+k'}^{m+m'+1}$ in the following way. At first, up to permutation of the coordinates $\mathbb{R}_{k+k'}^{m+m'+1}$ coincides with $\mathbb{R}_{k}^{m} \times \mathbb{R}_{k'}^{m'+1}$, which we write as $\mathbb{R}_{k+k'}^{m+m'+1} \cong \mathbb{R}_{k}^{m} \times \mathbb{R}_{k'}^{m'+1}$. Then, a vector in $\mathbb{R}_{k+k'}^{m+m'+1}$ will be written as a pair (η, μ) with respect to the product $\mathbb{R}_{k}^{m} \times \mathbb{R}_{k'}^{m'+1}$.

Lemma C.1 Let $\mathbb{S}_{k,k'}^{m,m'} := \mathbb{S}^{m+m'} \cap (\mathbb{R}_k^m \times \mathbb{R}_{k'}^{m'+1}) \cong \mathbb{S}_{k+k'}^{m+m'}$. If we define

$$\Psi: \mathbb{S}_{k,k'}^{m,m'} \smallsetminus \left(\{0\} \times \mathbb{S}_{k'}^{m'}\right) \to \mathbb{S}_{k}^{m-1} \times \mathbb{S}_{k'+1}^{m'+1}, \quad \Psi(\eta,\mu) = \left(\frac{\eta}{|\eta|}, \left(|\eta|,\mu\right)\right),$$

🖄 Springer

then Ψ extends to a diffeomorphism

$$\widetilde{\Psi} : [\mathbb{S}_{k,k'}^{m,m'} : \{0\} \times \mathbb{S}_{k'}^{m'}] \xrightarrow{\sim} \mathbb{S}_{k}^{m-1} \times \mathbb{S}_{k'+1}^{m'+1}$$
(36)

Up to a suitable permutation of coordinates, we permute the coordinates and if we identify \mathbb{S}_k^m with $\{0\} \times \mathbb{S}_k^m$ as subset of $\mathbb{S}_{k+k'}^{m+m'}$, we obtain a diffeomorphism:

$$\widetilde{\Psi}: [\mathbb{S}^{m+m'}_{k+k'}: \{0\} \times \mathbb{S}^{m'}_{k'}] \xrightarrow{\sim} \mathbb{S}^{m-1}_{k} \times \mathbb{S}^{m'+1}_{k'+1}$$

A proof of this result with our notations can be found in [6, Lemma 4.14] In the following proposition, we shall identify X/Y with Y^{\perp} , for some fixed scalar product $\langle \cdot, \cdot \rangle$ on X.

Proposition C.2 Let X be a real finite vector space and $Y \neq 0$ be a linear subspace of X. Then, there is a diffeomorphism

$$\Psi_Y = (\psi_Y, \chi) \colon [\overline{X} : \mathbb{S}_Y] \to \overline{X/Y} \times \overline{Y}$$

such that the restriction of ψ_Y to $\overline{X} \setminus \mathbb{S}_Y$ is the extension of the canonical projection $\pi_Y \colon X \to X/Y$ and $\chi(y) = y$ if $y \in Y$. Moreover, the blow-down map $\beta_{\overline{X},\mathbb{S}_Y}$ is determined by $\Xi_Y := \beta_{\overline{X},\mathbb{S}_Y} \circ \Psi_Y^{-1} \colon \overline{X/Y} \times \overline{Y} \to \overline{X}$ which is given by:

$$\Xi_Y(z, y) = \begin{cases} y & \text{if } y \in \mathbb{S}_Y, \\ z + \sqrt{1 + \langle z, z \rangle} y & \text{if } (z, y) \in X/Y \times Y. \end{cases}$$

Proof The proof is a modification of the proof of [6, Proposition 5.2]. The map ψ_Y is the same as the one defined in the aforementioned Proposition and used before in this paper. Let us recall briefly the construction of this map ψ_Y and then introduce the map χ . Let dim Y = p and dim X = n. Without loss of generality, we may assume that $Y = \mathbb{R}^p \subset X = \mathbb{R}^n$. Then, $\Theta_p(\mathbb{S}_Y) = \partial \mathbb{S}_1^p \cong \mathbb{S}^{p-1}$ may be viewed as subset of $\Theta_n(\mathbb{S}_X) = \partial \mathbb{S}_1^n \cong \mathbb{S}^{n-1}$. Using this identification, we can obtain

$$[\overline{X}:\mathbb{S}_Y]\simeq^{\Theta_n^\beta}[\mathbb{S}_1^n:\mathbb{S}_1^{p-1}],$$

where Θ_n^{β} is the lift of Θ_n to the two blow-up spaces. Using Lemma C.1 for m = n - p + 1 and m' = p, we have the diffeomorphism $[\mathbb{S}_1^n : \mathbb{S}^{p-1}] \simeq \mathbb{S}_1^{n-p} \times \mathbb{S}_1^p$. We go back to the compactified vector spaces using $(\Theta_{n-p}^{-1}, \Theta_p^{-1}) : \mathbb{S}_1^{n-p} \times \mathbb{S}_1^p \to \overline{X/Y} \times \overline{Y}$. Composing these diffeomorphisms, we obtain:

$$[\overline{X}:\mathbb{S}_Y]\simeq^{\Theta_n^\beta}[\mathbb{S}_1^n:\mathbb{S}^{p-1}]\simeq^{\Psi}\mathbb{S}_1^{n-p}\times\mathbb{S}_1^p\simeq^{(\Theta_{n-p}^{-1},\Theta_p^{-1})}\overline{X/Y}\times\overline{Y}.$$

At this point, the reader might want to check the definition of Θ_n and Θ_n^{-1} defined in (8) and (9), as well as Ψ defined in (36). If $v = (v_Y, v_\perp) \in Y \oplus Y^\perp$, we obtain

$$\widetilde{\Psi} \circ \Theta_n(v) = \Psi\left(\frac{1}{\langle v \rangle}(1,v)\right) = \left(\frac{1}{\langle v_\perp \rangle}(1,v_\perp), \frac{1}{\langle v \rangle}(\langle v_\perp \rangle, v_Y)\right).$$
(37)

For the first component, we obtain $\Theta_{n-p}^{-1}\left(\frac{1}{\langle v_{\perp}\rangle}(1, v_{\perp})\right) = v_{\perp} = \pi_Y(v)$. Hence, we have the required property for Ψ_Y . In the second component, if $v \in Y$, that is $v_{\perp} = 0$, we obtain $\langle v_{\perp} \rangle = 1$ and $\langle v \rangle = \langle v_Y \rangle$ then

$$\Theta_p^{-1}\left(\frac{1}{\langle v_Y \rangle}(1,v_Y)\right) = v_Y.$$

Hence, we have $\chi(y) = y$ for $y \in Y$. A long computation gives the required propriety for $\Xi_Y = \beta_{\overline{X}, \mathbb{S}_Y} \circ \Psi_Y^{-1}$.

Note that in contrast to the map ψ_Y , the constructions of the maps χ and Ξ_Y depend on more than the space *X* and its subspace *Y*, it also depends on the scalar product (and thus indirectly on the choice of the orthogonal complement). Thus, if $A : (X, Y) \rightarrow (X', Y')$ is an automorphism of pairs of vector space, and if we denote maps induced by *A* also by *A*, then we have the naturality relation $\pi_{Y'} \circ A = A \circ \pi_Y$, but in general χ and Ξ_Y are not natural, i.e., $\chi \circ A \neq A \circ \chi$ and $\Xi_{Y'} \circ A \neq A \circ \Xi_Y$.

References

- Ammann, B., Carvalho, C., Nistor, V.: Regularity for eigenfunctions of Schrödinger operators. Lett. Math. Phys. 101(1), 49–84 (2012)
- Ammann, B., Große, N., Nistor, V.: Well-posedness of the Laplacian on manifolds with boundary and bounded geometry. Math. Nachr. 292(6), 1213–1237 (2019)
- Ammann, B., Ionescu, A.D., Nistor, V.: Sobolev spaces on Lie manifolds and regularity for polyhedral domains. Doc. Math. 11, 161–206 (2006)
- Ammann, B., Lauter, R., Nistor, V.: On the geometry of Riemannian manifolds with a Lie structure at infinity. Int. J. Math. Math. Sci. 2004(1–4), 161–193 (2004)
- Ammann, B., Lauter, R., Nistor, V.: Pseudodifferential operators on manifolds with a Lie structure at infinity. Ann. Math. (2) 165(3), 717–747 (2007)
- Ammann, B., Mougel, J., Nistor, V.: A comparisons of the Georgescu and Vasy spaces associated to the *N*-body problems and applications. Ann. Henri Poincaré 23, 1141–1203 (2022). https://doi.org/ 10.1007/s00023-021-01109-1, arXiv:1910.10656, Shortlink https://rdcu.be/cyFzX
- Amrein, W., Boutet de Monvel, A., Georgescu, V.: C₀-groups, commutator methods and spectral theory of *N*-body Hamiltonians. Modern Birkhäuser Classics. Birkhäuser/Springer, Basel (1996). [2013] reprint of the 1996 edition
- Băcuță, C., Mazzucato, A.L., Nistor, V., Zikatanov, L.: Interface and mixed boundary value problems on *n*-dimensional polyhedral domains. Doc. Math. 15, 687–745 (2010)
- Boutet de Monvel-Berthier, A., Georgescu, V.: Graded C*-algebras and many-body perturbation theory. II. The Mourre estimate. méthodes semi-classiques, Vol. 2 (Nantes, 1991). Astérisque 210, 6–7, 75–96 (1992)
- Carvalho, C., Nistor, V., Qiao, Yu: Fredholm conditions on non-compact manifolds: theory and examples. In: Operator Theory, Operator Algebras, and Matrix Theory, Volume 267 of Operator Theory: Advances and Applications, pp. 79–122. Birkhäuser/Springer, Cham (2018)
- Castella, F., Jecko, T., Knauf, A.: Semiclassical resolvent estimates for Schrödinger operators with Coulomb singularities. Ann. Henri Poincaré 9(4), 775–815 (2008)

- Conlon, R., Degeratu, A., Rochon, A.: Quasi-asymptotically conical Calabi-Yau manifolds. Geom. Topol. 23, 29–100 (2019)
- Dereziński, J.: Asymptotic completeness of long-range N-body quantum systems. Ann. Math. (2) 138(2), 427–476 (1993)
- Dereziński, J., Faupin, J., Nguyen, Q., Richard, S.: On radial Schrödinger operators with a Coulomb potential: general boundary conditions. Adv. Oper. Theory 5(3), 1132–1192 (2020)
- Dereziński, J., Gérard, C.: Scattering Theory of Classical and Quantum N-Particle Systems. Texts and Monographs in Physics. Springer, Berlin (1997)
- Dereziński, J., Richard, S.: On Schrödinger operators with inverse square potentials on the half-line. Ann. Henri Poincaré 18(3), 869–928 (2017)
- 17. Felli, V., Ferrero, A., Terracini, S.: Asymptotic behavior of solutions to Schrödinger equations near an isolated singularity of the electromagnetic potential. J. Eur. Math. Soc. (JEMS) **13**(1), 119–174 (2011)
- Felli, V., Ferrero, A., Terracini, S.: On the behavior at collisions of solutions to Schrödinger equations with many-particle and cylindrical potentials. Discrete Contin. Dyn. Syst. 32(11), 3895–3956 (2012)
- Felli, V., Marchini, E., Terracini, S.: On Schrödinger operators with multisingular inverse-square anisotropic potentials. Indiana Univ. Math. J. 58(2), 617–676 (2009)
- Flad, H.-J., Hackbusch, W., Schneider, R.: Best *N*-term approximation in electronic structure calculations. II. Jastrow factors. M2AN Math. Model. Numer. Anal. 41(2), 261–279 (2007)
- Flad, H.-J., Harutyunyan, G., Schneider, R., Schulze, B.-W.: Explicit Green operators for quantum mechanical Hamiltonians. I. The hydrogen atom. Manuscr. Math. 135, 497–519 (2011)
- Flad, H.-J., Schneider, R., Schulze, B.-W.: Asymptotic regularity of solutions to Hartree–Fock equations with Coulomb potential. Math. Methods Appl. Sci. 31(18), 2172–2201 (2008)
- Fournais, S., Hoffmann-Ostenhof, M., Hoffmann-Ostenhof, T., Østergaard Sørensen, T.: Sharp regularity results for Coulombic many-electron wave functions. Commun. Math. Phys. 255(1), 183–227 (2005)
- Fournais, S., Hoffmann-Ostenhof, M., Hoffmann-Ostenhof, T., Østergaard Sørensen, T.: Analytic structure of many-body Coulombic wave functions. Commun. Math. Phys. 289(1), 291–310 (2009)
- Fournais, S., Hoffmann-Ostenhof, M., Hoffmann-Ostenhof, T., Østergaard Sørensen, T.: Analytic structure of solutions to multiconfiguration equations. J. Phys. A 42(31), 315208, 11 (2009)
- Georgescu, V.: On the essential spectrum of elliptic differential operators. J. Math. Anal. Appl. 468(2), 839–864 (2018)
- Georgescu, V., Iftimovici, A.: Localizations at infinity and essential spectrum of quantum Hamiltonians. I. General theory. Rev. Math. Phys. 18(4), 417–483 (2006)
- Grieser, D.: Basics of the *b*-calculus. In: Approaches to Singular Analysis (Berlin, 1999), Volume 125 of Operator Theory: Advances and Applications, pp. 30–84. Birkhäuser, Basel (2001). arXiv:math/0010314
- Grieser, D.: Scales, blow-up and quasimode constructions. In: Geometric and Computational Spectral Theory, Volume 700 of Contemporary Mathematics, pp. 207–266. American Mathematical Society, Providence (2017). arXiv:1607.04171
- Grieser, D., Talebi, M., Vertman, B.: Spectral geometry on manifolds with fibered boundary metrics I: low energy resolvent. J. Éc. polytech. Math. 9, 959–1019 (2022). arXiv:2009.10125
- Helffer, B., Siedentop, H.: Regularization of atomic Schrödinger operators with magnetic field. Math. Z. 218(3), 427–437 (1995)
- Hunsicker, E., Li, H., Nistor, V., Uski, V.: Analysis of Schrödinger operators with inverse square potentials I: regularity results in 3D. Bull. Math. Soc. Sci. Math. Roumanie (N.S.) 55(103)(2), 157– 178 (2012)
- Hunsicker, E., Li, H., Nistor, V., Uski, V.: Analysis of Schrödinger operators with inverse square potentials II: FEM and approximation of eigenfunctions in the periodic case. Numer. Methods Partial Differ. Equ. 30(4), 1130–1151 (2014)
- Jecko, T.: A new proof of the analyticity of the electonic density of molecules. Lett. Math. Phys. 93(1), 73–83 (2010). arXiv:0904.0221v8
- 35. Joyce, D.: A generalization of manifolds with corners. Adv. Math. 299, 760-862 (2016)
- Kottke, C.: Blow-up in manifolds with generalized corners. Int. Math. Res. Not. IMRN 2018(8), 2375–2415 (2018)
- Kottke, C.: Functorial compactification of linear spaces. Proc. Am. Math. Soc. 147(9), 4067–4081 (2019)

- Kottke, C., Melrose, R.: Generalized blow-up of corners and fiber products. In: Microlocal Methods in Mathematical Physics and Global Analysis, Trends Mathematics, pp. 59–62. Birkhäuser/Springer, Basel (2013)
- 39. Kottke, C., Rochon, F.: Quasi-fibered boundary pseudodifferential operators. arXiv:2103.16650 (2021)
- Laurent-Gengoux, C., Pichereau, A., Vanhaecke, P.: Poisson structures. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 347. Springer, Heidelberg (2013)
- Li, H., Nistor, V.: Analysis of a modified Schrödinger operator in 2D: regularity, index, and FEM. J. Comput. Appl. Math. 224(1), 320–338 (2009)
- 42. Melrose, R.: Differential analysis on manifolds with coners. Book in preparation. Manuscript available at http://www.math.mit.edu/~rbm/book.html
- Nistor, V.: Desingularization of Lie groupoids and pseudodifferential operators on singular spaces. Commun. Anal. Geom. 27(1), 161–209 (2019)
- Vasy, A.: Asymptotic behavior of generalized eigenfunctions in N-body scattering. J. Funct. Anal. 148(1), 170–184 (1997)
- 45. Vasy, A.: Structure of the resolvent for three-body potentials. Duke Math. J. 90(2), 379-434 (1997)
- Vasy, A.: Propagation of singularities in many-body scattering. Ann. Sci. École Norm. Sup. (4) 34(3), 313–402 (2001)
- Vasy, A.: Geometry and analysis in many-body scattering. In: Inside Out: Inverse Problems and Applications, Volume 47 of Mathematical Sciences Research Institute Publications, pp. 333–379. Cambridge University Press, Cambridge (2003)
- Yserentant, H.: On the regularity of the electronic Schrödinger equation in Hilbert spaces of mixed derivatives. Numer. Math. 98(4), 731–759 (2004)
- 49. Yserentant, H.: Regularity and Approximability of Electronic Wave Functions. Lecture Notes in Mathematics, vol. 2000. Springer, Berlin (2010)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.