Correction to: CYP2E1-mediated oxidative stress regulates HO-1 and GST expression in maneb- and paraquat-treated rat polymorphonuclear leukocytes

Israr Ahmad¹ · Smriti Shukla¹ · Deepali Singh^{1,2} · Amit Kumar Chauhan^{1,2} · Vinod Kumar^{1,2} · Brajesh Kumar Singh¹ · Devendra Kumar Patel¹ · Haushila Prasad Pandey³ · Chetna Singh^{1,2}

Published online: 19 July 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to:

Molecular and Cellular Biochemistry (2014) 393:209–222
<https://doi.org/10.1007/s11010-014-2062-y>

In the original publication of the article, the wrong β -actin blots were pasted in Figs. 1b and 2c. The correct versions of Figs. 1b and 2c are given in this correction.

The original article can be found online at <https://doi.org/10.1007/s11010-014-2062-y>.

✉ Chetna Singh
singhchetnaitrc@rediffmail.com

¹ Council of Scientific and Industrial Research-Indian Institute of Toxicology Research (CSIR-IITR), Mahatma Gandhi Marg, Post Box- 80, Lucknow, Uttar Pradesh 226 001, India

² Academy of Scientific and Innovative Research (AcSIR), New Delhi, India

³ Banaras Hindu University (BHU), Varanasi, Uttar Pradesh 221 005, India

Fig. 1 b Effect of MB and/or PQ on CYP2E1 and GSTA4-4 protein expression in rat PMNs following 1, 2 and 3 weeks of exposure. Upper panel shows representative western blots of CYP2E1, GSTA4-4 and β -actin protein expression and lower panel shows densitometric analysis of the same. ($n = 4$) (* $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ as compared with control; # $p < 0.05$, ## $p < 0.01$ and ### $p < 0.001$ as compared with MB treated groups; § $p < 0.05$ as compared with PQ treated groups). (Control, maneb-treated, paraquat-treated and maneb+paraquat treated groups are represented as Con/C, MB/M, PQ/P and MB+PQ/MP respectively)

Fig. 2 c Effect of DAS and AG pre-treatment on MB and/or PQ-induced increase in CYP2E1/GSTA4-4 protein expression following 2 weeks of exposure. Upper panel shows representative western blot of CYP2E1/GSTA4-4/ β -actin and lower panel shows densitometric analysis of the same. ($n = 4$) (** $p < 0.001$ as compared with control; # $p < 0.01$ and ### $p < 0.001$ as compared with MB treated

groups; \$\$\$ $p < 0.001$ as compared with PQ treated groups and &&& $p < 0.001$ as compared with MB+PQ treated groups). (Control, maneb-treated, paraquat-treated and maneb+paraquat treated groups are represented as Con/C, MB/M, PQ/P and MB+PQ/MP respectively)