Correction to: TAK1 is involved in sodium L-lactate-stimulated p38 signaling and promotes apoptosis

Qingen Da^{1,2} · Zilong Yan¹ · Zhangfu Li¹ · Zhen Han² · Mingming Ren² · Lei Huang² · Xiaowei Zhang³ · Jikui Liu¹ · Tao Wang²

Published online: 22 May 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to:

Molecular and Cellular Biochemistry (2021) 476:873–882
<https://doi.org/10.1007/s11010-020-03952-y>

In the original publication of the article, Fig. 2A was published incorrectly. The correct version of Fig. 2 is provided in this correction.

The original article can be found online at <https://doi.org/10.1007/s11010-020-03952-y>.

✉ Jikui Liu
Liu8929@126.com

✉ Tao Wang
szwangtao@126.com

¹ Department of Hepatobiliary Surgery, Peking University Shenzhen Hospital, Shenzhen Peking University, The Hong Kong University of Science and Technology Medical Center, Shenzhen, China

² Department of Cardiovascular Surgery, Peking University Shenzhen Hospital, Shenzhen Peking University, The Hong Kong University of Science and Technology Medical Center, Shenzhen 518000, China

³ School of Basic Medical Sciences, Peking University, Beijing, China

Fig. 2 TAK1 inhibitor TAK1 reduced L-lactate-triggered phosphorylation of p38. **A** HeLa cells were treated by vehicle, 25 μ M SB203580, 20 μ M Asiatic acid, 15 μ M NQDI-1, and 15 μ M TAK1 with and without 20 mM L-lactate and lysed. Western blot analyses of the protein and phosphorylated protein expression were performed. **B** and **C** The protein bands were digitized by ImageJ and plotted to present the protein level of p-p38 (**B**) and p-TAK1 (**C**). The protein bands were calculated as a ratio to total protein (TAK1 and p38) as an internal control. Quantitative data are shown as the mean $\pm$ SD of three independent experiments. * $p < 0.05$, ** $p < 0.01$ versus vehicle cells

12.25