



## Correction to: Combined therapeutic efficacy of carvacrol and X-radiation against 1,2-dimethyl hydrazine-induced experimental rat colon carcinogenesis

Sivaranjani Arivalagan<sup>1</sup> · Nisha Susan Thomas<sup>1</sup> · Balaji Chandrasekaran<sup>1</sup> · Vijay Mani<sup>1</sup> · Aktarul Islam Siddique<sup>1</sup> · Thayalan Kuppsamy<sup>2</sup> · Nalini Namasivayam<sup>1</sup>

Published online: 30 November 2021  
© Springer Science+Business Media, LLC, part of Springer Nature 2021

**Correction to:**  
**Molecular and Cellular Biochemistry (2015) 410:37–54**  
<https://doi.org/10.1007/s11010-015-2536-6>

In the original publication of the article, Fig. 2b was published incorrectly. The correct version of Fig. 2 is provided in this correction.

---


The original article can be found online at <https://doi.org/10.1007/s11010-015-2536-6>.

---

✉ Nalini Namasivayam  
nalininam@yahoo.com

<sup>1</sup> Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalai Nagar, Tamil Nadu 608 002, India

<sup>2</sup> Medical Physics Division, Dr. Kamakshi Memorial Hospital, No: 1, Radial Road, Pallikaranai, Chennai, Tamil Nadu 600 100, India



**Fig. 2** **a** Topographical view of normal crypts and ACF in the colonic mucosa of experimental rats ( $\times 10$ ). **A, B** Colon of control and control+CVC-treated rats show normal crypts. **C** Topographical view of ACF (arrows) with multiple (eighteen) crypts in the colon of a rat treated with DMH. **D** Topographical view of ACF (arrow) with eight crypts in the colon of a rat treated with DMH+X-radiation. **E** Topographical view of ACF (arrow) with four crypts in the colon of a rat treated with DMH+CVC. **F** Topographical view of ACF (arrow) with two crypts in the colon of a rat treated with DMH+CVC+X-radiation. **b** Topographical view of dysplastic ACF in the colonic mucosa of experimental animals ( $\times 20$ ). **A**: Normal

crypts in the unsectioned colon of control rats. **B** Normal crypts in the unsectioned colon of the rats supplemented with CVC alone. **C** DACF (arrow) with thirteen crypts resist methanol, decolouration after methylene blue staining in the unsectioned colon of the rat treated with DMH alone. **D** DACF (arrow) with seven crypts in the unsectioned colon of DMH-treated rat exposed to radiation alone. **E** DACF (arrow) with four crypts in the unsectioned colon of DMH-treated rat supplemented with CVC. **F** DACF (arrow) with three crypts in the unsectioned colon of DMH-treated rat treated with both CVC and X-radiation

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.