Correction to: Synergistic effects of curcumin with emodin against the proliferation and invasion of breast cancer cells through upregulation of miR-34a

Jiaoli Guo^{1,2} · Wenping Li³ · Hongliu Shi² · Xinhua Xie^{1,2} · Laisheng Li⁴ · Hailin Tang^{1,2} · Minqing Wu^{1,2} · Yanan Kong^{1,2} · Lu Yang^{1,2} · Jie Gao^{1,2} · Peng Liu¹ · Weidong Wei¹ · Xiaoming Xie^{1,2}

Published online: 15 December 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Mol Cell Biochem (2013) 382:103–111
<https://doi.org/10.1007/s11010-013-1723-6>

In the original publication of the article, Fig. 3 was published incorrectly. The correct version of Fig. 3 is provided in this correction.

The original article can be found online at <https://doi.org/10.1007/s11010-013-1723-6>.

✉ Weidong Wei
weiwd@sysucc.org.cn

✉ Xiaoming Xie
xiexm@sysucc.org.cn

¹ Department of Breast Oncology, Sun Yat-Sen University Cancer Center, 651 East Dong Feng Road, Guangzhou 510060, People's Republic of China

² State Key Laboratory of Oncology in South China, Sun Yat-Sen University Cancer Center, Guangzhou 510060, People's Republic of China

³ Department of Breast Disease, Maternal and Children Metabolic-Genetic Disease Key Laboratory of Guangdong Women and Children Hospital, Guangzhou 510623, People's Republic of China

⁴ Department of Laboratory Medicine, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510080, People's Republic of China

Fig. 3 Curcumin and emodin suppress breast cancer cell invasion. **a** Representative microscopic fields of invasive cells on the bottom of Matrigel-coated transwell cell culture chambers. MDA-MB-231 cells were treated with 30 μ M curcumin, 80 μ M emodin, or both; MDA-MB-435 cells were treated with 34 μ M curcumin, 70 μ M emodin, or both for 24 h. **b** Quantification of cell invasion was performed

by counting the number of invasive cells. Cells pretreated with curcumin, emodin, or both were seeded onto Matrigel-coated transwell plates and cultured for 24 h. After removing the non-migratory cells, the migratory cells on the bottom side of the membrane were counted. The results presented are the average of three experiments each expressed as the mean $\pm$ SD. * P < 0.05

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.