



## Correction to: *Celastrus paniculatus* oil ameliorates NF-KB mediated neuroinflammation and synaptic plasticity in the scopolamine-induced cognitive impairment rat model

Khushboo Govind Faldu<sup>1</sup> · Snehal Sanjay Patel<sup>2</sup> · Jigna Samir Shah<sup>1</sup>

Published online: 24 April 2023

© Springer Science+Business Media, LLC, part of Springer Nature 2023

### Correction to: Metabolic Brain Disease

<https://doi.org/10.1007/s11011-023-01186-7>

Images 7(C) and 7(E) were incorrectly inserted into the original Fig. 7. The corrected figure is shown below. The authors sincerely apologize for this mistake.

---


The online version of the original article can be found at <https://doi.org/10.1007/s11011-023-01186-7>.

---

✉ Jigna Samir Shah  
[jigna.shah@nirmauni.ac.in](mailto:jigna.shah@nirmauni.ac.in)

<sup>1</sup> Department of Pharmacology, Institute of Pharmacy, Nirma University, Sarkhej - Gandhinagar Hwy, Gota, Ahmedabad 382481, Gujarat, India

<sup>2</sup> Institute of Pharmacy, Nirma University, Ahmedabad, Gujarat, India



**Fig. 7** Histopathological examination of hippocampus region of the rat brain Hippocampus of 1 rat brain per group was utilized for the histopathological examination.

Figure 7 A: Normal Control, Fig. 7B: Disease Control Animals treated with Scopolamine (2 mg/kg IP). Figure 7 C: Disease treated with

Donepezil (1 mg/kg) from Day 8, Fig. 7D: Disease treated with *Celastrus paniculatus* oil (2 g/kg) from Day 0 and Fig. 7E: Disease treated with *Celastrus paniculatus* oil (2 g/kg) from Day 8. Magnification: 40x.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.