

## Bond dissociation enthalpies calculated by the PM3 method confirm activity cliffs in radical scavenging of flavonoids

Dragan Amić · Bono Lučić · Goran Kovačević ·  
Nenad Trinajstić

Published online: 27 November 2008  
© Springer Science+Business Media B.V. 2008

**Erratum to: Mol Divers**  
**DOI 10.1007/s11030-008-9095-7**

The term “ $I_{3',4'\text{-diOH or 3-OH}}$ ” was misrepresented throughout the original article. In addition to that error in Eq. 4, the minus sign was missing before the term “ $0.251(\pm 0.241)\text{BDE}_{\text{min}}$ ” in that equation. The correct equation is shown below.

$$\text{RSA} = 85.27 (\pm 71.66) + 66.77 (\pm 8.97)$$

$$I_{3',4'\text{-diOH or 3-OH}} - 0.251 (\pm 0.241) \text{BDE}_{\text{min}}$$

---

The online version of the original article can be found under  
doi:[10.1007/s11030-008-9095-7](https://doi.org/10.1007/s11030-008-9095-7).

---

D. Amić (✉)  
Faculty of Agriculture, The Josip Juraj Strossmayer University,  
P.O. Box 719, 31107 Osijek, Croatia  
e-mail: [damic@pfos.hr](mailto:damic@pfos.hr)

B. Lučić · G. Kovačević · N. Trinajstić  
The Rugjer Bošković Institute, P.O. Box 180, 10002 Zagreb,  
Croatia