

Erratum to: Genetic analysis reveals a novel locus in *Vicia faba* decoupling pigmentation in the flower from that in the extra-floral nectaries

Hamid Khazaei · Donal M. O’Sullivan ·
Mikko J. Sillanpää · Frederick L. Stoddard

Published online: 24 June 2014
© Springer Science+Business Media Dordrecht 2014

Erratum to: Mol Breeding
DOI 10.1007/s11032-014-0100-9

incorrectly with the headings at the top. Please find below the corrected version of Table 1.

In the original publication of the article, the lower columns in Table 1 (cross D and D’) aligned

The online version of the original article can be found under doi:[10.1007/s11032-014-0100-9](https://doi.org/10.1007/s11032-014-0100-9).

H. Khazaei (✉) · F. L. Stoddard
Department of Agricultural Sciences, University
of Helsinki, P O Box 27, Latokartanonkaari 5,
00014 Helsinki, Finland
e-mail: hamid.khazaei@helsinki.fi

D. M. O’Sullivan
School of Agriculture, Policy and Development,
University of Reading, Whiteknights,
P O Box 237, Reading, UK

M. J. Sillanpää
Department of Mathematical Sciences, University
of Oulu, P O Box 3000, 90014 Oulu, Finland

M. J. Sillanpää
Department of Biology and Biocenter Oulu, University
of Oulu, P O Box 3000, 90014 Oulu, Finland

Table 1 Observed offspring segregation ratios, value of χ^2 test and corresponding P value for coloured and colourless stipule spot pigmentation among F_1 – F_3 generations as well as number of independent genes segregating and segregating gene name in each population

Cross	F_1^a	F_2 populations		F_3 populations		$\chi^2_{(3:1)}$		$\chi^2_{(5:3)}$		Number of independent genes segregating	Segregating gene name
		Coloured	Colourless	Coloured	Colourless	Coloured	Colourless	P	P		
(A) Mélodie/2 × ILB 938/2 ^b	Coloured	261	89	208	142	0.034	0.853	1.409	0.235	1	<i>ssp1</i>
(A') ILB 938/2 × Mélodie/2 ^c	Coloured	263	88	205	144	0.001	0.975	2.106	0.147	1	<i>ssp1</i>
(B) Mélodie/2 × Disco/2	Coloured	109	41	–	–	0.436	0.509	–	–	1	<i>ssp2 (zt2)</i>
(B') Disco/2 × Mélodie/2	Coloured	107	43	–	–	1.076	0.300	–	–	1	<i>ssp2 (zt2)</i>
(C) ILB 938/2 × Aurora/2	Coloured	271	69	210	138	4.016	0.045	0.690	0.406	1	<i>ssp1</i>
(C') Aurora/2 × ILB 938/2	Coloured	259	91	–	–	0.187	0.666	–	–	1	<i>ssp1</i>
Pooled data		1,270	421	623	424	0.001	0.922	4.012	0.045		
$\chi^2_{(25:39)}$											
(D) ILB 938/2 × Disco/2	Coloured	163	130	120	169	0.046	0.831	0.735	0.391	2	<i>ssp1 + ssp2 (zt2)</i>
(D') Disco/2 × ILB 938/2	Coloured	163	128	110	179	0.007	0.935	0.121	0.728	2	<i>ssp1 + ssp2 (zt2)</i>
Pooled data		326	258	230	348	0.043	0.835	0.129	0.719		

^a Observations are from 25 to 30 F_1 plants

^b F_4 (coloured 151:colourless 129, $\chi^2_{(9:7)}$: 0.613, $P = 0.434$); F_5 (coloured 109:colourless 102, $\chi^2_{(17:15)}$: 0.182, $P = 0.670$); F_6 (coloured 102:colourless 101, $\chi^2_{(1:1)}$: 0.005, $P = 0.944$)

^c F_4 (coloured 157: colourless 134, $\chi^2_{(9:7)}$: 0.624, $P = 0.429$); F_5 (coloured 161:colourless 144, $\chi^2_{(17:15)}$: 0.014, $P = 0.906$)