Correction to: CEPO (carbamylated erythropoietin)-Fc protects hippocampal cells in culture against beta amyloid-induced apoptosis: considering Akt/GSK-3 β and ERK signaling pathways

Etrat Hooshmandi^{1,2} · Maryam Moosavi³ · Hermann Katinger⁴ · Shima Sardab⁵ · Rasoul Ghasemi⁶ · Nader Maghsoudi^{1,7}

Published online: 20 May 2020
© Springer Nature B.V. 2020

Correction to: Molecular Biology Reports
<https://doi.org/10.1007/s11033-020-05309-6>

Unfortunately, the original version of this article contained a mistake in the arrangement of representative cell images in Fig. 2. In this figure, the same representative image for A β group was mistakenly placed for A β + LY group. The corrected form of Fig. 2 is provided in this correction.

The original article can be found online at <https://doi.org/10.1007/s11033-020-05309-6>.

✉ Rasoul Ghasemi
rghasemi60@sbmu.ac.ir

✉ Nader Maghsoudi
Nmaghsoudi@sbmu.ac.ir

¹ Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

² Clinical Neurology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

³ Nanobiology and Nanomedicine Research Centre, Shiraz University of Medical Sciences, Shiraz, Iran

⁴ Department of Biotechnology, University of Natural Resources and Life Sciences, Vienna, Austria

⁵ Biology Department, University of Tehran, Tehran, Iran

⁶ Neurophysiology Research Center and Physiology Department, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁷ Neurobiology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Fig. 2 The representative images of morphological changes of hippocampal cells in different groups 24 h after cell treatments

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.