Correction to: Inhibition of semaphorin 4D enhances chemosensitivity by increasing 5-fluorouracil-induced apoptosis in colorectal cancer cells

Golnaz Rashidi¹ · Mahsa Rezaeepoor¹ · Chiman Mohammadi² · Ghasem Solgi¹ · Rezvan Najafi^{2,3}

Published online: 6 January 2021
© Springer Nature B.V. 2021

Correction to:
Molecular Biology Reports (2020) 47:7017–7027
<https://doi.org/10.1007/s11033-020-05761-4>

The author(s) wish to point out that an error appeared in Fig. 5 of the original published article. This error does not affect the paper's conclusion and the correct Fig. 5 is shown below.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Golnaz Rashidi and Mahsa Rezaeepoor have contributed equally to this work.

The original article can be found online at <https://doi.org/10.1007/s11033-020-05761-4>.

✉ Rezvan Najafi
najafi2535@gmail.com; re.najafi@umsha.ac.ir

¹ Department of Immunology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

² Department of Molecular Medicine and Genetics, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

³ Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

Fig. 5 Effect of SEMA4D knockdown on sensitivity of 5-FU-treated SW48 cell to cell invasion. Following treatment of SW48 cells with 5-FU in the presence and absence of siRNA SEMA4D, transwell invasion assay was performed. The data are expressed as the

mean $\pm$ standard deviation and each experiment was repeated three times. ** $P < 0.01$ and *** $P < 0.001$ vs. control group, $^{\dagger\dagger\dagger}P < 0.001$ vs. cells treated with 5-FU alone