



# Melatonin and REGN-CoV2 combination as a vaccine adjuvant for Omicron variant of SARS-CoV-2

Ismail Celil Haskologlu<sup>1</sup> · Emine Erdag<sup>2</sup> · Serkan Sayiner<sup>3,4</sup> · Nurettin Abacioglu<sup>1</sup> · Ahmet Ozer Sehirli<sup>5</sup>

Received: 20 February 2022 / Accepted: 23 March 2022 / Published online: 7 April 2022  
© The Author(s), under exclusive licence to Springer Nature B.V. 2022

## Abstract

The omicron variant (B.529) of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which emerged in late 2021, caused panic worldwide due to its contagiousness and multiple mutations in the spike protein compared to the Delta variant (B.617.2). There is currently no specific antiviral available to treat Coronavirus disease 2019 (COVID-19). However, studies on neutralizing monoclonal antibodies (mAb) developed to fight COVID-19 are growing and gaining traction. REGN-COV2 (Regeneron or imdevimab-casirivimab combination), which has been shown in recent studies to be less affected by Omicron's RBD (receptor binding domain) mutations among other mAb cocktails, plays an important role in adjuvant therapy against COVID-19. On the other hand, it is known that melatonin, which has antioxidant and immunomodulatory effects, can prevent a possible cytokine storm, and other severe symptoms that may develop in the event of viral invasion. Along with all these findings, we believe it is crucial to evaluate the use of melatonin with REGN-COV2, a cocktail of mAbs, as an adjuvant in the treatment and prevention of COVID-19, particularly in immunocompromised and elderly patients.

**Keywords** COVID-19 · Omicron variant · Melatonin · REGN-COV2 · Cytokine storm

## Abbreviations

ACE2 Angiotensin converting enzyme 2  
AI Artificial intelligence  
BMAL1 Brain and muscle ARNT-like protein 1

CD147 Cluster of differentiation 147  
COVID-19 Coronavirus disease 2019  
COX-2 Cyclooxygenase-2  
FDA Food and drug administration  
IL Interleukins  
IP10 Interferon- $\gamma$ -inducible protein 10  
iNOS Inducible nitric oxide synthase  
mAb Monoclonal antibodies  
MMPs Matrix metalloproteinases  
MPO Myeloperoxidase  
RAAS Renin–angiotensin–aldosterone system  
RBD Receptor binding domain  
REGN-COV2 Regeneron or imdevimab-casirivimab  
ROS Reactive oxygen species  
SARS-CoV-2 Severe acute respiratory syndrome coronavirus 2  
TGF- $\beta$  Transforming growth factor beta  
TNF- $\alpha$  Tumor necrosis factor- $\alpha$   
VEE Venezuelan equine encephalomyelitis  
WHO World health organization

✉ Ismail Celil Haskologlu  
ismailhaskologlu@gmail.com

Emine Erdag  
emine.erdag@neu.edu.tr

Serkan Sayiner  
serkan.sayiner@neu.edu.tr

Nurettin Abacioglu  
nurettin.abacioglu@neu.edu.tr

Ahmet Ozer Sehirli  
ahmetoz.sehirli@neu.edu.tr

<sup>1</sup> Faculty of Pharmacy, Department of Pharmacology, Near East University, Nicosia, Cyprus

<sup>2</sup> Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Near East University, Nicosia, Cyprus

<sup>3</sup> Faculty of Veterinary Medicine, Department of Biochemistry, Near East University, Nicosia, Cyprus

<sup>4</sup> Diagnostic Laboratory, Animal Hospital, Near East University, Nicosia, Cyprus

<sup>5</sup> Faculty of Dentistry, Department of Pharmacology, Near East University, Nicosia, Cyprus

## Background

Since the World Health Organization (WHO) reported the first case of COVID-19 on March 11, 2020 [1], previously identified dominant variants of SARS-CoV-2 virus were alpha (B.7), beta (B.351), gamma (P.1) and delta (B.617.2) [2]. The Omicron SARS-CoV-2 variant, first identified in South Africa in November 2021, is said to be more contagious than previous variants [3]. Omicron is known to have a much higher number of mutations compared to another previously dominant variant, Delta [4]. Fifteen of the mutations detected in the omicron variant are in the receptor-binding domain associated with increased viral binding affinity and antibody escape [5]. Therefore, understanding the mutational hotspots of the virus plays an important role in designing effective therapeutic and preventive strategies against the new variant. Variations in spike glycoprotein sequences and their implications suggest that vaccines and coronavirus-specific binding inhibitors and adjuvant treatment options may be required for the omicron variant of SARS-CoV-2 [3–5]. Vaccine-induced immunity aims to neutralize the SARS-CoV-2 spike protein and the receptor-interacting angiotensin converting enzyme 2 (ACE2) [6]. Since the spike protein in Omicron's version contains more modifications than other variants, it could be viewed as a potential anti-vaccine immune escape option. However, immunological escape from memory T cells is unlikely to occur for non-surface proteins, after infection or vaccine-induced immunity [7, 8].

On the other hand, recent data obtained during the pandemic period suggests that complications and deaths from COVID-19 disease caused by the SARS-CoV-2 virus might be related to the high viral load that infected people are exposed to [9]. However, clinical and experimental studies continue to develop new agents against COVID-19 from a prophylactic and therapeutic point of view. Monoclonal antibodies (mAb), which can bind to and neutralize the SARS-CoV-2 virus in infected patients, are one of the new classes of treatment being studied against COVID-19 [10, 11]. REGN-COV2 (imdevimab-casirivimab combination) is one of the neutralizing mAb treatments approved by the Food and Drug Administration (FDA) for emergency use in patients with mild to moderate complications of COVID-19 [12].

One of the most important conditions that characterizes a severe SARS-CoV-2 infection is the event known as a "cytokine storm" [13]. It is an aggressive inflammatory response with the release of large amounts of pro-inflammatory cytokines accompanied by COVID-19 infection [13]. This hyperinflammatory state is often characterized by pulmonary infiltration [13, 14]. With the entry of the SARS-CoV-2 virus into alveolar epithelial cells,

the immune system stimulates CD14 + and CD16 + monocytes, alveolar macrophages and Th17 lymphocytes to release large amounts of pro-inflammatory cytokines and chemokines, including interleukins (IL) such as IL-1 $\beta$ , IL-6 and IL-8, tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interferon- $\gamma$ -inducible protein 10 (IP10) [13, 14]. The resulting cytokine/chemokine storm causes severe damage to the lungs, endothelial and epithelial cells. Consequently, alveolar edema may develop with a breakdown in the integrity of the blood/air barrier [14]. Additionally, fibrinogen factors such as transforming growth factor beta (TGF- $\beta$ ) in the presence of a cytokine storm can significantly inhibit gas exchange in the lungs due to pulmonary fibrosis [14].

Despite all these treatment strategies, due to a severe cytokine storm, COVID-19 can have fatal outcomes such as tissue damage, lung failure and multi-organ failure [13, 14]. Complementary adjunctive and immunomodulatory therapies are required to reverse immune system dysfunction and cytokine dysregulation. Melatonin is also known to have anti-inflammatory [15], antioxidant [16], immunomodulatory [17] and antiviral [18] infectious activities. It may be useful to use melatonin together with REGN-COV2 in elderly and other high-risk patients as an adjuvant to vaccines against the Omicron variant of SARS-CoV-2 infection.


## REGN-COV2 therapy and omicron variant

Corona viruses are known as RNA viruses that cause zoonotic infections; they are particularly common in birds and mammals [19]. It is known that coronaviruses act by binding to the serine protease receptors TMPRSS2 and ACE2 on cells of the respiratory system due to the spike (S) proteins they carry [20]. Neutralizing monoclonal antibodies (mAb) used to fight the SARS-CoV-2 virus is produced by them against the receptor-binding domain of the virus spike (S) protein. Anti-RBD mAbs act by targeting the receptor-binding domain of protein S, preventing its binding to ACE2 on host cells and neutralizing its ability to fuse with the receptor [21], as shown in Fig. 1. Humanized mouse model technology or using sera from patients recovering from COVID-19 are known as preferred sources for obtaining neutralizing mAbs [22]. Concurrent studies continue to investigate the efficacy of neutralizing antibodies in hospitalized COVID-19 patients with severe symptoms [23, 24]. The REGN-COV2 antibody cocktail consists of a combination of casirivimab and imdevimab, two potent neutralizing IgG1 mAbs that bind to different sites of the spike protein receptor-binding domain [25]. The main reason for using neutralizing antibodies in combination rather than alone is to avoid simultaneous inactivation of both antibodies due to mutations in SARS-CoV-2 protein S. Furthermore, the

**Fig.1** REGN-COV2 (Casirivimab/imdevimab) antibody cocktail binds to the receptor binding domain (RBD) of SARS-CoV-2 virus and by neutralizing the RBD, the binding ability of the virus to ACE2 receptor is inhibited

**SARS-CoV-2**


**RBD- Antibody complex**



**REGN-COV2  
'antibody cocktail'  
Casirivimab/imdevimab**

**Antibodies block the RBD  
of spike protein,  
«neutralizing» its ability to  
bind ACE2**

**ACE2 receptor**



combination of casirivimab and imdevimab was tested on SARS-CoV-2-infected rhesus monkeys and golden hamsters, which served as models for SARS-CoV-2 disease, resulting in antibody-mediated cytotoxicity and cellular phagocytosis in infected cells [26]. In both models tested, there was evidence that prophylactic and therapeutic treatment with REGN-COV2 reduced viral load and lung disease severity compared to the placebo group [26].

In a recent study, it was found that the effect on REGN-COV2, also known as Regeneron mAb cocktail, is milder compared to other monoclonal antibodies [27]. In the same study, an artificial intelligence (AI) model was trained using tens of thousands of experimental data points. Based on the different three-dimensional structures of RBD-antibody complexes, we investigated how RBD mutations in the omicron variant might affect the activity of existing antibody drugs [27]. Looking at 15 known

RBD mutations (S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, N501Y, Y505H) in Omicron [4, 27] were the RBDACE2 complex and various RBD-antibody complexes also examined. The results showed that the Celltrion antibody regdanvimab could be significantly inactivated by Omicron RBD mutations E484A, Q493R and Q498R [27]. It has been shown that the E484A mutation can also greatly reduce the potency of Rockefeller University mAbs [27]. In contrast to Eli Lilly's mAb cocktail, it was clear that the antibody REGN10 (Imdevimab) did not compete directly with ACE2 and it was found that the Regeneron mAbs did not overlap with each other [27]. This appears to be crucial for REGN10933 (casirivimab) as it can directly neutralize the virus. Based on these results, omicron mutations may have a much milder effect on the REGN-COV2 cocktail compared to other mAb cocktails tested [27]. These results

support that REGN-COV2 can be used as a first option in monoclonal antibody-assisted therapies.

## Melatonin and importance of circadian rhythm in COVID-19

Chronotherapy could play an important role in the treatment of COVID-19 patients, a new concept in medicine that is gaining popularity today [28]. According to this concept, patients receive medication according to their circadian rhythm. In this way, as well as being able to achieve the same effectiveness at a lower dose, the medication administered can also reduce the side effects of other medications used at the same time. All these factors could optimize the therapeutic effect of the chronotherapy method. The SARS-CoV-2 virus uses endogenous cholesterol, for invasion of lung epithelial cells [29]. It is known that cholesterol production in the body peaks at night [30]. A recent study reported that statin drugs that decrease cholesterol biosynthesis by inhibiting HMGCoA reductase minimize the severe side effects of COVID-19 [31]. As a result of this study, it was observed that nighttime administration of the same dose of statins significantly reduced cholesterol levels in COVID-19 patients compared to the morning test group. Combined with all these findings, the same method could be applied to other drug groups that can be used in the treatment of SARS-CoV-2 infection and maximize the therapeutic effect.

The SARS-CoV-2 virus disrupts the function of circadian clock genes known as brain and muscle ARNT-like protein 1 (BMAL1) [32]. The decrease in the level of the BMAL1 gene, which is a regulator of the circadian rhythm, triggers the events leading to the cytokine storm by NF $\kappa$ B and causes the activation of pro-inflammatory cytokines [33]. Zhuang et al. demonstrated the relationship and role of circadian rhythms in altering the susceptibility of lung epithelial cells to SARS-CoV-2 infection [34]. According to this study, deletion of the essential circadian transcription activator BMAL1 resulted in lower expression of the primary viral receptor ACE2 and viral entry into lung epithelial cells.

Decreased BMAL1 levels lead to disruption of the renin–angiotensin–aldosterone system (RAAS) pathway and tissue damage [35]. BMAL1 deficiency also triggers increased production of pro-inflammatory cytokines such as TNF $\alpha$  [36]. Since elevated cytokine levels are important biomarkers that indicate the severity of coronavirus infection, it is important to develop a treatment strategy that takes into account BMAL1 and NF $\kappa$ B signaling in the management of COVID-19 disease [33, 36]. For the management of the NF $\kappa$ B signaling pathway, melatonin acts as a key molecule to antagonize the reduction in the levels of the Bmal1 gene, which induces a series of events leading to the cytokine storm [33]. The protective and anti-inflammatory effects of

melatonin make it a notable therapeutic option to enhance host defense response against viral invasive conditions in COVID-19.

Melatonin is a hormone produced by the pineal gland at night and plays an important role in the circadian rhythm [37]. It is known that melatonin reduces the hyperinflammatory response by inhibiting not only NF $\kappa$ B activity but also the interferon gamma response [38]. The fact that melatonin is readily available, inexpensive, and has a high safety profile makes it a suitable drug of choice for use against infections.


In addition to its anti-inflammatory effect, melatonin is known for its antidepressant, anxiolytic, immunomodulatory and antioxidant effects [39]. Another important role of melatonin is that it has a potent hydroxyl radical scavenging effect, particularly by increasing the activity of antioxidant enzymes such as glutathione in the antioxidant system [40]. In addition, various immune functions are modulated by melatonin. Exogenous administration of melatonin has been shown to increase antibody production [41]. Melatonin has stimulatory and suppressive effects on the immune system [42]. Specifically, it stimulates the immune system by increasing T cell activation and humoral response. In addition, it shows an immunosuppressive effect by reducing the activities of inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) enzymes, which play an important role in inflammation [43].

## The effects of melatonin in viral infections

Melatonin has indirect antiviral effects due to its anti-inflammatory, antioxidant, and immune-boosting properties [15]. Recent studies have shown that melatonin regulates levels of cytokines that play important roles in the immune system, such as interleukin 2 and IFN $\gamma$ , in mice during Venezuelan equine encephalomyelitis (VEE) virus infection [44]. As a result of the studies, it was found that melatonin reduces viral potency and also reduces acute lung damage in respiratory syncytial virus models by inhibiting oxidative damage. A recent study showed that interleukin 2 (IL2) and IFN $\gamma$  are specific biomarkers of the cellular response to SARS-CoV-2 [45]. Related to these findings, melatonin may reduce the effectiveness of the virus by showing similar activity in the SARS-CoV-2 virus.

SARS-CoV-2 has been reported to use the Cluster of differentiation 147 (CD147) S protein pathway to enter host cells [46]. CD147 is a glycoprotein that causes a cytokine storm and tissue damage in the lungs upon viral invasion. CD147 plays an important role in inflammation mediated by pro-inflammatory cytokines. In patients with COVID-19 infection, these cytokines are released uncontrollably, causing a cytokine storm [47]. Cytokine storm syndrome is caused by a variety of inflammatory conditions, including

**Fig.2** Melatonin prevents the cytokine storm and inflammation by inhibiting ROS production, overactivity of myeloperoxidase and excess cytokine release



severe systemic inflammation, hemodynamic instability, and multiple organ failure. With an age-related weakened immune system and declining melatonin levels, middle-aged and elderly patients with chronic diseases are much more susceptible to respiratory arrest from SARS-CoV-2 infection. In some clinical studies, melatonin has been shown to stimulate a large decrease in the levels of cytokines, IL-6, IFN- $\gamma$  and C-reactive protein, which have important effects on the CD147-mediated inflammatory pathway [48].

### Antioxidant and anti-inflammatory effects of melatonin in cytokine storm

As the SARS-CoV-2 virus develops cytokine storm and acute respiratory distress syndrome by binding to ACE2 receptors, the oxidative reactions that occur in cytokine storm syndrome cause reactive oxygen species (ROS)-mediated lung damage [49]. The immune system can become more susceptible to SARS-CoV-2 infection as it becomes compromised with age and major comorbidities such as diabetes, cancer, heart problems [50]. Factors implicated in the onset of COVID-19 infection include a particularly compromised immune response, pathogenicity of new viral variants, and uncontrolled production of ROS associated with a cytokine storm [49]. By using melatonin, reactive

oxygen species and the production of free metal ions can be significantly reduced. Therefore, harmful conditions such as DNA damage, protein oxidation and lipid peroxidation can also be prevented. Reactive oxygen species cause an increase in the expression of matrix metalloproteinases (MMPs) [51]. ROS removed by melatonin supplementation may reduce the deleterious effects of MMP overexpression. In addition, melatonin can protect against inflammation in the lungs resulting from COVID-19 by suppressing oxidative stress and cell apoptosis [52]. In previous studies it was reported that the antioxidant capacity of melatonin is higher compared to other known ROS scavengers such as lycopene, methionine, taurine and uric acid [49, 53]. In addition, the severity of the effects caused by the pro-inflammatory cytokines released in the cytokine storm following SARS-CoV-2 infection is related to the degree of inflammation. Myeloperoxidase (MPO) activity and ROS production have important effects that enhance the inflammatory immune response [54]. Inhibition of MPO and removal of unwanted ROS are important therapeutic targets against SARS-CoV-2 infection, as shown in Fig. 2.

Melatonin acts by inhibiting allosteric binding and chlorination activity at the entrance of the myeloperoxidase heme pocket [55]. The cytokine storm that develops due to COVID-19 causes the MPO enzyme to work overactive. MPO hyperactivity is one of the main sources of HOCl, one of the



important reactive oxygen species [56]. Therefore, melatonin plays a beneficial role in disease treatment by scavenging released HOCl or reducing ROS-induced metal release [49, 57]. In addition, melatonin, which plays an important role in ROS detoxification, can be considered a powerful adjuvant to combat COVID-19 infection.

In addition, when melatonin is used with other drugs in the treatment of COVID-19, it enhances the effects of other drugs and reduces the potential for their side effects [58]. Melatonin has been shown to be safe for short-term use, even at high doses [59]. Therefore, the use of melatonin can be suggested as a safe treatment modality along with other drugs against SARS-CoV-2 infection. the treatment of COVID-19 as an adjuvant therapy.

## Conclusion

While vaccines are the best strategy to prevent COVID-19, combination therapy of melatonin and REGN-CoV2 before or after exposure to the Omicron variant may offer significant benefits for elderly and chronically ill patients. Nursing ward admissions have increased due to the greater contagion of the Omicron variant. Therefore, infections caused by Omicron should not be underestimated and the importance of vaccination especially in high-risk patients should be emphasized. Additionally, when we consider the cost and time involved in developing a vaccine that targets new variants, identifying and expanding treatment options is extremely important. In conclusion, combined therapy of melatonin and REGN-CoV2 is an attractive approach with potential benefits in both prophylactic and therapeutic strategies as an adjuvant to vaccines for immune boosting and circadian rhythm regulation against the Omicron variant.

**Author contributions** All authors have equally contributed to the research and preparation of the article.

**Funding** No funding was received to assist with the preparation of this manuscript.

## Declarations

**Conflict of interest** The authors declare that they have no competing interests.

**Ethical approval** Not applicable.

**Consent for publication** Not applicable.

## References

1. Parasher A (2021) COVID-19: Current understanding of its pathophysiology, clinical presentation and treatment.

- Postgrad Med J 97:312–320. <https://doi.org/10.1136/postgradmedj-2020-138577>
2. Aleem A, Akbar Samad AB, Slenker AK (2022) Emerging Variants of SARS-CoV-2 And Novel Therapeutics Against Coronavirus (COVID-19). StatPearls Publishing, In StatPearls
  3. Wang L, Cheng G (2021) Sequence analysis of the emerging SARS-CoV-2 variant Omicron in South Africa. *J Med Virol* 94:1728–1733
  4. Kim S, Liu Y, Ziarnik M, Cao Y, Zhang XF, Im W (2022) Binding of Human ACE2 and RBD of Omicron Enhanced by Unique Interaction Patterns Among SARS-CoV-2 Variants of Concern. Preprint. bioRxiv: the preprint server for biology <https://doi.org/10.1101/2022.01.24.477633>
  5. Lupala CS, Ye Y, Chen H, Su XD, Liu H (2022) Mutations on RBD of SARS-CoV-2 omicron variant result in stronger binding to human ACE2 receptor. *Biochem Biophys Res Commun* 590:34–41. <https://doi.org/10.1016/j.bbrc.2021.12.079>
  6. Martínez-Flores D, Zepeda-Cervantes J, Cruz-Reséndiz A et al (2021) SARS-CoV-2 vaccines based on the spike glycoprotein and implications of new viral variants. *Front Immunol* 12:701501. <https://doi.org/10.3389/fimmu.2021.701501>
  7. Sekine T, Perez-Potti A, Rivera-Ballesteros O et al (2020) Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *Cell* 183:158–168.e14. <https://doi.org/10.1016/j.cell.2020.08.017>
  8. Grifoni A, Weiskopf D, Ramirez SI et al (2020) Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 181:1489–1501.e15. <https://doi.org/10.1016/j.cell.2020.05.015>
  9. Fajnzylber J, Regan J, Coxen K et al (2020) SARS-CoV-2 viral load is associated with increased disease severity and mortality. *Nat Commun* 11:5493. <https://doi.org/10.1038/s41467-020-19057-5>
  10. Kreuzberger N, Hirsch C, Chai KL et al (2021) SARS-CoV-2-neutralising monoclonal antibodies for treatment of COVID-19. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD013825.pub2>
  11. Liu LD, Lian C, Yeap LS, Meng FL (2020) The development of neutralizing antibodies against SARS-CoV-2 and their common features. *J Mol Cell Biol* 12:980–986. <https://doi.org/10.1093/jmcb/mjaa070>
  12. Heustess AM, Allard MA, Thompson DK, Fasinu PS (2021) Clinical management of COVID-19: a review of pharmacological treatment options. *Pharmaceuticals (Basel)* 14:520. <https://doi.org/10.3390/ph14060520>
  13. Ragab D, Salah Eldin H, Taeimah M, Khattab R, Salem R (2020) The COVID-19 cytokine storm; what we know so far. *Front Immunol* 11:1446. <https://doi.org/10.3389/fimmu.2020.01446>
  14. Kim JS, Lee JY, Yang JW et al (2021) Immunopathogenesis and treatment of cytokine storm in COVID-19. *Theranostics* 11:316–329. <https://doi.org/10.7150/thno.49713>
  15. Zhang R, Wang X, Ni L et al (2020) COVID-19: melatonin as a potential adjuvant treatment. *Life Sci* 250:117583. <https://doi.org/10.1016/j.lfs.2020.117583>
  16. Marseglia L, Gitto E, Laschi E et al (2021) Antioxidant effect of melatonin in preterm newborns. *Oxid Med Cell Longev* 2021:6308255. <https://doi.org/10.1155/2021/6308255>
  17. Álvarez-Sánchez N, Cruz-Chamorro I, López-González A et al (2015) Melatonin controls experimental autoimmune encephalomyelitis by altering the T effector/regulatory balance. *Brain Behav Immun* 50:101–114. <https://doi.org/10.1016/j.bbi.2015.06.021>
  18. Bahrapour Juybari K, Pourhanifeh MH, Hosseinzadeh A et al (2020) Melatonin potentials against viral infections including COVID-19: current evidence and new findings. *Virus Res* 287:198108. <https://doi.org/10.1016/j.virusres.2020.198108>

19. Wang W, Xu Y, Gao R et al (2020) Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA* 323:1843–1844. <https://doi.org/10.1001/jama.2020.3786>
20. Marchi R, Sugita B, Centa A et al (2021) The role of microRNAs in modulating SARS-CoV-2 infection in human cells: a systematic review. *Infect Genet Evol* 91:104832. <https://doi.org/10.1016/j.meegid.2021.104832>
21. Wang Z, Schmidt F, Weisblum Y et al (2021) mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *Nature* 592:616–622. <https://doi.org/10.1038/s41586-021-03324-6>
22. Hansen J, Baum A, Pascal KE et al (2020) Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science* 369:1010–1014. <https://doi.org/10.1126/science.abd0827>
23. Phan AT, Gukasyan J, Arabian S, Wang S, Neeki MM (2021) Emergent inpatient administration of casirivimab and imdevimab antibody cocktail for the treatment of COVID-19 pneumonia. *Cureus* 13:e15280. <https://doi.org/10.7759/cureus.15280>
24. O'Brien MP, Forleo-Neto E, Musser BJ et al (2021) Subcutaneous REGEN-COV antibody combination to prevent Covid-19. *N Engl J Med* 385:1184–1195. <https://doi.org/10.1056/NEJMoa2109682>
25. Taylor PC, Adams AC, Hufford MM et al (2021) Neutralizing monoclonal antibodies for treatment of COVID-19. *Nat Rev Immunol* 21:382–393. <https://doi.org/10.1038/s41577-021-00542-x>
26. Baum A, Ajithdoss D, Copin R et al (2020) REGN-COV2 antibodies prevent and treat SARS-CoV-2 infection in rhesus macaques and hamsters. *Science* 370:1110–1115. <https://doi.org/10.1126/science.abe2402>
27. Chen J, Wang R, Gilby NB, Wei GW (2022) Omicron variant (B.1.1.529): infectivity, vaccine breakthrough, and antibody resistance. *J Chem Inf Model* 62:412–422. <https://doi.org/10.1021/acs.jcim.1c01451>
28. Tamimi F, Abusamak M, Akkanti B et al (2020) The case for chronotherapy in Covid-19-induced acute respiratory distress syndrome. *Br J Pharmacol* 177:4845–4850. <https://doi.org/10.1111/bph.15140>
29. Kočar E, Režen T, Rozman D (2021) Cholesterol, lipoproteins, and COVID-19: basic concepts and clinical applications. *Biochim Biophys Acta Mol Cell Biol Lipids* 1866:158849. <https://doi.org/10.1016/j.bbalip.2020.158849>
30. Schroor MM, Sennels HP, Fahrenkrug J et al (2019) Diurnal variation of markers for cholesterol synthesis, cholesterol absorption, and bile acid synthesis: a systematic review and the bispebjerg study of diurnal variations. *Nutrients* 11:1439. <https://doi.org/10.3390/nu11071439>
31. Giri A, Srinivasan A, Sundar IK (2021) COVID-19: sleep, circadian rhythms and immunity - repurposing drugs and chronotherapeutics for SARS-CoV-2. *Front Neurosci* 15:674204. <https://doi.org/10.3389/fnins.2021.674204>
32. Borrmann H, McKeating JA, Zhuang X (2021) The circadian clock and viral infections. *J Biol Rhythms* 36:9–22. <https://doi.org/10.1177/0748730420967768>
33. Sehirli AO, Chukwunyeru U, Aksoy U, Sayiner S, Abacioglu N (2021) The circadian clock gene Bmal1: role in COVID-19 and periodontitis. *Chronobiol Int* 38:779–784. <https://doi.org/10.1080/07420528.2021.1895198>
34. Zhuang X, Tsukuda S, Wensch F et al (2021) The circadian clock component BMAL1 regulates SARS-CoV-2 entry and replication in lung epithelial cells. *iScience* 24:103144
35. Mercan M, Şehirli AO, Chukwunyeru U, Abacıoğlu N (2021) Acute kidney injury due to COVID-19 and the circadian rhythm. *Med Hypotheses* 146:110463. <https://doi.org/10.1016/j.mehy.2020.110463>
36. Vieira E, Mirizio GG, Barin GR, de Andrade RV et al (2020) Clock genes, inflammation and the immune system-implications for diabetes, obesity and neurodegenerative diseases. *Int J Mol Sci* 21:9743. <https://doi.org/10.3390/ijms21249743>
37. Grivas TB, Savvidou OD (2007) Melatonin the “light of night” in human biology and adolescent idiopathic scoliosis. *Scoliosis* 2:6. <https://doi.org/10.1186/1748-7161-2-6>
38. Yang L, Xie X, Tu Z, Fu J et al (2021) The signal pathways and treatment of cytokine storm in COVID-19. *Signal Transduct Target Ther* 6:255. <https://doi.org/10.1038/s41392-021-00679-0>
39. Emet M, Ozcan H, Ozel L et al (2016) A review of melatonin, its receptors and drugs. *Eurasian J Med* 48:135–141. <https://doi.org/10.5152/eurasianjmed.2015.0267>
40. Sehirli AO, Sayiner S, Serakinci N (2020) Role of melatonin in the treatment of COVID-19; as an adjuvant through cluster differentiation 147 (CD147). *Mol Biol Rep* 47:8229–8233. <https://doi.org/10.1007/s11033-020-05830-8>
41. Perflyeva YV, Ostapchuk YO, Abdolla N et al (2019) Exogenous melatonin up-regulates expression of CD62L by lymphocytes in aged mice under inflammatory and non-inflammatory conditions. *Immunol Invest* 48:632–643. <https://doi.org/10.1080/08820139.2019.1586918>
42. Srinivasan V, Maestroni GJ, Cardinali DP et al (2005) Melatonin, immune function and aging. *Immun Ageing* 2:17. <https://doi.org/10.1186/1742-4933-2-17>
43. Carrillo-Vico A, Lardone PJ, Álvarez-Sánchez N et al (2013) Melatonin: buffering the immune system. *Int J Mol Sci* 14:8638–8638. <https://doi.org/10.3390/ijms14048638>
44. Negrette B, Bonilla E, Valero N et al (2001) Melatonin treatment enhances the efficiency of mice immunization with venezuelan equine encephalomyelitis virus TC-83. *Neurochem Res* 26:767–770. <https://doi.org/10.1023/a:1011645400123>
45. Costela-Ruiz VJ, Illescas-Montes R, Puerta-Puerta JM et al (2020) SARS-CoV-2 infection: the role of cytokines in COVID-19 disease. *Cytokine Growth Factor Rev* 54:62–75. <https://doi.org/10.1016/j.cytogfr.2020.06.001>
46. Wang K, Chen W, Zhou Y-S et al (2020) SARS-CoV-2 invades host cells via a novel route: CD147-spike protein. *bioRxiv*. <https://doi.org/10.1101/2020.03.14.988345>
47. Mahmudpour M, Roozbeh J, Keshavarz M et al (2020) COVID-19 cytokine storm: the anger of inflammation. *Cytokine* 133:155151. <https://doi.org/10.1016/j.cyto.2020.155151>
48. Farias TSM, da de, Paixao RI, Cruz MM, et al (2019) Melatonin supplementation attenuates the pro-inflammatory adipokines expression in visceral fat from obese mice induced by a high-fat diet. *Cells* 8:1041. <https://doi.org/10.3390/cells8091041>
49. Camp OG, Bai D, Gonullu DC, Nayak N, Abu-Soud HM (2021) Melatonin interferes with COVID-19 at several distinct ROS-related steps. *J Inorg Biochem* 223:111546. <https://doi.org/10.1016/j.jinorgbio.2021.111546>
50. Williamson EJ, Walker AJ, Bhaskaran K et al (2020) Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 584:430–436. <https://doi.org/10.1038/s41586-020-2521-4>
51. Ueland T, Holter JC, Holten AR et al (2021) Distinct and early increase in circulating MMP-9 in COVID-19 patients with respiratory failure. *J Infect* 81:e41–e43. <https://doi.org/10.1016/j.jinf.2020.06.061>
52. Solun B, Shoenfeld Y (2020) Inhibition of metalloproteinases in therapy for severe lung injury due to COVID-19. *Med Drug Discov* 7:100052. <https://doi.org/10.1016/j.medidd.2020.100052>
53. Ogino T, Than TA, Hosako M et al (2009) Taurine chloramine: a possible oxidant reservoir. *Adv Exp Med Biol* 643:451–461. [https://doi.org/10.1007/978-0-387-75681-3\\_47](https://doi.org/10.1007/978-0-387-75681-3_47)
54. Björnsdóttir H, Welin A, Michaëlsson E et al (2015) Neutrophil NET formation is regulated from the inside by myeloperoxidase-processed reactive oxygen species. *Free Radic Biol Med* 89:1024–1035. <https://doi.org/10.1016/j.freeradbiomed.2015.10.398>

55. Galijasevic S, Abdulhamid I, Abu-Soud HM (2008) Melatonin is a potent inhibitor for myeloperoxidase. *Biochemistry* 47:2668–2677. <https://doi.org/10.1021/bi702016q>
56. Pullar JM, Vissers MC, Winterbourn CC (2000) Living with a killer: the effects of hypochlorous acid on mammalian cells. *IUBMB Life* 50:259–266. <https://doi.org/10.1080/713803731>
57. Banerjee J, Maitra D, Diamond MP, Abu-Soud HM (2012) Melatonin prevents hypochlorous acid-induced alterations in microtubule and chromosomal structure in metaphase-II mouse oocytes. *J Pineal Res* 53:122–128. <https://doi.org/10.1111/j.1600-079X.2012.00977.x>
58. Molina-Carballo A, Palacios-López R, Jerez-Calero A et al (2022) Protective effect of melatonin administration against SARS-CoV-2 infection: a systematic review. *Curr Issues Mol Biol* 44:31–45. <https://doi.org/10.3390/cimb44010003>
59. Andersen LP, Gögenur I, Rosenberg J, Reiter RJ (2016) The safety of melatonin in humans. *Clin Drug Investig* 36:169–175. <https://doi.org/10.1007/s40261-015-0368-5>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.